

L 29981-66 EMT(1) AT

ACC NR: AP6012482

SOURCE CODE: UR/0181/66/008/004/1182/1187-

65
64
B

AUTHOR: Gifeysman, Sh. N.

ORG: Kishinev State University (Kishinevskiy gosudarstvennyy universitet)

TITLE: Ionization energy of shallow traps in ionic crystals

SOURCE: Fizika tverdogo tela, v. 8, no. 4, 1966, 1182-1187

TOPIC TAGS: ionic crystal, ionization, impurity center, crystal lattice defect, phonon interaction, color center, polaron

ABSTRACT: This is a continuation of earlier work by the author (with Yu. Ye. Perlin, ZhETF v. 49, 1237, 1965), dealing with the ionization energy of impurity centers in ionic crystals. Unlike the earlier study, which was devoted to the case of a strong coupling between the localized electron and the lattice defect, the present study deals with weak coupling, and with arbitrary electro-phonon coupling. The calculation is based on the theory of shallow traps such as F centers, developed by Yu. Ye. Perlin, the author, and others (FTT v. 7, 1467, 1965; Uch. zap. Kishinevsk. gos. univ. v. 80, 3, 1965). The ionization energy is found to be equal to the Coulomb term of the polaron with effective mass corrected with allowance for the deformation of the polaron state by the field of the defects. The results are compared with the limiting case

Card

1/2

L 29981-66

ACC NR: AP6012482

of strong coupling with defects, such as crystals of the type A^{III}B^V and A^{II}B^{VI}. The ground-state energies obtained in the two limiting cases of strong and weak coupling of the localized electron with the lattice defect turn out to be quite close to each other. The only difference between the two couplings is that the external values lie in different regions. Comparison with experimental data by others shows good agreement for some crystals. The author thanks Yu. Ye. Perlin for continuous interest in the work and valuable remarks. Orig. art. has: 32 formulas and 2 tables.

SUB CODE: 20/ SUBM DATE: 17Sep65/ ORIG REF: 004/ OTH REF: 007

Card

2/2 *dc*

July/9

(A, N)

SOURCE CODE: UR/0181/66/008/011/3324/3334

AUTHOR: Perlin, Yu. Ye.; Gifcysman, Sh. N.

ORG: Kishinev State University (Kishinevskiy gosudarstvenny universitet)

TITLE: Diamagnetism of bound polarons

SOURCE: Fizika tverdogo tela, v. 8, no. 11, 1966, 3324-3334

TOPIC TAGS: polaron, diamagnetism, impurity center, ionic crystal, electron spin, magnetic susceptibility, ionization

ABSTRACT: This is a continuation of earlier work (FTT v. 7, 1467, 1965 and elsewhere) dealing with hydrogenlike local centers in ionic crystals. The present article extends the earlier results to include the case of a weak homogeneous external magnetic field and the case of a bound polaron. Inasmuch as the Lagrangian formalism of quantum mechanics has been developed only for systems having a classical Lagrangian, the present investigation is limited to diamagnetic effects of the localized electron which are not coupled to the spin. The partition function of an impurity semiconductor in a magnetic field is represented in the form of a Feynman functional integral over the trajectories of the localized electron. In the zeroth approximation, the system is described by a trial Lagrangian in which the Coulomb terms are imitated by elastic interactions. The dependence of the trial-Lagrangian parameters on the magnetic field is disregarded. In the first approximation in the difference between the true and trial action the authors calculate the free-energy correction which is

Card 1/2

GIGA, V., general-major; YURCHENKO, Ya., dotsent; KULIKOV, I., kand.-
pedagogicheskikh nauk

"Pedagogy; essays on the theory and practice of academic instruction
and training of Soviet soldiers" by A.G.Bazanov. Reviewed by
V.Giga, IA.IUrchenko, I.Kulikov. Voen. vest. 41 no.2:120-123
F '62. (MIRA 15:3)
(Russia--Army--Education, Nonmilitary) (Bazanov, A.G.)

GIGA, V., general-major; KALININ, V., podpolkovnik

Search for new methodological forms. Voen. vest. 42 no.8:77-79
Ag '62. (MIRA 15:7)
(Military education)

S/137/60/000/012/003/041
A006/A001

Translation from; Referativnyy zhurnal, Metallurgiya, 1960, No. 12, p. 77,
28600

AUTHORS: Ponomarev, V.D., Giganov, G.P.

TITLE: Separation of Tantalum and Niobium by Liquid Extraction

PERIODICAL: Izv. AN KazSSR Ser. metallurgii, obogashcheniya i ogneuporov, 1959,
No. 1 (4), pp. 3 - 15 (Kaz. summary)

TEXT: Laboratory investigations were made on extraction of Ta and Nb from a mineral acid mixture by tributylphosphate. It was established that Nb is more completely extracted into the organic phase at a concentration of HF 4 - 6 n. and H₂SO₄ 6 - 10 n. in the initial solution. At a Nb₂O₅ content in the initial solution as high as 30 - 35 g/l its extraction into the organic phase is 98%. Separation of Ta and Nb, if they are jointly present in the solution, can be performed by two ways: 1) in the absence of H₂SO₄ in the solution containing HF 0.5 - 2 n., mainly Ta is transferred into the organic phase, whereas Nb re-

V
Card 1/2

S/137/60/000/012/003/041
A006/A001

Separation of Tantalum and Niobium by Liquid Extraction

mains in the aqueous phase; 2) from the initial solution containing 6 n. HF and 8 n. H₂SO₄, Ta and Nb are jointly transferred into the organic phase, and Nb is then selectively washed out of the organic phase with 0.5 n. HF or pure water.

G.S.

Translator's note: This is the full translation of the original Russian abstract.

Card 2/2

S/137/61/000/001/003/043
A006/A001

Translation from: Referativnyy zhurnal, Metallurgiya, 1961, No. 1, p. 17,
1G158

AUTHORS: Giganov, O.P., Ponomarev, V.D., Khan, O.A.

TITLE: On the Composition of Niobium and Tantalum Complexes Extracted
With β -Butylphosphate

PERIODICAL: Izv. AN KazSSR Ser. metallurgii, obogashcheniya i ogneuporov, 1960,
No. 3 (6), pp. 73 - 78 (Kaz. summary)

TEXT: The authors describe a graphical method of determining the composition of Ta and Nb complexes extracted from a mixture of H_2SO_4 and HF acids with the aid of tributylphosphate. An investigation of the distribution coefficient of each of the metals and acids depending on the amount of tributylphosphate in the organic phase (kerosene) has shown that the acids are extracted in the form of complexes: H_2SO_4 . tributylphosphate and HF . tributylphosphate. Ta apparently forms a dissolvate H_2TaP_7 . 2-tributylphosphate, and Nb a complex H_2NbF_7 . tributylphosphate. Highest distribution coefficients are obtained for

Card 1/2

S/137/61/000/001/003/043
A006/A001

On the Composition of Niobium and Tantalum Complexes Extracted With 3-Butylphosphate

Ta in the presence of H_2SO_4 in the initial solution and minimum HF concentration; for Nb in the presence of H_2SO_4 and HF concentration > 7%. At a low HF concentration niobium is present in the solution in the form of a non-extractable complex $H_2NbOF_5 \cdot H_2O$. ✓

M. L.

Translator's note: This is the full translation of the original Russian abstract.

Card 2/2

S/828/62/000/000/007/017
E039/E420

AUTHORS: Giganov, G.P., Ponomarev, V.D., Khan, O.A.

TITLE: On the conditions for the extraction and separation of tantalum and niobium and the formation of complexes

SOURCE: Razdeleniye blizkikh po svoystvam redkikh metallov.
Mezhvuz. konfer. po metodam razdel. blizkikh po svoyst.
red. metallov. Moscow, Metallurgizdat, 1962, 79-97

TEXT: As no previous work on this part of the subject has been published the authors present results of an investigation on the formation of complexes with HF, H₂SO₄, Ta and Nb and on the conditions for extraction of the separate metals. The experiments are carried out at 20°C in polyethylene vessels and the phase separation accomplished in a graduated polyethylene funnel. The duration of mixing is 10 minutes and the time of separation ~ 1 hour. The initial ratio of phases is 1:1. Methods of analysis are discussed in detail and the influence of various parameters on the extraction of Ta and Nb are examined. In particular the dependence of the distribution coefficient K_p and the specific electrical conductivity of the organic phase on Card 1/2

S/828/62/CC/CC/CC7/317
E039/E420

On the conditions for ...

the concentration of HF, the concentration of the metals in solution and the concentration of tributylphosphate (TBP) is examined. It is determined that in the presence of 3M H₂SO₄ in an aqueous solution of HF the limiting concentration of Nb and/or Ta in undiluted TBP is 1M or 133 g/litre of Nb₂O₅ and 225 g/litre of Ta₂O₅. Niobium is extracted from HF solution (with or without H₂SO₄) by TBP in the form of HNbF₆·3TBP. From a saturated solution of Nb containing a solid phase Nb is transferred to the ether phase in the form of oxyfluoride complexes H₂NbO₅·3TBP and HNbOF₄·3TBP. At low concentrations of HF in aqueous solution Ta is extracted as HTaF₆·3TBP and at high concentrations of HF and from solutions containing H₂SO₄ the Ta is transferred in the form H₂TaF₇·3TBP. The optimum conditions for separation of Nb and Ta by extraction with TBP from HF-H₂SO₄ solutions are studied. If in the initial solution the ratio Ta₂O₅:Nb₂O₅ = 2 or more, the largest separation coefficient is obtained by extraction from a weak acid solution 1M HF - 0.5M H₂SO₄. When the ratio of Nb₂O₅:Ta₂O₅ = 2 or more in the initial solution, it is necessary to extract from a solution with excess acid by 6M HF - 3M H₂SO₄. There are 9 figures and 5 tables.

S/817/62/005/000/007/012
A006/A101

AUTHORS: Giganov, G. P., Potomarev, V. D.

TITLE: The extraction of hydrofluoric acid with tributylphosphate (TBP)

SOURCE: Akademiya nauk Kazakhskoy SSR. Institut metallurgii i obogashcheniya. Trudy. v. 5, 1962, Tsvetnaya metallurgiya, 108 - 114

TEXT: To investigate complex formation in the system HF-TBP the authors studied the extraction of hydrofluoric acid from aqueous solutions with concentrations varying from 1 to 20 mole. After one-hour separation of phases, specific electric resistivity of the organic phase was measured at 21 and 41°C. Equilibrium concentration of HF was determined by titration at 21 and 41°C. coefficient was calculated. The results obtained show that hydrofluoric acid is extracted with TBP in the form of monosolvate from aqueous solutions from $\frac{1}{4}$ to 14 mole, and for all the investigated HF concentrations in aqueous solutions in the extrahent. The shape of the curves of specific electric conductivity and of the temperature coefficient of electric conductivity indicates the formation of a new complex at 4 mole HF concentration in the equilibrium aqueous

Th
pha
of 1
the
basis:
repre
aqueo
are 6

in the
concentrations. There

Card 1/2

S/317/62/005/000/003/012
A006/A101

AUTHORS: Giganov, G. P., Ponomarev, V. D.

TITLE: Sulfuric acid extraction with tributylphosphate (TBP)

SOURCE: Akademiya nauk Kazakhskoy SSR. Institut metallurgii i obogashcheniya. Trudy, v. 5, 1962, Tsvetnaya metallurgiya, 115 - 118

TEXT: The authors studied sulfuric acid extraction in a concentration range of from 1 to 5.5 mole. The concentration of the acid in equilibrium phases was determined by titration with caustic soda from methyl orange. The density of the organic phase was determined pycnometrically, and viscosity was measured with a capillary viscosimeter. The results are given in graphs. To determine the degree of solvation of the sulfuric acid in the organic phase, the experimental data were used to calculate the logarithmic dependence of distribution upon equilibrium concentration of TBP. The data obtained lead to the conclusion that at up to 3 mole concentration in the aqueous phase, the sulfuric acid is extracted in the form of complex $H_2SO_4 \cdot TBP \cdot 2H_2$. From higher-concentrated solu-

Card 1/2

S/617/62/005/000/008/012

Sulfuric acid extraction with tributylphosphate (TBP) A006/A101

tions the sulfuric acid is transferred to the organic phase in the form of compound $H_2SO_4 \cdot TBP \cdot H_2O$. The formation of a new chemical compound at 3 mole H_2SO_4 concentration in the aqueous phase is indirectly proved by a sharp increase in electric conductivity of the organic phase, a maximum on the curve of the temperature coefficient of electric conductivity and a minimum on the curve of the relative temperature coefficient of viscosity. This is explained by a lesser degree of hydration of the extracted complex and a higher degree of its dissociation. On the basis of the different hydration degree of the sulfuric acid, it can be assumed that the degree of its solvation with tributylphosphate will also be different. To extract a higher-hydrated molecule, two ester molecules are required; in case of lesser hydration one ester molecule is needed. There are 5 figures and 2 tables.

✓
—

Card 2/2

S/817/62/005/000/009/012
A006/A101

AUTHORS: Giganov, G. P., Ponomarev, V. D.

TITLE: Niobium extraction with tributylphosphate (TBP)

SOURCE: Akademiya nauk Kazakhskoy SSR. Institut metallurgii i obogashcheniya. Trudy. v. 5, 1962, Tsvetnaya metallurgiya, 119 - 124

TEXT: Graphoanalytical and physico-chemical methods of measuring the electric conductivity of the organic phase were employed to investigate complex formation in systems HF-Nb₂O₅-TBP and HF-H₂SO₄-Nb₂O₅-TBP. The authors studied niobium extraction depending on the concentration of hydrofluoric acid, on the metal concentration in the solution, and on tributylphosphate concentration in the extractant. At relatively low Nb concentrations in the initial solution the composition of the complex to be extracted was determined in three series of experiments using solutions with different concentrations of hydrofluoric and sulfuric acids, and equal Nb content. The experimental investigation yielded the following results. It was established that in the presence of 3 mole sulfuric acid in the aqueous solution, the extremal niobium concentration in undiluted

Card 1/2

Niobium extraction with tributylphosphate (TBP)

S/817/62/005/000/009/012
A006/A101

tributylphosphate is 1 mole/l or 133 g/l Nb_2O_5 . In the absence of sulfuric acid in the aqueous solution, the extremal Nb content in the extractant can only attain 0.74 mole (98 g/l Nb_2O_5). Out of hydrofluoric acid solutions, with and without sulfuric acid, niobium is extracted with tributylphosphate in the form of a trisolvate ($\text{HNbF}_6 \cdot 3\text{TBP}$). From niobium-saturated solutions with a solid phase, niobium is transferred to the ester phase in the form of oxifluoride complexes $\text{H}_2\text{NbOF}_5 \cdot 3\text{TBP}$, $\text{HNbOF}_4 \cdot 3\text{TBP}$. There are 2 tables and 3 figures.

Card 2/2

S/817/62/005/000/010/012
A000/A101

AUTHORS: Diganov, G. P., Ponomarev, V. D.

TITLE: Tantalum extraction with tributylphosphate (TBP)

SOURCE: Akademiya nauk Kazakhskoy SSR. Institut metallurgii i obogashcheniya. Trudy. v. 5, 1962, Tsvetnaya metallurgiya, 125 - 129

TEXT: To investigate tantalum extraction, the gravimetric method was used for large tantalum amounts and the colorimetric method for smaller quantities. The authors investigated tantalum extraction, depending on the concentration of hydrofluoric acid, the concentration of the extractant, the concentration in the solution, and the composition of metal in the solution, and TBP were in g/l: HF - 4; Ta - 0.113; HF - 4; H₂SO₄ - 3; Ta - 0.244; HF - 12; Ta - 0.2; HF - 16; Ta - 0.2. TBP concentration varied between 3.3 to 0.73 mole/liter. It was found that the limit concentration of tantalum in the tributylphosphate, in the presence of 3 mole/liter sulfuric acid in the initial solution, is 1 mole/liter (225 g/l Ta₂O₅). Without sulfuric acid in the initial

1/2

Tantalum extraction with tributylphosphate (TBP)

S/817/62/000/000/010/012
A006/A101

talum content in the extractant is 0.81 mole/liter (1.9 g/l Ta₂O₅). It was es-
tablished that tantalum was transferred to the organic phase in the form of
solvates: HTaF₆.3TBP and H₂TaF₇.3TBP. There are 3 figures and 2 tables.

Card 2/2

GIGANOV, G.P.; PONOMAREV, V.D.; KHAN, O.A.

Conditions for the extractive separation and composition of tantalum and niobium complex ores. Trudy Alt. GMNII AN Kazakh. SSR 14:
39-51 '63. (MIRA 16:9)
(Tantalum--Metallurgy) (Niobium--Metallurgy)

GIGAS, E.

Klectronic Measurement of Distances. Geodetski list, Zagreb (Federation of
Geodetic Engineers and Geometers), p. 520, N. 7, No 11/12, Nov/Dec 1953, p. 3,
vol. 8, No 1/4, Ja/Ap 1954.

GIOVAS, F.

Measuring distances with an interferometer.

P. 113 (Geodezia es Kartografla. Vol 9, no 3, 1957, Budapest, Hungary)

Monthly Index of East European Accessions (FFAJ) LC. Vol. 7, no. 2,
February 1958

MIRZASHVILI, V.I.; GIGAURI, G.N.

Characteristics of the growth of spruce stands in Georgia and their
optimal cutting age. Trudy Inst. leza AN Gruz. SSR 10:15-29 '62.
(MIRA 17:3)

CIGAURI, G.N.; MIRZASHVILI, V.I.

Characteristics of the growth of pine stands in Georgia and their
optimal cutting age. Trudy Inst. lesa AN Gruz. SSR 10:31-42 '62.
(MIRA 17:3)

MIRZASHVILI, V.I.; GIGAURI, G.N.

Nature of growth of the fir woods of Georgia and the optimal age
for their cutting. Trudy Inst. lesa AN Gruz. SSR 12:3-18 '63.
(MIRA 18:2)

GIGAURI, G.N.

Nature of growth and taxonomic structure of *Opuntia hastanea*
sativa Mill. (cactus). Trudy Inst. Fiz. Akad. Nauk Gruz. SSR 12:93-102
1963.

GIGAUBI, V.S.; KOLYUTSKAYA, G.D.; OVCHINNIKOV, G.N.; STEPAN'YAN, Yu.L.;
GEBEL', G.Ya.; NIKOLAYEVA-KOMBERG, M.I.

Use of the RO-1 apparatus in the clinic of the Scientific Research
Institute of Clinical and Experimental Surgery of the Ministry of
Public Health of the R.S.F.S.R. Report No.2. Nov. med. tekhn. no.3:
58-60 '65. (MIRA 19:1)

GIGAURI, V.S.

Histochemical study of liver protein following sensitization. Zhur. mikrobiol., epid. i immun. 32 no.9:74-77 S '61. (MLRA 15:2)

1. Iz kafedry patologicheskoy fiziologii I Moskovskogo ordena Lenina meditsinskogo instituta imeni Sechenova,
(LIVER) (PROTEINS) (ANAPHYLAXIS)

GIGAURI, V.S.

Producing blood serum without hemolysis. Lab. delo 2 no.4:46-47
Ap '62. (MIR 15:5)

1. Kafedra patologicheskoy fiziologii (zav. - prof. S.M.Pavlenko)
I Moskovskogo ordena Lenina meditsinskogo instituta imeni I.M.Sechenova.
(SERUM) (HEMOLYSIS AND HEMOLYSINS)

GIGAURI, V.S.; YURIN, R.F. (Moskva)

Blood coagulation system and its disorders; a teaching schedule.
Pat. fiziol. i eksp. terap. 7 no.4:72-73 Jl-Ag '63.

(MIRA 17:9)

1. Iz kafedry patologicheskoy fiziologii (zav.- zasluzhennyy
deyatel' nauki RSFSR prof. S.M. Pavlenko) I Moskovskogo ordena
Lenina meditsinskogo instituta imeni Sechenova.

GIGAURI, V.S.; LIUSHITS, Ye.V.; TREKOVA, N.A.

Effect of muscle relaxants on the cardiovascular system. Trudy
I-^{go} MMI 33:41-47 '64. (MIRA 18:3)

ABINDELL, RICHARD, JR.

Accompanying document and interpretation, related to the subject, attached
below, on the hypothesis that resistance efforts to prevent further
expansion of Soviet influence in Central America were
unproductive. Study is to be completed by [redacted]

GIGAURI, V.S.

Dynamics of the distribution of radioactive phosphorus P^{32} in
the organism of rats in a state of hypothermia. Trudy 1-go MM
(MIRA 18:3)
33:120-123 '64.

GIGAURI, V.S.; OVCHINNIKOV, G.S.; MURDASOVA, I.V.

Clinical and experimental experience in the use of the RN-59
respiratory apparatus. Trudy 1-go MMI 33:226-231 '64.

(MIRA 18:3)

GIGAURI, V.S.

Protein function of the liver in sensitization. Izv. Akademii, epid. i immun. 42 no.1:27-31 Ja '65. (XAI 3²;6)

I. I. Moskovskiy ordena Lenina meditsinskij institut im. I.M. Sechenova.

GIMURI, Yu. A.

"The Problem of the Selection and Application of the "Best" Nutritive Substances of Yams or Sweet Potato. Nutritive Value of the Yam and its Nutritional Value," Sov. Med. Sci., Voil'naia Med. Rev. i Tret'iaia, 1951, 1953. (ZMZh, No. 1, Sep 53)

SO: Sov. 432, N° 1ar 55

USSR/ Chemistry - Coal	
Orgnol/1	Pub. 116 - 25/39
Authors	Kuznetsov, V. I.; Devorova, R. P.; Fadeycheva, A. G.; <u>Gigel', T. B.</u> ; and Chernysh, N. K.
Title	Complex interaction of brown coal in the Ukr. SSR. Part 13, Tars from semicoking of brown coal with the solid heat carrier - semicoke.
Periodical	Ukr. Khim. zhurn. 21/6, 804-809, Dec 1955
Abstract	Tars obtained by semicoking of brown coal with the solid heat-carrier (semicoke) were found to offer a higher yield of benzene and lower yield of paraffins than tar obtained during the semicoking of the very fine fractions as compared with tar obtained with a gaseous heat carrier. The primary decomposition products of brown coal with a gaseous heat carrier - semicoke - during the semicoking of brown coal with a solid heat carrier - semicoke submit to cracking to a greater extent than during semicoking with a gaseous heat carrier. The increase in fractions in tars of unsaturated compounds was found to be due to cracking. The phenols obtained from such fractions offer a somewhat lower yield of phenol-cresol fractions; and the paraffin yield is much lower. Tables; graph.
Institution	Acad. of Sc., Ukr. SSR, Inst. of Heat Power Engineering, Lab. for Chem. Proc.
Submitted	June 17, 1955

GIGIASHVILI, B. F.

4689 Gigiashvili, B. F. kak mekhanizirovat' vodoshabzhyeniye
vashivotnovodstva. tbitlisi. izd-vo gruz s-kh. in-ta, 1954. 12 s. s ill 20 sm. (upr.
s-kh propagandy m-vasovkhozov gruz. ssr) 2,000 ekz bespl.-na obl. avt. ne
ukazany.--na gruz yaz.-154-567511 636,0025/628.18:636

GIGIASHVILI, B. F.

4690 Gigiashvili, B. F. kompleksnaya mekhanizatsiya zhivotnovodcheskikh
farm. tbilisi, izd-vo gruz. s-kh in-ta, 1954. 8 s. szill 20sm. (upr. s. kh. propaganda
m-va sovkhозov gruz ssr) 2,000 ekz. Bespl-naobl avt ne ukazano

124 57-2-2098 D

Translation from: Referativnyy zhurnal. Mekhanika 1957. Nr 2. p 89 (USSR)

AUTHOR: Gigiberiya, G. Ya.

TITLE: Investigation of the Dynamics of Air Inclusions in Hydraulic
High pressure Structures (Issledovaniye dinamiki vozdushnykh
vkluchenii v napornykh gidrotehnicheskikh sooruzheniyakh)

ABSTRACT: Bibliographic entry on the author's dissertation for the degree
of Candidate of Technical Sciences presented to the Azerb.
industr. in-t (Azerbaijhan Industrial Institute). Tbilisi 1956

ASSOCIATION: Azerb. industr. in-t (Azerbaijhan Industrial Institute),
Tbilisi

1. Structures--Analysis 2. Air--Properties 3. Dynamics--Theory

Card 1/1

SOV/124-58-1-874

Translation from: Referativnyy zhurnal, Mekhanika, 1958, Nr 1, p 117 (USSR)

AUTHOR: Gigiberiya, G. Ya.

TITLE: Problems of the Hydraulics of Air Inclusions (Voprosy gidravliki vozdushnykh vkljucheniy)

PERIODICAL: Tr. In-ta energ. AN GruzSSR, 1956, Vol 10, pp 67-78

ABSTRACT: The author obtains the rate of displacement of an air inclusion in a sloping high-pressure conduit through an application of Hamilton's principle of least action. He treats the drag force of the air inclusion relative to the water flow erroneously as a quantity of motion, whereas that velocity can be obtained simply by equating that drag force to the projection on the axis of the conduit of the lifting force. As a result of this error the author's formula, in its initial form, contains an incorrect numerical coefficient; this, however, does not affect the further reasonings, since this error is automatically corrected during the computation of the parameters of the definitive formulas from the test data. The experimental material relates to pipes 54 and 45 mm in diameter. The lack of rigorousness of the theoretical reasonings (e.g., the determination of the length of an air inclusion by dividing

Card 1/2

Problems of the Hydraulics of Air Inclusions

SOV/124-58-1-874

its volume by its middle section) and the narrowness of the range covered by the test data render the use of the results obtained for other conditions difficult.
Bibliography: 6 references.

N. A. Kartvelishvili

Card 2/2

SOV/124-58-4-4100

Translation from: Referativnyy zhurnal, Mekhanika, 1958, Nr 4, p 60 (USSR)

AUTHOR: Gigiberiya, G. Ya.

TITLE: Water Hammer During the Escape of Air Inclusions From a High-pressure Water Conduit (Gidravlicheskiy udar pri vykhode vozдушных включений из напорного водовода)

PERIODICAL: Tr. In-ta energ. AN GruzSSR, 1956, Vol 10, pp 195-203

ABSTRACT: On the basis of his previous study (Tr. In-ta energ. AN GruzSSR, 1956, Vol 10, pp 67-68; RZhMekh, 1958, Nr 1, abstract 874), the author suggests a method for the determination of the law of fluctuation of the discharge of water in a section of a high-pressure main connected to a sufficiently large reservoir during the escape from the conduit of entrained air inclusions. This law is a boundary condition for the calculation of the water hammer which may be obtained by any of the known methods. In the example given by him, the author uses the familiar method of M. A. Mostkov.

1. Pipelines--Pressure 2. Air--Performance
3. Mathematics

N. A. Kartvelishvili

Card 1/1

GIGIBERIYA, G.Ya.

Steady-state flow of air and water in siphon. Trudy Inst.energ.
AN Gruz.SSR 16:183-192 '62. (MIRA 16:4)
(Fluid dynamics) (Siphons)

GIGIBERIY¹, SH. S.

Gigiberiya, Sh. S.: "Ways of improving the quality of the tea leaf in machine harvesting" Byulleten' Vsesoyuz. nauch.-issled. in-ta chaya i subtrop. kul'tur, 1948, No. 3, p. 95-101.

SO: U-3042, 11 March 53, (Letopis 'nykh Statey, No. 10, 1949).

"APPROVED FOR RELEASE: 09/24/2001

CIA-RDP86-00513R000515020015-2

KHAPARDEEV, Sh.; GIGBERTYA, V.

Rustavi Metallurgical Plant. Metallurg 9 no.11:2 -31 0 164.
(MIRA 18:2)

APPROVED FOR RELEASE: 09/24/2001

CIA-RDP86-00513R000515020015-2"

GIGIN, I.

Obsolete terminology. Mol. prom. 13, No 6, 1952.

"APPROVED FOR RELEASE: 09/24/2001

CIA-RDP86-00513R000515020015-2

GIGINEYSHVILI, G.M.

Vocal audiometry with the use of Georgian words. Trudy Tbil.
(MIRA 16:2)
GIDUV 6:343-348 '62.
(AUDIOMETRY)

APPROVED FOR RELEASE: 09/24/2001

CIA-RDP86-00513R000515020015-2"

GIGDZEYSHVILI, G.M., aspirant

Objective assessment of speech audimetry. Vestn. otorinolaring.
25 no.3 t1'7-19 '63 (MIRA 17 t1)

1. Iz kliniki bolezney ukha, gorla i nosa (zav. - prof. S.M. Khechinashvili) Tbilisskogo instituta usovershenstvovaniya vrachey.

GIGINEISHVILI, G.N.

Discharge conditions of the Kelasuri and Besleti Rivers. Trudy
Inst. geog. AN Gruz. SSR 17:201-211 '62. (MIRA 16:7)

(Kelasuri River—Hydrology)
(Besleti River—Hydrology)

VLADIMIROV, I.A.; MELIKISHVILI, T.S.; GIGIMYREVILI, G.N.

Hypsometry of river basins in Georgia. Soob. AN Gruz. SSR 28 no.4:
L25-430 A₁ 1:2.

(MIREA 18:1)

i. AN Gruzinskoy SSR, Institut geografii im. Vakhtangi, Tbilisi.
Submitted January 0, 1970.

GIGINEYSHVILI, G.N.

Regime of rivers and springs in the karst zone of Abkhazia
from Gumitsa to Madzhara. Trudy Inst. geog. AN Gruz. SSR
(MIRA 17:6)
18:243-244 '64.

GIGINEYSHVILI, G.N.

Average perennial drainage and its distribution within the year
in the small rivers of Abkhazia. Trudy Inst. geog. AN Gruz. SSR
(MIRA 18:5)
20:227-234 '64.

KASHAKASHVILI, N.V., prof., oty.red.; GAMBASHIDZE, R.B., kand.nauk, oty.
red.; AGLADZE, R.I., prof., red.; BERIDZE, V.M., prof., red.;
GIGINEISHVILI, K.M., red.; GONIASHVILI, T.B., kand.nauk, red.;
TAVADZE, F.I., prof., red.; KKEKLIDZE, M.A., doktor nauk, red.;
MIKELADZE, G.Sh., kand.nauk, red.; NADIRADZE, Ye.M., kand.nauk,
red.¶

[Metallurgical terminology] Metallurgicheskaja terminologija.
Otv.red. N.V.Kashakashvili i R.B.Gambashidze. Tbilisi, 1959.
(MIRA 13:2)
324 p.

1. Akademija nauk Gruzinskoy SSR, Tiflis. Institut jazykoznanija.
(Metallurgy--Dictionaries)
(Russian language--Dictionaries--Georgian)
(Georgian language--Dictionaries--Russian)

AMERICAN, U. S.

AMERICAN INDEPENDENT, U. S. Corporation "American Independent" 1973
(Regd. No. 100-1000), D. V., 1973, C. 1973, Inc., 1973, -
expiring Mar. 1978.

AMERICAN, U. S. 1973

GIGINEYSHVILI, M.S.

1. GIGINEYSHVILI, M., METRASHVILI, N.V.
2. USSR (600)
4. Tuberculosis
7. Atypical forms of acute military tuberculosis of the lung, Probl. tub. no. 6
1952.
9. Monthly List of Russian Accessions, Library of Congress, April 1953, Uncl.

KETSKYAN, S.P.; KIRBLADZE, D.Ch.; GIGRELLIYA, G.N.

qualitative composition of phytochromes of bark and wood plastids.
Zhob. Akad. Nauk Gruz. SSR 36 no.1:161-168 (1964).
(MIRA 18:3)

1. Institut Botaniki Akademii Nauk Gruzinsskoy SSR. Submitted March 23, 1964.

GIGINEYSHVILI, N.S.

Structure of intramural innervating mechanisms of the uterus in
animals before pregnancy, at various stages in pregnancy, and
during and after labor. Soob.AN Gruz.SSR 24 no.4:485-492 Ap
'60. (MIRA 13:7)

1. Tbilisskiy gosudarstvennyy meditsinskiy institut. Predstavлено
академиком V.K.Zhgenti.
(UTERUS-- INNERVATION)

GIGINEVILY, N. S.

Giginevily, N. S. "On 'specific genes' and the constitution
of gray karakul sheep," Karakulievodstvo i zverovedstvo, 1949, No. 2,
p. 21-26.

SS: 6-3736, 21 May 53, (Letopis 'Zhurnal 'nyki Statey, No. 17, 1949).

GIGINEYSHVILI, N. S.

Early determination of vitality in grey karakul lambs. Kar. i zver., 5, No 1,
1952.

1. GIGIKH-S.VILLI, N. A.
2. USSR (600)
4. Karakul Sheep
7. Developing a methodology for selecting grey karakul sheep with the aim of increasing the vitality of the offspring. Trudy Inst. morf. zhiv. no 7'52.
)
9. Monthly List of Russian Accessions, Library of Congress, March 1953, Unclassified.

1. GIGINEYSHVILI, N. S.
2. TCSR (600)
4. Karakul Sheep
7. Production standards for karakul fur need to be thoroughly revised. Kar. i zver
6 no. 1, 1953
9. Monthly List of Russian Accessions, Library of Congress, June 1953, Uncl.

GIGINEVSHVILI, N. S.

Gray karakul sheep Moskva, Gos. izd-vo sel'khoz. lit-ry, 1954. (Mic 55-3591)

Collation of the original, as determined from the film: 319 p.

Microfilm Slavic 387 AC

USSR/Farm Animals. Sheep and Goats.

Q

Abs Jour: Ref Zhur-Biol., No 17, 1958, 70761.

Author : Gidneyashvili, N. S.

Inst.

Title : Results of Selection Work for the Creation of
New hues of "Sur" Karakul Lambskin.

Orig Pub: Karakulevodstvo i zverovodstvo, 1956, No 2,
13-17.

Abstract: In breeding for a series of years of colored Karakul
sheep of the "kambar" type with selection of rams
which possessed lambskins of heterogeneous color
of the fibers in length (growing lighter toward the
ends), with the use of related breeding, permitted
the obtaining of Karakul sheep which gave lambs
with new colorings of the skins. The greatest

Card : 1/2

GIGINEYSHVILI, N. S.

USER / Farm Animals. Small Horned Stock.

Q

Abs Jour: Ref Zhur-Biol., No 9, 1958, 40458.

Author : Gigineyshvili, N. S.

Inst : Not given.

Title : Hybridization in Karakul Breeding.

Orig Pub: Karakulevodstvo i zverovodstvo, 1957, No 4,
14-20.

Abstract: In order to increase the vitality of the gray Karakul sheep, an experiment in hybridization was started in 1953 with Argali Ovis ammon bochariensis Nasonov. The aim of the experiment was to obtain hybrids with 1/8 of Argali blood and 7/8 of Karakul ewes' blood, uniting the valuable fur qualities of the gray Karakuls with an increased constitutional Argali toughness. The adult Argali males were hunted and

Card 1/3

USSR / Farm Animals. Small Horned Stock.

Q

Abs Jour: Ref Zhur-Biol., No 9, 1958, 40458.

Abstract: shot, and their spermatic cords were amputated and sent to the sovkhoz Ak-Kapchigay (Uzbekistan), where the sperm was extracted for the artificial insemination of gray Karakul ewes. In all, during 1953 - 1956, 300 ewes were inseminated, of which 37 hybrids F₁ were born, including 6 stillborn ones; 9 died, 15 were sacrificed for research, and 7 remained alive. In 1955, 170 gray Karakul ewes and 22 black ones were inseminated by the young hybrid ram Kok-Kiyik. 100 ewes (62.1%) brought forth lambs. The offspring were obtained also from two hybrid ewes inseminated by the Karakul rams. Of 34 lambs F₁, 14 had gray coloration, 16 had black coloration, 3 were "guligaz" (mixture of white and gray hair), and one was of brown coloration. In F₂, from uniform (by color) mating, 22 black

Card 2/3

34

OIGINEYSHVILI, N. A., BUTARIN, N. S., LOGINOV, N. V., LOPYRIN, A. I.,
RAKHIMOV, A. A.

"An attempt making creative use of the Mitshurin method of remote
hybridization for the purpose of the selection of agricultural animals."

reported at Conference on Problem of Heredity and Variability, held at
Institute of Genetics, AS USSR, 8-14 Oct 1957
Vestnik AN SSSR, 1958, Vol. 28, No. 1, pp. 127-129 (author Kushner, KH, F.)

GIGINEISHVILI , N.S.

[Karakul sheep in foreign countries] Karakulevodstvo za
rubezhom. Moskva, Izd-vo sel'khoz.lit-ry, zhurnalov i
plakatov, 1962. 477 p. (MIRA 16:8)
(Karakul sheep)

GIGINEYSHVILI, V. M.

Def. at
Tbilisi stage

Application for degree of
Candidate Paulo Henrique Sciences

GIGIMEYSHVILI, V.M.

Forecasting hail phenomena in eastern Georgia. Trudy Tbil. MIGNI
no. 2:44-67 '57. (MIRA 11:4)
(Georgia—Hail)

GIGINEYSHVILI, V. M.

Chirakadze, G. I. and Gigineyshvili, V. M.

"Scheme of the Radiation Method of Flotting the Slipperiness of Ice in Transcaucasia and the Characteristic of Slush and its Distribution in Transcaucasia."

Report presented at the Scientific Session of Tbilisi Scientific Research Institute for Hydrometeorology, May 1957. (Meteorologiya i Gidrologiya, No. 1, 1958).

PAPINASHVILI, K.I.; GIGINEYSHVILI, V.M.; NAPETVARIDZE, Ye.A.

Aerosynoptic conditions for the formation and thawing of the snow
cover of the Transcaucasian mountains. Trudy Tbil. NIGMI no.3:22
'58. (MIRA 11:10)

I. Tbilisskiy nauchno-issledovatel'skiy gidrometeorologicheskiy
institut. (Transcaucasia--Snow)

GIGINEYSHVILI, V. M.: Doc Geogr Sci (diss) -- "Hail damage in eastern Georgia in its climatic aspect". Tbilisi, 1959. 26 pp (Moscow State Order of Lenin and Order of Labor Red Banner U im M. V. Lomonosov), 150 copies (KL, № 11, 1959, 116)

GIGINEISHVILI, V. M.

307/30-52-2-24/23

367)

Author: G. M.

Title: Geological Meeting of the Tbilisi Scientific Research Institute of Hydrogeology (Armenian Scientific Research Institute of Hydrogeology)

Periodicals:

Abstract:

In May 1959 the Tbilisi nauchno-tekhnicheskoye gidrogeologicheskoye institut (Tbilisi Hydrogeological Research Institute) held a meeting in which the following representatives participated: Representatives of the Tbilisi hydrometeorological Observatory (Central Forecasting Institute), Gidrometeorologicheskaya observatory (Meteo-physical Observatory) and the local administrative organs of the Armenian hydrological service of the Transcaucasian Republic. On the occasion of the fifth anniversary of the Tbilisi Branch of the Institute, Prof. N. S. Polubarnova-Kochina, head of the Institute, delivered a speech on the occasion of the event. Dr. F. N. Polubarnova (wife on the character of temperature distribution and hydrocirculation of the atmosphere above the Antarctic). Dr. A. Shchegolev and Prof. A. Saptevants spoke on the characteristics of the circulation processes above Transcaucasia. M. M. Zhdanovskaya reported on the application of synoptic processes carried out by Dr. E. A. Buzikashvili and two papers on theoretical problems of dynamic meteorology. Prof. G. Gicheishvili and Prof. G. Dzhaparidze spoke on the present state of the Tbilisi hydrogeological institute. L. A. Shchegolev spoke on the great seismic activity in Armenia. Georgia, L. M. Martishvili on the hydrogeological variability in Georgia, Prof. N. G. Polyakova (GIG) on the hydrogeological variability in the case of precipitation and temperature. G. G. Gicheishvili and Prof. G. Dzhaparidze on the influence of precipitation on the hydrogeological processes of Georgia. Dr. J. Petrosyan on the hydrogeological situation in the Alpine zone of the Armenian Republic. Dr. L. Dzhaparidze on the radiosensitivity of the atmosphere in the case of nuclear weapons. Prof. G. Gicheishvili on different methods of surface runoff (GIG or the Grusinian hydrology GIG) in the period of the rainy season in Tbilisi. Prof. S. G. Tigranyan spoke on forecasting the amount of snow cover developed by him. Prof. V. P. Park-

Card 1/3

reported on the application of synoptic processes carried out by Dr. E. A. Buzikashvili and two papers on theoretical problems of dynamic meteorology. Prof. G. Gicheishvili and Prof. G. Dzhaparidze spoke on the present state of the Tbilisi hydrogeological institute. L. A. Shchegolev spoke on the great seismic activity in Armenia. Georgia, L. M. Martishvili on the hydrogeological variability in Georgia, Prof. N. G. Polyakova (GIG) on the hydrogeological variability in the case of precipitation and temperature. G. G. Gicheishvili and Prof. G. Dzhaparidze on the influence of precipitation on the hydrogeological processes of Georgia. Dr. J. Petrosyan on the hydrogeological situation in the Alpine zone of the Armenian Republic. Dr. L. Dzhaparidze on the radiosensitivity of the atmosphere in the case of nuclear weapons. Prof. G. Gicheishvili on different methods of surface runoff (GIG or the Grusinian hydrology GIG) in the period of the rainy season in Tbilisi. Prof. S. G. Tigranyan spoke on forecasting the amount of snow cover developed by him. Prof. V. P. Park-

Card 2/3

1960 as a method for the calculation of the volume of rain water supply in floods. Dr. G. G. Gicheishvili reported on the use of the Arseniyanaya SGB on the basis of the atmospheric circulation in hydrological forecasting. The representative of the Arseniyanaya SGB M. M. Shekharashvili reported on the characteristics of the formation of spring floods on the rivers of Armenia. A. A. Petrosyan spoke on the effect of the spring precipitation on the water supply of the belt between 1600 and 2100 m in the formation of the water supply for spring floods. Dr. G. G. Gicheishvili spoke on the method of forecasting easily measurable humidity in the soil before grain outcrops. Prof. S. G. Tigranyan spoke on the periods of the spring and spring floods in Transcaucasia. O. M. Sandoleashvili and A. A. Petrosyan (GIG of the Armenian SGB) and T. K. Chernykh spoke on the meteorological conditions of the Labatinskaya basin in the Arseniyanaya SGB. In all, 27 papers were read.

Card 3/3

GIGINEYSHVILI, V.M.; NIKOLAYSHVILI, G.V.

Conditions promoting the development of low-level clouds at a height of up to 600 meters on the principal air routes of Transcaucasia and their prediction. Trudy Tbil. NIGMI no.7: 38-48 '60. (MIRA 14:8)
(Transcaucasia--Clouds)

GIGINEYSHVILI, V.M.

Solid deposits on electric transmission and communication
lines in Transcaucasia. Trudy Tbil. NIGMI no. 8:21-33 '61.
(MIRA 15:3)

(Transcaucasia--Electric lines--Overhead)
(Transcaucasia--Ice)

ABZIANIDZE, T.S.; GIGINEYSHVILI, V.M., doktor geogr. nauk, kand. fiziko-matem. nauk, otv. red.; GOGESHVILI, E. red.; SAGARADZE, Sh., tekhn. red.

[Critique of Newton's laws and the construction of the Keplerian ellipse] Kritika zakonov N'jutona i postroenie Keplerova ellipsa. Tbilisi, Izd-vo Gruzinskogo in-ta im. V.I.Lenina. Pt.1. [Universal gravitation] O sile vsemirnogo tiagoteniia. 1961. 89 p. (MIRA 15:6)

(Gravitation) (Orbits)

GIGINEYSHVILI, V.M.; NAPETVARIDZE, Ye.A.; PAPINASHVILI, K.I.

Atmospheric processes as a factor affecting glacier fluctuations
in the Greater Caucasus. Trudy Tbil.NIGMI no.8:3-9 '61.
(MIRA 15:3)

(Caucasus--Glaciers)

ACCESSION NR: AR4008222

S/0169/63/000/011/B035/B035

SOURCE: RZh. Geofizika, Abs. 11B222

AUTHOR: Gigineyshvili, V. M.

TITLE: Hailstorms in Armenia

CITED SOURCE: Tr. Zakavkazsk. n.-i. gidrometeorol. in-ta, vy* p. 11, 1963, 45-53

TOPIC TAGS: meteorology, hailstorm, thunderstorm, hail, hailstorm chart, Armenian meteorology, Armenian hailstorm, mountain hailstorm

TRANSLATION: The orography of Armenia is conducive to hailstorm processes. Hailstorms occur most often in mountain and high-mountain regions. The greatest frequency of hailstorms in Armenia (14.6 days) occurs in the high-mountain region of Agarats (3228 m) and the regions of Kalinino and Shakhnazer (1500-1600 m) -- 7 days per year. 20-year observations were used to construct of hailstorm occurrence in Armenia. In order to determine the hailstorm trajectories, the author analyzed all of the observational material from 1950 to 1959. The orographic peculiarities of a particular location dictate the placing of trajectories in

Card 1/2

ACCESSION NR: AR4008222

individual groups. There are seven such groups. These hailstorm groups are interesting because on the territory assigned them, the hailstorms develop, attain their greatest intensity, and die out, rarely passing into a region belonging to another group. The hailstorm trajectories chiefly lie in a west-to-east direction and are of relatively short length. The greatest frequency of hailstorms is in the central and northwestern regions. The western and southwestern parts are subject to hailstorms whose foci are located on the territory of adjacent countries. The remaining regions are subject to hailstorms whose foci are located in the mountain ranges in the inner part of Armenia. N. Myachkova.

DATE ACQ: 09Dec63

SUB CODE: AS

ENCL: 00

Card 2/2

GIGINEYSHVILI, V.M.; AYVAZYAN, V.Ye.

Aerosynoptic conditions of the origin of hail phenomena in
Armenia. Trudy TbilNIGMI no.12:3-20 '63.
(MIRA 18:5)

GIGINEYSHVILI, V.M.

Appearance of local characteristics of circulation processes
of the atmosphere under conditions of a mountainous relief
for the purpose of forecasting heavy precipitation and the
significant rain floods on the principal and mountain rivers
of Georgia related to it. Trudy ZakNIGMI no.18:3-36 '65.
(MIRA 19:1)

L 2307-66 E.I.(1)/FCC GW/JXT(CZ)
ACC NR: AT6006487

SOURCE CODE: UR/3061/65/000/018/0003/0036

26

AUTHOR: Gigineyshvili, V. M.

B+1

ORG: none

TITLE: Evaluation of local characteristics of atmospheric circulation processes in mountainous regions as a means of forecasting heavy precipitation and floods in the mountain rivers of Georgia

SOURCE: Tiflis. Zakavkazskiy nauchno-issledovatel'skiy gidrometeorologicheskiy institut. Trudy, no. 18(24), 1965. Voprosy gidrometeorologii (Problems in hydro-meteorology), 3-36

TOPIC TAGS: weather forecasting, atmospheric precipitation, atmospheric circulation

ABSTRACT: The investigation analyzes the high altitude formation of a warm front over Western Georgia and its precipitation characteristics; the spring floods of the Chkherimela River resulting from melting snow; the effect of cyclonic occlusion processes over the Suramskiy Range and other mountainous regions of Georgia. Fundamental cold fronts and their effects on precipitation and river floods, and the effect

UDC: 551.51 : 551.509 : 551.48

Card 1/2.

L 29907-66
ACC NR: AT6006487

O

of the cold-front wave disturbances on the Georgian climate were also studied. The study shows that 1) the high-altitude warm front does not induce high precipitation unless it is followed by the fundamental cold front; 2) cyclone occlusion processes are responsible for high precipitation and floods of the mountain rivers; 3) the wind-type cold fronts formed in the coastal parts of Western Georgia usually bring heavy precipitation; 4) two main cold fronts passing over Western Georgia are responsible for heavy precipitation; and 5) cold-front wave disturbances induce heavy and unevenly distributed precipitation. Orig. art. has: 14 figures and 21 tables.

SUB CODE: Q4/ SUBM DATE: 22Jun65/ ORIG REF: 007/ OTH REF: 000

Card 2/2 (1)

GIGITASHVILI, M.S.

Aepidium therapy of hymenolepiasis; preliminary report. Med. paraz.
i paraz. bol. no.2:136-137 Ap-Je '54. (MLRA 7:8)

1. Iz Instituta malyarii i meditsinskoy parazitologii imeni prof.
S.S.Virsaladze Ministerstva zdravookhraneniya Grusinskoy SSR (dir.
instituta G.M.Marashvili)

(TAPEWORM INFECTON, therapy,
*Dryopteris filix-mas extracts)

(PLANTS,
*Drypteris filix-mas extract, ther. of hymenolepiasis)

ST' AGBUJI, V. T.

AGBUJI, V. T.: "Morphological and volumetric changes in the blood in encylostomiasis." Georgian State University, Institute of Medical Literature. Tbilisi State Medical Inst., Tbilisi, 1954. (Dissertation for the degree of Candidate in medical Sciences).

SC: Entzmann [unclear], No. 23, 1954

GIGITASHVILI, M.S.

Geophagia in Schylostomiasis, Med.paraz. i paraz.bol. 25 no.3:273
Jl-8 '56. (MIRB 9:10)
(DIRT-EATING) (HOOKWORM DISEASE)

GIGITASHVILI, N.S.

Comparison of methods for treating teniasis. N.S. Gigitashvili.
Med. paraz. i paraz. bol. 27 no.2:218 Mr-Ap '58 (MIRA 11:5)

l. In Nauchno-issledovatel'skogo instituta malyarii i meditsinskoy
parasitologii imeni prof. Virsaladze.
(TAPEWORMS)

GORDADZE, G.N.; GIGITASHVILI, M.S.

Epileptoid seizures during hymenolepiasis induced by dwarf tapeworms.
Med.paraz. i paraz.bol. 28 no.4:430-434 Jl-Ag '59. (MIRA 12:12)

1. Iz Nauchno-issledovatel'skogo instituta malyarii i meditsinskoy
parazitologii imeni S.S. Virsaladze Ministerstva zdravookhraneniya
Gruzinskoy SSR (dir. instituta - prof. G.M. Maruashvili).
(TAPEWORM INFECTION complications)
(EPILEPSY etiology)

GIGITASHVILI, M.S.; ZIRAKISHVILI, L.M.

Treatment of cases of ancylostomiasis with carbon tetrachloride
through a duodenal tube. Med.paraz.i paraz.bol. 29 no.4:416-418
(MIRA 13:11)
Jl.-Ag '60.

1. In Instituta meditsinskoy parazitologii i tropicheskoy medi-
tsiny imeni S.S. Virsaladze Ministerstva zdravookhraneniya
Gruzinskoy SSR (dir. instituta I.I. Topuriya, na v. klinicheskim
otdeleniyem T.K. Zhordaniya).
(HOOKWORM DISEASE) (CARBON TETRACHLORIDE)

GIGITASHVILI, M.S.; ABULADZE, T.Ye.; MDIVANI, I.M.

Four cases of fascioliasis in man treated with emetine and
carbon tetrachloride. Med.paraz.i paraz.bol. no.5:526-527
'61. (MIRA 14:10)

1. Iz klinicheskogo otdeleniya Nauchno-issledovatel'skogo instituta
meditsinskoy parazitologii i tropicheskoy meditsiny Gruzinskoy SSR
imeni S.S. Virsaladze (dir. instituta I.I. Tsvuriya, rukovoditel'
otdeleniya T.K. Zhordaniya).
(LIVER FLUKE) (EMETINE) (CARBON TETRACHLORIDE)

CHIKHVA, N.N.; CHIKHVA, V.I.; KALININ, L.V.; KARAKHAN, A.A.; KARATASHKA, L.A.; KHOLODINA, D.S.; KREK, V.Ye.; TAKAISHVILI, M.L.; TIKHONOV, A.Z.; TIKHONOV, I.Ya.; AMUTIAN, R.A.; BAGDASHVILI, N.I.; BULGRENKO, V.A.; FUNSKAYA, I.G.; BURMISTROVA, O.G.; POGORELSKAYA, S.A.; D'YACHENKO, Z.P.; TOFURIYA, I.I.; MATABELI, G.V.; GIGITASHVILI, N.S.; VACHNADZE, Z.G.; MAGHRIB, N.P.; NABIVEV, E.G.; BICERKOV, V.T.

Abstrakti. Zhur. mikrobiol., v.10. i izdani. 41 no.1(1964-1977)
Ap. 104. (V.1A 18:4)

1. Moskovskiy institut epidemiologii i mikrobiologii (for Kurnosova). 2. Filatovskoye rayonnoye bol'ničnoe Molodёžnogo SSR i Vladičinskoy meditsinskoy institut imeni Pirogova (for Boncharenko). 3. Stavropol'skiy institut včetaini i svyazotek (for Bakuriani). 4. Kalinubelskiy oblastnyj otdel naravosobrannosti (for Yarunina, Tsvetkova). 5. Dantsevskiy meditsinskiy institut (for Manolova). 6. Tbilisijskaya rayonnoye imeni L. Mirzasejanishevskoye sanitarno-epidemiologicheskoye tsentral'noye (for Shapua, Babashvili). 7. Kemerovskiy naftosibkiy institut (for Protsenko). 8. Turkmenskiy meditsinskij institut (for Funskaya, Burmistrova). 9. Gor'kovskiy Institut epidemiologii i mikrobiologii i Gor'kovskaya rayonnoye sanitarno-epidemiologicheskaya stantsiya (for Pogorelskaya, D'yachenko). 10. Institut naftosibskoy parazitologii i tropicheskoy meditsiny imeni Vsesoyuznogo Ministerstva prirodorazvedenija i Gruzinskoy SSR (for Tepuriya, Matabeli, Gigitashvili, Vachnadze). 11. Almatinskij institut universitetovozrastnoj vrachey (for Natiyev).

GIGGLOV, M. G.

Giglov, M. G.: "Comparative evaluation of the surgical treatment methods of Echinococcus," (Report, Trudy III Zakavkazsk. s"yezda khirurgov, Yerevan, 1949 (on cover: 1949), p. 270-278

SO: U-5240, 17 Dec. 53, (Letopis 'zurnal 'nykh Statey, No. 25, 1949).

GIGOROVICI, R.; CROITORU, N.; DEVENYI, A.

Structure and electric conductivity of the silver thin layers.
Studii cerc fiz 11 no.4:897-920 '60. (EEAI 10:8)

1. Institutul de fizica, Bucuresti.
(Electric conductivity) (Thin films) (Silver)

ORLOVA, T.Yu.; GIGER'YEV, A.I.; NOVOSELOVA, A.V.

Aluminum alkoxacetates. Izv. AN SSSR. Neorg. mat. 1 no.5:633-637
Mys '65. (MIRA 18:10)

1. Moskovskiy gosudarstvennyy universitet imeni Lomonosova,
khimicheskiy fakul'tet.

GIGOV, A.

BULGARIA/Zooparasitology - General Problems.

G-1

Abs Jour : Ref Zhur - Biol., No 6, 1958, 24315

Author : Gigov, A.

Inst :

Title : Parasitic Protozoa in the Human Oral Cavity.

Orig Pub : Stomatologiya (Bulg.), 1956, No 6, 375-377

Abstract : No abstract.

Card 1/1

ANGELOV, S., Akad.; GIGOV, A.; NIKOLOV, P.; AMIDZHIN, S.

Toxoplasmosis and its studies in Bulgaria. Suvrem. med.,
Sofia 7 no.11:79-83 1956.

(TOXOPLASMOSIS, epidemiology,
in Bulgaria (Bul))

ANGELOV, St., Akad.; GULUBOV, S., D-r.; GIGOV, A., D-r.; NIKOLOV, P., D-r.

Toxoplasmosis in domestic animals in Bulgaria. Izv. Mikrob. inst.,
Sofia no.8:35-40 1957.

(TOXOPLASMOSIS

in domestic animals, statist. in Bulgaria)

(ANIMALS, dis.

toxoplasmosis in domestic animals, statist. in Bulgaria)

GIGOV, A.

Country : TURKEY
City : Ankara
Address : Ankara International Pediatric Hospital, Marmaray
Date, year : 1986, No. 2-1986, C. 147
Name : GIGOV, A.; GIGOV, A.; GIGOV, A.
Title : Head of Dept., Admin AB
Other : Student of Foreign Languages in Ankara

Report No. : 1986, Vol. 1, No. 1, P. 2-13.

Report No. 1986, Vol. 1, No. 1, P. 2-13. The report is dated 1986, No. 2-1986, C. 147. The subject is GIGOV, A. The report is in English. The subject is a child, female, born in 1981. The subject has brown hair. In the address of the child, there is no name. In the address of the child, there is no name. The complement-fraction that can alter the name.

*GIGOV, A.

RECOVERY:

TIME: 10:00

PLACE: P. L. B. S. Butcher, 2000 14th St., N.W.

TESTS:

TESTS:

TESTS:

TESTS:

TESTS: The test for phenyl isocyanide, the latter suffices to detect phenyl isocyanide in hair cellulose. In three out of four coats which were in the sample, the ninhydrin and carbonyl reactions were positive. The isocyanide-diazotization test was used to investigate the nature of the stains with ammonium hydroxide, etc., and by colorimetric, titration, refractive, absorption, etc. In 15% of the samples the protein was positive, of 259 samples infected with oil-powdered and other mortal

TESTS: TESTS:

1970-1971
1971-1972

July 1971, 1972, 1973, 1974, 1975

1971-1972
1971-1972
1971-1972

1971-1972

In 1971-1972, the indirect agglutination test was performed on peripheral blood samples from 30 patients and 16 cattle. A positive result in 22 out of 46 cases indicated a high incidence of spontaneous abortions and premature parturition. In 1971-1972, the complement-fixation test was positive in 11. Positive allergic reactions were noted in 5 heads of cattle (out of 11), 10 goats (out of 11), 1 sheep (out of 12).