

GOL'DBERG, V.V.

Mapping of Legendre's or n-dimensional projective space onto
a stationary hypersurface. Sib. mat. zhurn. 5 no. 1 3-53
Ja-F '64. (MIR 17.7)

RECORDED BY: [Redacted] DATE: [Redacted]
TIME: [Redacted]

REPORTER: [Redacted] DATE: [Redacted]
TIME: [Redacted]
MATERIAL: [Redacted]

OKLAHOMA CITY, OKLAHOMA, U.S.A.

Wide-faced meadowlarks begin appearing in the area about July 15.
no. 63-37 34-3 162.

EWT(m)/BDS--AFFTC/ASD--DM

L 1120L-63

ACCESSION NR: AP3001178

S/0089/53/014/005/0432/0484

55

AUTHOR: Artemov, K. P.; Gol'dberg, V. Z.; Rudakov, V. F.

TITLE: Elastic and inelastic scattering of Alpha particles by Al²⁷

SOURCE: Atomnaya energiya, v. 14, no. 5, 1963, 482-484

TOPIC TAGS: elastic scattering, inelastic scattering, Alpha particles, excited states of Al²⁷

ABSTRACT: The Alpha particles were accelerated to 40, 38, and 36 Mev in the 1.5 m cyclotron of the Institute for Atomic Energy. A high pressure ionization chamber was used for detection of Alpha particles. In agreement with the results of other workers, the angular distribution of scattered particles showed a "diffraction pattern," the maxima of the inelastically scattered particles coincided with the minima of those scattered elastically. The results are interpreted on the basis of theory by other authors, among them S. I. Brodov (Zh. experim. i theoret. fiz., 31, 901, 1956). The radius of interaction of the Alpha particle with the Al²⁷ nucleus is found to be 5.5 fermi. Conclusions are made concerning the excited states in Al²⁷. "The authors are grateful to S. I. Brodov for the discussion of results of the work." Orig. art. has: 3 figures and 6 references.

Card 1/2

8/0056/64/047/002/0571/0576

ACCESSION NR: AP4043632

AUTHORS: Gol'dberg, V. Z.; Rudakov, V. P.; Serikov, I. N.

TITLE: Analysis of elastic scattering of He-3 and Alpha particles
on the basis of the optical model of the nucleus

SOURCE: Zh. eksper. i teor. fiz., v. 47, no. 2, 1964, 571-576

TOPIC TAGS: helium, elastic scattering, alpha particle reaction,
differential cross section, optical potential

ABSTRACT: Although a detailed analysis of elastic scattering of alpha particles by many nuclei from Cl²² to Th²³² was made by Igo and Thaler (Phys. Rev. v. 106, 126, 1957), no such analysis was made for the elastic scattering of He³. Earlier calculations, made on the basis of a limited experimental material, have led to parameters that vary erratically from nucleus to nucleus. The authors have therefore used the optical model to attempt a more systematic

Card 1/2

ACCESSION NR: AP4043632

analysis on the data concerning elastic scattering of He^3 by different nuclei, and calculated the differential cross sections for this scattering. New data obtained on the differential cross sections at the laboratory of the authors (V. M. Pankratov and I. N. Serikov, ZhETF, v. 44, 187, 1963) and by Gonzalez-Yidal et al. (UCRL-9566, 1961) have been used in the calculations. The results show that a single set of parameters for the optical potential can be used to describe satisfactorily the experimental data over the wide range of nuclei from Be^9 to Bi^{209} . A comparison is given of the parameters of the potentials describing the elastic scattering of He^3 and of alpha particles by Al^{27} . "The authors thank V. A. Belyakov, P. E. Nemirovskiy, and I. S. Shapiro for useful discussions." Orig. art. has: 3 figures and 3 tables.

ASSOCIATION: None

ENCL: 00

SUBMITTED: 28Jan64

OTHER: 010

SUB CODE: NP

NR REF SOV: 002

Card 2/2

ARTEMOV, K.P.; GOL'DBERG, V.Z.; ISLAMOV, E.I.; RUDAKOV, V.P.; SEMIKOV, I.N.

Elastic scattering of He³ ions on Be⁹, N¹⁴, and O¹⁶. IAI. fiz.
(MIRA 16:5)
1 no.4:620-632 Ap '65.

APPROVED FOR RELEASE: Thursday, September 26, 2002 CIA-RDP86-00513R000515620006-6"

AUTHORS: Samoylov, A., Gol'dberg, Ya.

SOV/29-58-6-22/23

TITLE: Forks No Longer Break (Vilki perestali lomat'sya)

PERIODICAL: Tekhnika molodezhi, 1958, Nr 8, pp. 40-40 (USSR)

ABSTRACT: Many cyclists know very well that while they ride along country roads or paved streets at high speed, the fork of their bicycle, which is subjected to considerable vibrational stress, breaks easily. The authors found a simple and reliable method of counteracting this danger by fitting out the front fork of their bicycles, which are provided with a "D-4" motor, with a damper or shock-absorber ("amortizer"). This makes it possible to ride along any kind of road at top speed. Besides, this simple device prolongs the life of the motor. The device is then described. The authors express the opinion that factory-produced forks are more simple and of lighter weight than those made by hand. There is 1 figure.

1. Bicycles--Equipment

Card 1/1

BATALOV, N.; GOL'DBERG, Ya.

34 times, such is the increase in volume of transports in twenty years. *Grazhd. av.*, 21 no.10-1-3 C '64. (MIRA 18:3)

1. Komandir Litovskiy oddel'nyy avisegruppy grazhdanskoy aviatsii (for Batalov). 2. Zamestiteli komandira po politicheskoy chasti Litovskiy oddel'nyy avisegruppy grazhdanskoy aviatsii (for Gol'dberg).

GOL'DBERG, Ya. M.

GOL'DBERG, Ya. M. "Penicillin therapy of typhoid fever", Trudy Kishinevsk gos. med. in-ta, Vol. 1, 1949, p. 115-21.

SO: U-3:61, 10 April 53 (Letopis - Zhurnal 'nykh Stat'ev No. 11, 1949)

GOL'DBERG, Ye.D.

[Hematological changes in acute radiation sickness caused by a
25 Mev betatron] Gematologicheskie sviggi pri ostroii luchevoi
bolezni, vyzvannoi na betatrone 25 MEV. Tomsk, Izd-vo Tomskogo
univ., 1960. 39 p. (MIRA 13:9)

(RADIATION SICKNESS) (BLOOD)

PHASE OF EXPLOITATION 307/5042

Author: V. I. D.

General information about acute radiation sickness caused by betatron 25 Mev. (H. L. S. et al.) Opis iktivnosti Acute Radiation Sickness Caused by Emission of 25 Mev. Betrons. Ed. Tomsk. Izd. na Tomskogo univ., 1960. 40 p. 5,000 copies printed.

Soviet Army Academy, Tomsk, Mathematics Institute and Siberian filial Vsesoyuznaya radiobiological laboratory.

Ed.: L. M. Tsyplakova. Profs. Dr. math. Ed. A. T. Osoevskiy

PURPOSE: This article was intended for research scientists and engineers concerned with the biological effects of ionizing radiation.

COVERAGE: The article presents the results of biological experiments on radiation damage caused by the ionizing radiation produced by a 25 Mev betatron. Most of the work is concerned with the effects on the blood and the circulatory system. The book also contains a survey of the existing literature on the effects of betatron radiations on the blood system. The fol-

Card 1/2

Hematological Complications (Cont.)

SOV/5042

N. M. Ogiyenko, M. N. Meysel, V. A. Sondak, Ye. S. Kirpichnikova and
N. N. Kurshakova (nucleic acids in blood cells). There are 59 references:
37 Soviet, 9 German, 7 English, 4 Swiss, 1 Italian, and 1 Czech.

TABLE OF CONTENTS:

Introduction	3
Brief Information on the Betatron	5
Data in the Literature on the Effect of the Radiation From a Betatron on the Blood System	7
Characteristics of the Blood Count in Normal Guinea Pigs	10
Characteristic Data on the Changes in the Blood and Bone Marrow of Guinea Pigs in Acute Radiation Sickness	13
Kurlov Bodies in Radiation Sickness	16

Card 3/4

GOL'DBERG, Ye.D.

Blood picture and bone marrow hemopoiesis in guinea pigs in acute radiation sickness caused by irradiation on a 25 Mev. betatron. Med. rad. 5 no.1:28-35 Ja '60. (MIREA 15:3)

1. Iz kafedry patofiziologii (zav. - prof. D.I. Gol'dberg) i 2-y betatronnoy laboratorii (zav. - kand.med.nauk G.P. Garganeyev) Tomskogo meditsinskogo instituta.
(SLCDD CELIS)
(RADIATION SICKNESS)

GOL'DBERG, D.I., nasl. chet' nauki ASFSR, prof.; GOL'DBERG, Ye.D.;
TOROPTSEV, I.V., prof., red.; OSOVSKIY, A.T., tekhn. red.

[Handbook of hematology with an atlas of microphotographs]
Spravochnik po hematologii s atlasom mikrofotogramm. Tom 1,
Izd-vo Tomskogo univ., 1961. 121 p. (MFA 15:10)

1. Chlen-korrespondent Akademii meditsinskikh nauk SSSR (for
Toroptsev).

(HEMATOLOGY)

GOL'DBERG, Ye.D.; GOLOSOV, O.S.; POTEKHIN, K.G.

Hematological indices in workers of roentgenological and radiologica departments. Med.rad. no.5:49-54 '61. (MIRA 14:11)

1. Iz kafedry patofiziologii Tomskogo meditsinskoj instituta i travmatologicheskoy bol'nitsy Prokop'yevskaya.
(BLOOD CELLS--RADIOGRAPHY) (RADIOLOGISTS)

GOL'DBERG, Ye.D.

Blood picture in healthy guinea pigs. Biul. eksp. biol. i
med. 52 no.7:115-118 Jl '61. (MIRA 15:3)

1. Iz lefedy patofiziologii (zaveduyushchiy - prof. D.I.
Gol'dberg) Tomskogo gosudarstvennogo meditsinskogo instituta.
Predstavlena akademikom V.N. Chernigovskim.
(BLOOD--EXAMINATION)
(GUIDEA PIGS)

H/021/62/000/006/001/002
D296/D307

AUTHORS: Gol'dberg, Ye.D., Golosov, O.S. and Potekhin, K.G.

TITLE: Hematological indices found in X-ray and radiotherapy departmental staff

PERIODICAL: Magyar Radiologia, no. 6, 1962, 321-326

TEXT: The authors analyzed the blood of 130 patients exposed to continuous small doses of ionizing radiation by reason of their occupation, and of 75 healthy control subjects not previously exposed to radiation. It was found that the staff of X-ray and radiotherapy departments were on the average exposed to a daily dose of 0.02 - 0.03 r. Some of the subjects complained of occasional headaches, tiredness, and in a few cases skin changes, pigmentation and loss of hair could be observed. In 17% of the exposed persons the white cell count was decreased and in 6.1% it was increased. Among the staff of radiotherapy departments, neutropenia was found in 90.9% of those who worked in these departments for less than 5 years, but only in 75% of those working for more than 5 years.

Card 1/2

GOL'DBERG, Yevgeniy Daniilovich; TOROPTSEV, I. V., prof., med.;
MORDOVINA, L.G., red. izd-va

[Leukemia and radiation] Leikozy i radiatsiiia. Tomsk,
Izd-vo Tomskogo univ., 1963. 71 p. (MIRA 16:7)

1. Chlen-korrespondent Akademii meditsinskikh nauk SSSR
(for Toroptsev).
(LEUKEMIA) (RADIATION--PHYSIOLOGICAL AFFECT)

GOLD'BERG, D.I., prof.: LEVINA, G.P.; FALINGER, I.E.; KURPOVA, G.V.;
GOL'DBERG, Ye.I.; TUTERINA, V.I.; LAVROV, V.S.; TIKHIN, N.P.;
GOL'DBERG, A.I.; CHUDINA, Ye.A.

Clinical significance of erythrocytes try. Probl. germt. i perel.
krovi 9 no.19:6-14 (1984) (ZhM 18:3)

1. Tomskiy meditsinskiy institut.

"APPROVED FOR RELEASE: Thursday, September 26, 2002

CIA REF ID: A6520006
APPROVED FOR RELEASE: Thursday, September 26, 2002 CIA-RDP86-00513R000515620006-6"

ACC-NR: A00000000

SOURCE CODE: UF/0240/57/000/001/0023/0096

AUTHOR: G. V. Lurye, T. N. Kuznetsova

ORG: Moscow Municipal Sanitary-Epidemiological Station (Moskovskaja gorodskaya sanepidstantsiya)

TITLE: Determining small concentrations of carbon dioxide in the air with the FEK-N-54 photocalorimeter

SOURCE: Gigiyena i sanitariya, no. 1, 1967, 93-94

TOPIC TAGS: photocalorimetry, carbon dioxide sensor, colorimetry, carbon dioxide/FEK N 54 photocalorimeter

ABSTRACT: The following procedures are used in determining CO_2 concentration in air spectrophotometrically: The air sample is shaken up with an absorbent containing one part 0.125% bromothymole blue and 50 parts of NaHCO_3 . After interaction with the absorbent, optical density is measured at $\lambda_{\text{max}} = 600 \text{ nm}$ using an SF-5 spectrophotometer in a cell with a ten-mm layer. The concentration of CO_2 is then found using a graduated graph showing optic density as a function of CO_2 . Using a variant of this method, the author devised an approach employing the FEK-N-54 photocalorimeter. Air samples were collected in 50--100-ml syringes from closed spaces (dimensions are given). A cell with a ten-mm layer and

Card 1/3

UDC: 614:72:661.993-074

ACC NR: AP7003545

Table 1. Comparative evaluation of titro-metric and photometric or methods of determining CO₂ concentration in closed spaces.

No.	CO ₂ con-		O ₂
	centration	(vol.%)	
1	0.15	0.683	0.15
2	0.15	0.671	0.15
3	0.15	0.676	0.15
4	0.15	0.681	0.15
5	0.15	0.677	0.15
6	0.15	0.675	0.15
7	0.15	0.677	0.15
8	0.15	0.677	0.15
9	0.15	0.676	0.15
10	0.15	0.680	0.15

Card 2/3




Fig. 1. Standardized graph showing the dependence of the optical density of the absorbent on CO₂ concentration; ordinate--optical density; Abscissa--CO₂ concentration (in vol.%)

ACC NR: APT003545

no. 7 (orange) filter was used. Some results of a comparison of this method with a titrometric one are shown in Table I. An example of the graph described above is shown in Fig. 1. It was concluded that the author's method facilitates the determination of CO_2 concentration in closed spaces with $\pm 3.6\%$ accuracy during a five-min test. Orig. art. has: 1 table and 1 figure.

SUB CODE: 06/ SUBM DATE: 17Mar66/ ATD PRESS: 5112

Card 3/3

GOL'DBERG, Ye.N.

Correcting the output pulses. Avtom., telem. i sviaz' 2 no. 8:35
Ag '58. (MIRA 11:8)

1. Starshiy inzhener sluzhby signalizatsii i svyazi Estonskoy
dorogi.
(Railroads--Telephone)

GOL'DBERG, Ye.N.

Remote control of an audio generator in railroad radio communication. Avtom. telem. i sviaz'. 4 no.5:22-23 My '60.
(MIE: 13:8)

1. Nachal'nik Pyatnuskoy distantsii signalizatsii i svyazi
Sotseskoy dorogi.
(Railroads--Communication systems)

GOL'DBERG, Ye.N., inzh.

Pulse modulators using junction transistors. Sov. trud. LITZHT
no.224:75-99 '64. (MIRA 18:9)

GOL'DBERG, Yu., inzh.

Model plans for enlarging rural hospitals. Sel'. stroi. 15
no.4: insert: 1-3 Ap '61. (MIRA 14:6)
(Hospitals, Rural—Construction)

ACC NRI AP6030155

SOURCE CODE: UR/0120/66/000/004/0189/0193

AUTHOR: Gol'dberg, Yu. A.; Nasledov, D. N.; Tsarenkov, B. V.

ORG: Physico-Technical Institute, AN SSSR, Leningrad (Fiziko-tehnicheskiy institut
AN SSSR)

TITLE: The ohmic contact between gallium arsenide and indium

SOURCE: Pribory i tekhnika eksperimenta, no. 4, 1966, 189-193

TOPIC TAGS: gallium arsenide, indium, semiconductor research

ABSTRACT: The wetting of gallium arsenide surface with indium, and the extent of fusion and contact resistance as a function of temperature and fusion time were studied. It is shown that 100% wetting and minimum contact resistance occur at a temperature of 500°C and above. The GaAs-In junction was obtained by fusion in hydrogen. Hydrogen was used as the reducing medium to prevent the oxidation of In and GaAs at high temperatures. To prevent the explosion of the hydrogen-air mixture, a neutral gas was passed through the system before and after the hydrogen was turned on. The gases were dried by cooling them to a temperature of -196°C. Activated charcoal was used to purify H₂ and He at liquid nitrogen temperature. The following parameters were determined during the fusion process: the edge wetting angle, contact resistance, wetting coefficient, depth of fusion, and hole shape. The reduced resistance of the n-GaAs-In

UDC: 621.382.032.27

Cord 1/2

ACC NR: AP6030155

contact was 10^{-5} ohm·cm² while that of the p-GaAs-In contact was $10^{-4}\text{--}5\cdot10^{-5}$ ohm·cm². The author expresses his gratitude to A. D. Forelenk, Ye. A. Posse, and V. P. Yurochkin for their assistance. Orig. art. has: 5 figures.

SUB CODE: 20,09/ SUBM DATE: 16Jul65/ ORIG REF: 007/ OTH REF: 004

Card 2/2

GOL'DBERG, Yu.A., inzh.; SEMENOVKER, I.Ye., kand.tekhn.nauk; CHAKRYGIN,
V.G., kand.tekhn.nauk

Study of the operation of the radiational section of a PK-12
boiler. Teploenergetika 10 no.1:34-40 Ja '63. (MIRA 16:1)

1. TSentral'nyy nauchno-issledovatel'skiy kotloturbinnyy institut
imeni I.I.Polzunova i Vostochnyy filial Vsesoyuznogo nauchno-issle-
dovatel'skogo teplotekhnicheskogo instituta.
(Boilers)

GOL'DBERG, Yu.A., inzh.; SEMENOVSKIY, I.Ya., kand.tehn.nauk; MARYGIN, V.G.,
kand.tehn.nauk

Assurance of adequate temperature of the water walls of boilers operating
on pulverized coal. Elek. sta. 74 mili:ll-16 N 62.
(MIRA 1712)

ACC NR: A07001959

SOURCE CODE: UR/0120/66/000/006/0,6J/0184

AUTHOR: Gol'dberg, Yu. A.; Nasledov, D. N.; Tsarenkov, B. V.

CRA: Physicotechnical Institute, Academy of Sciences SSSR, Leningrad (Fiziko-
tekhnicheskiy institut AN SSSR)

TITLE: Thin multilayer gallium arsenide-metal contacts

SOURCE: Pribory i tekhnika eksperimenta, no. 6, 1966, 180-184

TOPIC TAGS: ohmic contact, multilayered ohmic contact, gallium arsenide, gold, tin,
nickel, zinc, silver, copper

ABSTRACT:

A method of manufacturing gallium arsenide-metal contacts by chemical deposition of thin metal layers has been developed. The method permits uniform coating of gallium arsenide with thin (about 1 μ) layers of various metals with a very small (1 μ) depth of fusion. The main advantage of the small depth of fusion is that the crystals can be cleaned thoroughly with the deposited metals. It was found that with only one metal, the contact was either nonohmic, not sufficiently low-ohmic, or technologically unsuitable. The best low-ohmic contacts were obtained with several layers of various metals deposited on gallium arsenide. For instance a contact

Card 1/2

UDC: 621.382.032.27

ACC NR: AP7001959

on n-type gallium-arsenide coated with Au-Sn-Ni-Au (deposited in that order) has a resistance (per unit area) of 10^{-5} ohm/cm²; a contact on p-type gallium arsenide coated with Au-Zn-Ni-Au has [a resistance of 10^{-4} ohm/cm². Orig. art. has: 2 figures and 2 tables.

SUB CODE: 11, 09/ SUBM DATE: 03Dec65/ ORIG REF: 004/ OTH REF: 007
ATD PRESS: 5111

Card 2/2

L 37687-66 ECO(k)-2/EWF(k)/EWT(1)/EWT(m)/FBD/T/EWP(t)/ETI IJP(c) WG/JD
ACC NR: AP6024502 SOURCE CODE: UR/0181/66/008/007/2251/2253

AUTHOR: Gol'dberg, Yu. A.; Nasledov, D. N.; Tsarenkov, B. V.

72
B

ORG: Physicotechnical Institute im. A. F. Ioffe, AN SSSR, Leningrad (Fiziko-tehnicheskiy institut AN SSSR)

TITLE: Dependence of electroluminescent parameters of GaAs lasers on the angle between the p-n junction plane and the resonator mirrors

SOURCE: Fizika tverdogo tela, v. 8, no. 7, 1966, 2251-2253

TOPIC TAGS: semiconductor laser, gallium arsenide laser, diode laser, laser output, gallium arsenide, laser, pn junction

ABSTRACT: The threshold current density and the output of diode lasers were investigated experimentally as a function of the angle ($\phi = 90^\circ + \theta$) between the p-n junction plane (100) and the resonator mirrors placed in the (110) plane. It was shown that: 1) the threshold current density decreased with an increase in the distance between mirrors 1 (Fig. 1), and with a decrease in the angle when $l = \text{const}$ (Fig. 2); and 2) quantum yield increased with a decrease in θ (Fig. 2). The maximum angle $\theta_{\max} = \frac{d}{l}$ (where $d = \text{width of active medium}$) for which the rereflected

Card 1/3

L 37687-66

ACC NR: AP6024502




Fig. 1. Dependence of threshold current density on the distance between mirrors




Fig. 2. Dependence of threshold current density (curve 1) (for $l = 0.7 \text{ mm}$) and quantum yield (curve 2) on the angle between the p-n junction plane and resonator mirrors

Card 2/3

L 37687-66

ACC NR: AP6024502

beam will travel the entire length of the active medium was estimated roughly at 11'—18', for $d = 2-3 \mu$ and $l = 0.5-0.7$ mm. Orig. art. has: 2 figures and 2 formulas. [YK]

SUB CODE: 20/ SUBM DATE: 26Jan66/ OTH REF: 002/ ATD PRESS: 504/

Card 3/3

GOL'DBERG, Yu.I. (Moskva); ALEXSAKHIN, S.P. (Moskva).

Vasilii Grigor'evich Chichigin. Mat.v shkole no.1:75-76 Ja-F '57.
(Chichigin, Vasilii Grigor'evich, 1885-) (MLRA 10:2)

G.D.B.C.G., P.M.L. (Lentova)

"(1) (a); tested for the presence of A.M. by A.M. Lamm, ave.
" (2) (b) (1) G. 150 mg. (3) (b) (1) vehicle no. 17-63-1
" (4) (b) (1) (5) (b) (1) (6) (b) (1) (7) (b) (1) (8) (b) (1)

(A) (1) (a)
" (2) (b) (1) (3) (b) (1) (4) (b) (1) (5) (b) (1) (6) (b) (1) (7) (b) (1) (8) (b) (1)

GOL'DBERG, Yu.M., inzh.

Installation of unprotected bus conductor lines. From energ.
18 no.1:38-40 Ja '63. (MIRA 16:4)
(Electric lines--Overhead)
(Bus conductors (Electricity))

SEMIOSHKO, V.M., gornyy inzh., GOL'DINERG, Yu.S., gornyy inzh.

Complete treatment by flotation of 2d- and 3d- class manganese
concentrates. Gorn. zhur. no.10-58-61 O '69. (MIRA 16:11)

1. Mekhanobrchermet, Krivoy Rog.

Tungstic anhydride. P. N. Lyubimov and Z. A. Goldberg. Russ. 60,442, Feb. 28, 1937. Wolframite or W-contg. shavings and scale are treated with Na₂CO₃ and KNO₃, the soln. of Na₂WO₄ obtained is treated with phenols, the product is filtered and H₂WO₄ is formed in the usual manner by action of HCl. H₂WO₄ is converted into WO₃ by heat treatment.

AIA-SLA - METALLURGICAL LITERATURE CLASSIFICATION

C2

SHAPOSHNIKOV, I.G.; GOL'DBERG, Z.A.

Absorption of sound in binary mixtures. Zhur. eksper. teor. Fiz. 23, No.
4, 425-9 '52. (MLRA 5:12)
(PA 56 no.668:5342 '53)

TOEFL DOCUMENT

Abs Jour: Radiation Dosimetry and Radiobiology

Author : ~~W. L. HARRIS~~

Inst : University of California, Berkeley, CA, USA

Title : Absorbed Dose and Dose Rate in Irradiated Cells
From Thermoluminescence Analysis

Orig Eds: MR

Abstract: Radiation dose and dose rate parameters can be determined from the amplitude and decay time of the thermoluminescence signal in the irradiated sample. The dose rate is proportional to the rate of change of the signal amplitude with time. The dose can be calculated from the dose rate and the dose rate constant. It is shown that the thermal energy released during irradiation must be taken into account in calculating the dose rate. A method is described for determining the dose rate and dose constants from the thermoluminescence signal. Following ESR, the dose rate is determined by the rate of spin trapping. The dose rate is determined by the rate of spin trapping. The dose rate is determined by the rate of spin trapping.

Card : 4/2

: POLAND/Acoustics.

J

Abs Jour : Referat Zhur-Fizika. 1987, No 4, 1014.

order, it is possible to separate from the hydrodynamic equations with viscosity the equations of first, second, and higher approximations. Terms with viscosity, as well as the thermal terms, should be included into the equations of the various approximations, depending on the value of the dimensionless parameter ν/c (ν is the kinematic viscosity) with respect to v/c (v is the amplitude of the vibrational velocity). Particular solutions are found for the velocity of the second approximation, in two cases: (1) $\nu/c \gg v/c$ (the viscous terms enter into the equation of the first and higher approximations) and (2) $\nu/c \ll v/c$ (viscous terms do not enter into the equation of second approximation). In case (2) the solution coincides with the second approximation of the Riemann solution, and in spite of the presence of viscosity, the waveform has a tendency to accumulate the discontinuity. In case (1) the author determines the increase in the coefficient of absorption γ , due to the appearance of the second parameter (see ~~paper-text~~, p. 251).

$$\frac{\gamma}{\gamma_0} = 1 + \left[1 + \rho_0 \left(\frac{e^{-z}}{z} \right) \right] \left[1 - \frac{1}{2} \left(\frac{e^{-z}}{z} \right)^2 \left(\frac{d}{dz} \ln \left(\frac{e^{-z}}{z} \right) \right) \right]$$

Card : 2/2

AUTHOR: Gol'dberg, Z.A.

46-2-7/23

TITLE: Second order magnitudes in acoustics. (Nekotoryye
velichiny vtorogo polya v akustike)

PERIODICAL: "Akusticheskiy Zhurnal" (Journal of Acoustics), 1957,
Vol. 3, No.2, pp. 149-153 (U.S.S.R.)

ABSTRACT: The second order acoustical magnitudes have been, for
the ideal medium, investigated elsewhere (bibliography in (2)).
It is nevertheless of interest to investigate them for the
case of a viscous thermo-conducting medium. It has been
shown (1) that three particular cases need to be considered
(1), every one of them described by equations of the first
and of the second order approximation. In (3) expressions
of the second order approximation have been obtained for one
viscous medium. In the present article the author, using
the notation and terminology of (4) and (1) analyses mathema-
tically the solution, obtained in (3) as applied to a plane
sound wave for the above 3 cases (1). Second order solutions
are found and analysed for the hydro-dynamic velocity v_2 ,
pressure p_2 and density ρ_2 for the three following cases:
 $N \ll \omega/c^2$. It is shown that for this condition the ampli-
tude of v_2 is linearly proportional to the coefficient of

Card 1/2

Second order magnitudes in acoustics. (Cont.) 46-2-7/23

viscosity and to the frequency and that at a certain distance z_0 from the source, the second order magnitudes decay faster than the magnitudes of the first order. z_0 is called the "relative form stabilising distance". ρ_2 and p_2 are related by the usual relationships of linear acoustics. The second considered case is the condition $N \approx v_0/c^2$ and expressions for the same quantities are derived. It is shown that for a wave of infinite duration the amplitude of v_2 in this case increases for every point in time.

Card 2/2 The third condition is $N > v_0/c^2$ and second order solutions are given in eq.(29).

There are 5 Slavic references.

ASSOCIATION: The Magnitogorsk State Teaching Institute. (Magnitogorskiy Gosudarstvennyy Pedagogicheskiy Institut)

SUBMITTED: November 5, 1956.

AVAILABLE: Library of Congress

43-4-2/17

AUTHOR: Gol'dberg, Z.A.

TITLE: On the Propagation of Finite Waves of Finite Amplitude
(O rasprostranenii planarnih voln o omezhnoj amplitudoj)

JOURNAL: Akusticheskiy Zhurnal, 1959, V.11, N.4, p.390-396
(USSR)

ABSTRACT: The propagation of plane waves of finite amplitude in a viscous, thermally conducting medium is considered from the point of view of first and second approximation acoustics. A study is made of the criterion which indicates when the accumulation of discontinuities in a plane wave is possible. The distance from the source to the place of accumulation of discontinuities is estimated. Expressions are obtained for the absorption coefficient in the case where discontinuity is impossible as well as the case where it is possible. The above criterion is:

$$0.43 \frac{p^2}{b\omega} \leq 1 , \quad (1)$$

for air and for water the criterion is:

Card 1/2

46-4-3717

On the Frequency Dependence of the Absorption Coefficient.

$$A.A. \cdot \frac{1}{\omega^2} \leq 1. \quad (7)$$

(The notation is defined in Ref. 6-5 of which the present paper is a continuation). The present theoretical results are in agreement with the experimental results reported in Refs. 5 and 7. Thus in Ref. 5 a linear dependence of the absorption coefficient on frequency was observed beginning with $\omega' = 2$ atm. This is shown to be in agreement with the present results. According to Eq. 7 the increase in the coefficient of absorption is proportional to acoustic pressure in the case of a rigid absorber, and in this case it is natural to assume that ω' is the inductance of the waves, and t is again as predicted by the present theory. There are publications, of which 6 and 8 are English and 3 English.

ASSOCIATION: Institute for Applied Mathematics
(Institut für Angewandte Mathematik)

SUBMITTED: November 5, 1955.

AVAILABLE: Library of Congress.
Card 2/2

1. Waves-Propagation-Theory 2. Absorption 3. Acoustics

AUTHOR: *Ivanov, A.M.*

TITLE: On Propagation of Plane Sound Waves of Finite Amplitude in a Viscous Heat Conducting Medium (O propagatsii nelinejnykh zvukovykh voln v onechnoy sredy v vyzkoye teploprovodimoye srede)

PUBLICATION: Akusticheskiy Zhurnal, 1960, Vol. 6, Nr 1, pp 118-120 (USSR)

ABSTRACT: The author uses Lagrange's variables x , t to discuss an acoustic field produced in the region $x > 0$ by a plane vibrating along the x -axis at the point $x = 0$ under the following initial and boundary conditions: (I) - displacement of the particles in the medium is $u(x, t) = 0$ at $t \leq 0$; (II) - at $t > 0$ we have $u(0, t) = f(t) = a(1 - \cos \omega t)$; (III) - there are no reflectors at $x > 0$, i.e. only the wave moving from the vibrating plane in the direction of positive x is considered. The equation of motion is solved using the Krylov-Bogolyubov method (Ref 3). The expression obtained for the vibrational motion of the particles of the medium shows that in propagation of waves of finite amplitude their profile is distorted because of differences in velocities on various points of this profile. The wave amplitude decreases with

Zero 1/2

"On Propagation of Plane Sound Waves of Finite Amplitude in a Viscous Heat Conducting Medium"

distance. The paper is entirely theoretical. Acknowledgment is made to N.N. Andreyev and the participants of his seminar for their advice. There are 6 Soviet references.

ASSOCIATION: Akusticheskiy institut AN SSSR, Moscow (Acoustic Institute of the Academy of Sciences of the U.S.S.R., Moscow)

SUBMITTED: December 23, 1957

JAN 1/6

S/046/60/006/003/004/C12
B006/B063

AUTHOR: Gol'dberg, Z. A.

TITLE: Interaction Between Plane Longitudinal and Transverse
Elastic Waves

PERIODICAL: Akusticheskiy zhurnal, 1960, Vol. 6, No. 3, pp. 307-310

TEXT: The present article describes a theoretical investigation of the interaction between elastic waves in an unbounded, isotropic solid. The equations of motion (5) used for this purpose take account of both the linear terms and the terms which are quadratic with respect to the derivatives of the deformation vector u . For the special case of plane waves the general system (5) can be transformed into the system (6) - (8). In a linear approximation, the latter system consists of three independent wave equations for u_x , u_y , and u_z . This means that longitudinal and transverse waves propagate without affecting one another. The terms which are quadratic in du_i/dx_k depend on all components of the displacement vector. Accordingly, one obtains an interaction of the two kinds of waves only. ✓C

Card 1/3

Interaction Between Plane Longitudinal and
Transverse Elastic Waves

S/046/006/003/004/012
B006/2063

in second approximation. A series of peculiarities appears in this connection. A consideration of the propagation of a transverse wave only shows that also a longitudinal wave occurs, whereas vice versa, during the propagation of a longitudinal wave, no transverse wave appears. The shape of the longitudinal wave changes during its propagation, while the transverse wave remains unchanged. These results are finally discussed. N. N. Andreyev and the participants in the author's seminar are thanked for their valuable remarks. There are 4 references; 2 Soviet and 2 US.

ASSOCIATION: Magnitogorskiy gosudarstvennyy pedagogicheskiy institut
(Magnitogorsk State Pedagogical Institute)

Card 2/3

Interaction Between Plane Longitudinal and
Transverse Elastic Waves

S/046/60/006/003/004/012
E006/B063

SUBMITTED: September 28, 1959

✓

Card 3/3

"APPROVED FOR RELEASE: Thursday, September 26, 2002

APPROVED FOR RELEASE: Thursday, September 26, 2002

CIA-RDP86-00513R000515620006-6"

GOLDBERG, Z. A. and NAUGOLNYKH, K. A.

"On the radiation pressure of standing waves"

report submitted for the 4th Intl. Congress of Acoustics,
Copenhagen, Denmark, 21-28 Aug 1962.

24,2500 (1057)

Re: 5
S/CHB/62, 147/1.1/336, C43
B102/B106

AUTHOR: Gel'fand, I. M.

TITLE: Waves with finite amplitude in magnetohydrodynamics

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, v. 41,
no. 1, 1961, pp. 1 - 28

TEXT: I. M. Gel'fand and N. B. Slobodcikov (DAN SSSR, 11, 74, 1959) have solved the problem of one-dimensional traveling wave propagating transverse to a magnetic field. The solution obtained is valid as long as no discontinuities arise. A similar problem is now considered in magnetohydrodynamics for weak shock waves or waves emitted by a harmonically vibrating plane. The waves are assumed to propagate across the field in a viscous conducting medium. The system of one-dimensional magneto-hydrodynamic equations given by L. D. Landau and Ye. M. Lifshits (Elektrodinamika sploshnykh sred - Electrodynamics of continuous media - Gostekhizdat, 1957) is approximately rewritten in Lagrange variables and solved neglecting terms smaller than certain other terms. The solutions permit studying the development of a shock wave as well as estimating the

✓

Card 1/3

Journal of the American Mathematical Society

1996-1997

duration of this process. Also the formation during propagation of an initially clumped field wave can be studied. In the latter case, for a wave field amplitude $\sim \sin(\omega_0 t)$ given in the plane $x = 0$,

$$v(a,t) = \frac{b_{00}}{\rho_0 u_0^2 r} \sum_{n=1}^{\infty} \frac{\sin(n\phi(t-a/r))}{\sin(n(\mu + 1/2R))}, \quad (37),$$

$$b_0 + \frac{1}{3} \Psi_0^2 + \frac{\pi}{8} + \text{Z}_0 \left(\frac{1}{c_p} - \frac{1}{c_n} \right) \frac{a_0^2}{a^2} = -\frac{c_0^2 H_0^2}{16 \pi^2 m^2}, \quad (12),$$

$$\epsilon = 1 + \rho_0 (\partial^2 p / \partial \mu^2)_{\mu=0} / 2u_0^2 + 3H_0^2 / 8\pi\rho_0 u_0^2.$$

is obtained. $R = \beta x_{\mu_0} u_{\mu_0}^2 / m$; $\tilde{V} = (v_1, \dots, v_n)$; $\tilde{U} = (u_1, \dots, u_n)$.

$$h = \frac{H}{u} v, \quad F = \left(\frac{\partial T}{\partial p} \right)_U p u_0^2, \quad \frac{\partial}{\partial t} = - H \frac{\partial}{\partial u}; \quad u = \sqrt{v^2 + \frac{H^2}{u_0^2}} e^{-\frac{H^2}{2u_0^2}t}, \quad v = \sqrt{v^2 + \frac{H^2}{u_0^2}} e^{-\frac{H^2}{2u_0^2}t}.$$

The result shows that at a distance of the order of λ_0 the sinusoidal wave has turned into a sawtooth wave and in further propagation becomes a simple sinusoidal wave. There are 1 figure and 10 references: 5 Soviet Card 2/7

2115

Journal of Health Politics, Policy and Law, Vol. 35, No. 4, December 2010
DOI 10.1215/03616878-35-4 © 2010 by The University of Chicago

3-1866-62-042-02-0356-146
3-1866-62

and so on. The fact that recent references to duality in harmonic analysis are based on first last J. B. Monticinos, *J. Math. Soc. Asia*, 1970, 1971; M. A. Ramanujan, *Surveys in Mathematics*, Cambridge University Press, 1971; E. Hille, *Functional Analysis*, Pure Appl. Math., 1973, 1973.

ДИАГНОСТИКА Наши клиники предлагают диагностические центры для диагностики и лечения онкологии.

2016-2017 学年第二学期期中考试

Chart 3 -

9/046/63/009/001/005/026
B104/B106

AUTHORS: Gol'dberg, Z. A., Naugol'nykh, K. A.

TITLE: Rayleigh sound pressure

PERIODICAL: Akusticheskiy zhurnal, v. 9, no. 1, 1963, 28-31

TEXT: The results of Rayleigh (Phil. Mag., 1905, 10, 564-574) obtained for the sound pressure on a fixed rigid wall for the case of a medium vibrating between two fixed plane rigid boundaries are generalized for a forced vibration of the medium produced by harmonical motion of one of the two boundaries. In linear approximation of the sound field the radiation pressure of a standing wave on the fixed boundary is

$$\overline{P} = \frac{\gamma + i}{i} \rho_0 v_0^2 \left(1 + \frac{\sin 2kt}{2kt} \right), \quad (9),$$

where $v_0 = A\omega/\sin(kl)$ is the particle velocity, γ is the mean shift of the particles from their equilibrium position, A is a constant, k is the

Card 1/2

Rayleigh sound pressure

S/046/63/009/001/005/026
B104/B186

wave number, ω the angular frequency. If $k_1 \ll 1$ or $k_1 \rightarrow n\pi$ (resonance), v_0 and with it \bar{p}' increases unlimitedly. In the case of spherical standing waves produced between two concentric spheres by vibration of the inner sphere the pressure at the unmoved outer sphere is

$$\bar{p}'(R_2) = p_0 \frac{A^0 k^3}{4} \frac{\sin^4 kR_1}{R_1^4} + c_1. \quad (16).$$

This formula is specialized for a quiet inner sphere and a vibrating outer sphere, for a zero radius of the inner sphere and for a standing wave between non-vibrating spheres.

ASSOCIATION: Akusticheskiy institut AN SSSR, Moskva (Institute of Acoustics AS USSR, Moscow)

SUBMITTED: June 25, 1962

Card 2/2

"APPROVED FOR RELEASE: Thursday, September 26, 2002
CIA-RDP86-00513R000515620006-6"

1. Description of the individual, group, or organization involved.

Individuals of the same name, age, sex, and ethnicity as the target.
For example, "John Doe, Jr."

2. Description of the individual, group, or organization involved.
For example, "John Doe, Jr."

ACC NR: AP6034020

SOURCE CODE: UR/0226/66/000/010/0071/0077

AUTHOR: Tumanov, V. I.; Gol'dberg, Z. A.; Chernyshev, V. V. Pavlova, Z. I. (Deceased)

ORG: All-Union Scientific Research Institute of Hard Alloys (Vsesoyuznyy nauchno-issledovatel'skiy institut tverdykh splavov)

TITLE: Thermal stability of alloys of tungsten-cobalt carbides

SOURCE: Poroshkovaya metallurgiya, no. 10, 1966, 71-77

TOPIC TAGS: thermal shock simulation, heat resistant alloy, tungsten carbide, cobalt, bend strength, grain size, grain structure, hardness

ABSTRACT: Thermal shock testing of alloys of tungsten-cobalt was made by water quenching samples from temperatures up to 1120°K. The furnace capacity was sufficiently great to test 20-40 samples simultaneously. Specimens were held 5 min in the furnace and 0.5 min in the quenching bath. Thermal shock stability was measured in terms of superficial cracks and the decrease in ultimate bend strength after thermal cycling. The cobalt content of the samples ranged from 1 to 30 wt %, while some samples containing 20-30% cobalt were alloyed with 0.6 or 2.1% titanium, chromium, or molybdenum. The porosity did not exceed 0.2 vol %. The first set of experiments was conducted on 8 × 5 × 35 mm samples quenched from 770°K. Thermal shock resistance increased sharply above 15% Co. Up to 6% Co the number of thermal shock cycles needed to induce macro-

Card 1/2

ACC NR: AP6034026

cracks was 7 or less; at 15 to 30% Co no cracking was observed after 500 cycles. Small grained samples had a lower thermal shock stability. A microstructural analysis was made on samples with 25% Co, alloyed with either Ti, Cr, or Mo. The volume distribution of grain size was given for the different alloys, before and after 300 cycles of thermal shock testing. The ultimate bend strength of 2 × 5 × 35 mm samples, quenched from 1120°K, is given as a function of the number of cycles. The greatest drop in strength occurred after 100 cycles. Alloy VK20 (20% Co) had the highest bend strength while VK30 (30% Co) had the lowest for all thermal shock cycles, ranging up to 500. The effect of thermal cycling on Vickers hardness was negligible. It is concluded that the mechanism of strength decrease during thermal cycling is associated with fine structural changes, which could not be observed by the techniques described above.
Orig. art. has: 2 figures, 4 tables.

SUB CODE: 11/ SUBM DATE: 04Apr64/ ORIG REF: 003/ OTH REF: 005

Card 2/2

GOL'DBERG, Z.N., inzh.

Conference on the mechanization and automation of coal mining.
Bezop.truda v prom. 3 no.4:34-36 Ap '59. (MIRA 12:6)
(Coal mines and mining)

GOL'DBURG, Z.N., inst.

All-Union conference on reducing air dustiness in mines. Bezp.
truda v prom. 5 no.4:35-36 Ap '61. (MTRA 14:3)
(Mine dusts--safety measures)

GOL'DBERG, Z.N., inzh.

A skillful organization of work is the basis of success. Bezop. truda
v prom. 6 no.3-3 4 Mr '62. (MRA 15:3)
(Donets Basin--Coal mines and mining)

GOL'DBERG, Z.N., inzh.

Scientific technical conference on safety engineering in the coal
mining industry. Bezop.trud v prem. o no.4:37-38 Ap '62.
(MIRA 15:5)
(Coal mines and mining--Safety measures)

1. Optical Cell, 1000 nm.
2. Ozone (100)
3. Glass, Variable
7. Spectroscopic investigation of a variable variable filter. [unclassified] 1953. NAV. Astron. Obs., U.S.A.F., Eng.
9. Monthly List of Russian Accessions, Library of Congress, [unclassified] 1953. Unclassified.

KRAT, V.A.; GOL'DBERG-ROGOZINSKAYA, N.M.

Investigating granulations of the sun's photosphere. Part 2.
Izv.GAO 20 no.2:17-21 '56. (MIRA 13:5)
(Sun)

GOL'DBERG-ROGOZINSKAYA, N.M.

Spectrophotometric investigation of the eclipsing variable RS
Vulpeculae. Izv.GAO 20 no.2:61-73 '56. (MIRA 13:5)
(Stars, Variable)

GOL'DBERG-ROGOZINSKAYA, N.M.

Determining photometric standards in areas of the Pulkova
Catalog of extragalactic nebulae. Izv.GAO 21 no.3:94-158.
(MIRA 1):4)
(Nebulae)

L-45334-50 EMT(1)/EMI(m)/EMI(t)/ETI 13P(c) SOURCE CODE: UR/0269/65/000/012/0051/0051
ACC NR: AR6015216

AUTHOR: Gol'dberg-Rogozinskaya, N. M.

TITLE: Helium lines in the spectra of chromospheric flares

SOURCE: Ref. zh. Astronomiya, Abs. 12.51.399

REF SOURCE: Izv. Gl. astron. observ. v Pulkove, v. 24, no. 2, 1965, 35-40

TOPIC TAGS: chromosphere, helium, spectrum, chromosphere flare, solar flare, ionized helium, neutral helium

ABSTRACT: An article of the same title by the author (Goldberg-Rogozinskaya, N. M., R. Zh. Astr, 1963, 2.51.429) is continued. Lines of neutral helium $\lambda\lambda 4471, 4713, 4922, 5016, 5876$, and a line of ionized helium $\lambda 4686$ were observed during three flares of 2 April, 30 April and 6 August 1960. The line profiles are wide and nonsymmetrical. Real interior movements explain the profiles observed better than do thermal processes. The electronic density n_e and the electronic temperature T_e in the flares is evaluated by the intensities of the lines. In the case of neutral

47
13

Card 1/2

UDC: 523.75

ACC NR: AR6015216

helium, $n_e = 10^{10}$, $T_e = 30,000K$, and in the case of ionized helium, $n_e = 10^{10}$
and $T_e = 50,000K$. [Translation of abstract] [GC]

SUB CODE: 03,20/ SUBM DATE: none/

Card 2/2

GOLDBERGEN

LUPASCU, Gh., membru coresp. al Academ. RPR; AGAVRILOAEI, A.; COSTIN, P.; ELIAS, M.; ZELIG, M.; RADCOV, G.; FEODOROVICI, St.; GOLDBERGEN, E.; SZABO, M.; STANCULESCU-ROSIU, I.

Study of pappataci fever. Bul. stiint. sect. med. 8 no.1:
265-275 Jan-Mar 56.

(FEVER

pappataci fever, epidemiol. & prev. in Rumania.)

PLATE 23

卷之三

Clinical Examinations, Hospital Patients
Number Participating (estimated) Clinical Examinations Participating, 1947
657 P. Clinic and University, No. of copies printed not stated. No.
Diagnosis: Data base is intended for management, physical therapy, exercise, and
social and vocational guidance.

Comments: The base consists of 10,000 by hospital patients admitted to the
institution, with approximately 10,000 additional patients, namely,
menial labor, etc., and their dependents. Patients in hospitals and
clinics or clinics are given as the total of each service. Data of the services
are summarized by performance. The general hospital and institutions to which the data
relate are not given.

Table of Contents:

Author, C. R.	CONTINUOUS DENSITY DISTRIBUTION LAW FOR THE WEIGHTS OF ADULTS. V. A STUDY OF ESTIMATION AND P. P. PLOTS. AN INVESTIGATION OF THE PROBLEMS OF ESTIMATING THE WEIGHTS FROM THE DISTRIBUTION OF INDIVIDUALS ON THE BASIS OF PREDICTED DENSITY	277
Dale, F.	DEPOSITIONS AND PROOFS IN THE POLICE COURT CASE	277
Dale, F. and G. Dean	A MORE PRACTICAL WAY OF SOLVING THE RELATED PROBLEMS INVOLVED IN THE POLICE COURT CASE	278
Dale, F. and G. Dean	THE RELATIONSHIP OF THE DEFENDED PERSON TO A PROSECUTED SUSPECT	279
Nikolsky, M.	ON WHICH DEAL CONDITIONS THREE-DIMENSIONAL APPROXIMATIONS BETWEEN THE FUNCTIONS OF TWO VARIABLES CAN BE APPLIED TO THE WAVE EQUATION	280
Rudnitskii, M.	COMPARISON OF MEASURED RESULTS WITH COMPUTED RESULTS OF THE PROPAGATION OF A ROLLING WAVE	281
Semenov, I.	DETERMINATION OF RADIAL ILLUMINATION BY MEANS OF LUMINES- CENT MATERIALS	282
Semenov, I. and G. Mikhlin	INFLUENCE OF THE POSITION OF THE POINT OF OBSERVATION ON THE RADIAL DISTRIBUTION OF LIGHT	283

PROTIC, Mihajlo F.; GOLDBERGER, Aleksandar

Rectal injuries. Med. Glasn. 8 no.1:11-14 Ja '54.
(RECTUM, wds. & inj.)

*
(WOUNDS AND INJURIES
*rectum)

PROTIC, Mihalio, F., Asist., Doc., Dr.; GOLDBERGER, Aleksander, dr.

Gallbladder perforation as a complication of typhoid fever. Med.
arh., Sarajevo 10 no.5:81-85 Sept-Oct '6.

1. Hirurska klinika Medicinskog fakulteta u Sarajevu. Sef: prof.
dr. Blagoje Kovacevic.

(TYPHOID FEVER, compl.

perf. of gallbladder, surg. (Ser))

(GALLBLADDER, perf.

in typhoid fever, surg. (Ser))

GOLDBERGER, Aleksandar, dr.

Surgery of the thyroid gland. Med. glasn. 14 no. 2a : 135-140
F '60.

1. Hirurško odjeljenje Opste Spomen-bolnice "Proleterskih
brigada" u Foci, Unravnik: dr A. Godberger.
(THYROID GLAND surg.)

GOLDBERGER, A.; HEGYMEGI KISS, P.; SZOKE, G.; TEGEDI, I.

Significance of bacilli secretion in infantile tuberculosis. Gyermekgyogyaszat. 2 no.9:273-276 Sept 51. (CIML 21:1)

1. Doctors. 2. Szabadsaghegyi State Children's Sanatorium (Director Head Physician--Dr. Istvan Flesch).

HOFFMANN, I.; GOLDBERGER, A.; SZOCSKA, M.; SZOKE, G.

Streptomycin therapy of infant tuberculosis. Gyermekgyogyaszat 4 no.4:
119-121 Apr 1953. (CIML 24:4)

1. Doctors. 2. Szabadsaghegyi State Children's Sanatorium (Director --
Head Physician -- Dr. Istvan Flesch).

TELEGDI, Istvan, dr.,; TOTH, Eva, dr.,; GOLDBERGER, Arpad, dr.,; TEGE,
Antal, dr.

Data on the anamnesis of children with pulmonary tuberculosis.
Otv. hetil. 97 no.6:153-155 5 Feb 56.

1. A Szabadsaghegyi Allami Gyermekszanatorium (igazgato-foorvos:
Flesch Istvan dr.) Primer tbc osztalyanak (foorvos:Telegdi Istvan
dr.) kozlemenye.

(TUBERCULOSIS, PULMONARY, in inf. & child
anamnestic data on 1100 child. (Hun))

GOLDBERGER, A.

Endemic struma. Bul s: Yng 7 no.6 173 D '62.

1. Medicinski fakultet, Sarajevo

#

GOLDBERGER, M. L.

Weak interactions. Postepy fizyki 12 no.4:403-413 '61.

GOLDBERGS, J., red.; PASTARE, D., tekhn. red.

[Soviet Baltic Republics in the fraternal family of Soviet peoples; materials of the interrepublic conference] Baltijas Padomju republikas PSRS tautu brālīgāja saime; notikusas starprepublikaniskas apspriedes materiāli. Riga, Latvijas Valsts izdevniecība. [In Latvian]. Vol.1. 1960. (MIRA 15:1)

1. Mezhrespublikanskiy seminar--soveshchaniye na temu "Sovetskaya Pribaltika v bratskoy sem'ye narodov SSSR, Riga, 1960.
(Latvia--Economic conditions)

GOLDBERGS, J., red.; ERENSTEINE, A., tokhn. red.

[Soviet Baltic Republics in the fraternal family of Soviet peoples; materials of the interrepublic conference] Baltijas Padomju republikas PSRS tautu brālīgāja saime; notikusas starp-republikaniskas apspriedes materiali. Riga, Latvijas Valsts izdevniecība. [In Latvian] Vol.3. 1960. (MIRA 15:1)

1. Mezhrespublikanskiy seminar-soveshchaniye na temu "Sovetskaya Pribaltika v bratskoy sam'ye narodov SSSR, Riga, 1960.
(Latvia—Economic conditions)

TURSEVICS, V.; GOLDBECK, J., red.; STIKALIS, A., tekhn. red.

[Let us mobilize hidden potentials] Mobilizesim iekcejas rezerves. Riga, Latvijas valsts izd-va, 1962. 46 p.
(MIRA 17:2)

GOL'DBERT, S. V.

Gol'dbert, Z. V. and Peterson O. P. "Histopathological changes in white mice upon the introduction of large doses of grippa virus into the stomach", Voprosy med. virusologii, Issue 1, 1945, p. 190-97, - Bibliog: 7 items.

SO: U-3042, 11 March 53, (Letopis zhurnal 'nykh Statey, No. 10, 1945).

GOL'DBETS, S. B.

USSR/Engineering - Refractories, Production Feb 52

"On Utilization of Chrome Ore Tailings Obtained in Fabrication of Chromomagnesite Brick," S. B. Gol'-dbets, Engr, Plant imeni Petrovskiy

"Ogneupory" No 2, pp 85-87

Describes expts for using chrome ore tailings in fabrication of chromite bricks or as chromite component in chromomagnesite bricks. In latter case, briquettes of chrome ore are preliminarily burned to 1,520-1,580° and crushed to fine powder. Results show that utilization of ore tailings is quite worth while. Gives physicochem characteristics of both types of brick.

204T21

APPROVED FOR RELEASE: Thursday, September 26, 2002

CIA-RDP86-00513R000515620006-6

CIA-RDP86-00513R000515620006-6"

220500
SACI, R. T. 141145Z, 10-1.

"gas-linder automobile," dated 1900, 1 copy. Description of
system used in USSR of use of natural gas or coal gas as fuel for
trucks.

SAMOL', G.I., kandidat tekhnicheskikh nauk; GOL'DBLAT, I.I., kandidat
tekhnicheskikh nauk; KOLOSOV, V.A., kandidat tekhnicheskikh nauk,
redaktor; POPOVA, S.M., tekhnicheskiy redaktor

[Gas cylinder automobiles] Gazoballonnye avtomobili. Izd. 2-e,
perer. Moskva, Gos. nauchno-tekhn. izd-vo mashinostroit. i sude-
stroit. lit-ry, 1953. 284 p. (MLRA 7:9)
(Automobiles--Engines (Compressed gas))

GOL'DBLAT, I.I.
SAMOL', Grigoriy Ivanovich, kandidat tekhnicheskikh nauk; GOL'DBLAT, Il'ya
Isaakovich, kandidat tekhnicheskikh nauk; GRUZINOV, V.I., redaktor;
MAL'KOVA, N.V., tekhnicheskiy redaktor

[Gas cylinder automobiles; chauffeur's manual] Gazoballonnye avto-
mobilii; posobie dlya shofera. Moskva, Avtotransizdat, 1954. 86 p.
(Automobiles - Engines (Compressed-gas)) (MLRA 8:4)

SAMOL', Grigoriy Ivanovich; GOL'DBLAT, Il'ya Isaakovich; IGUL'KIN, V. N.,
redaktor; MAL'KOVA, N.V., tekhnicheskij redaktor.

[Booklet on safety measures while driving compressed-gas auto-
mobiles] Pamiatka po tekhnike bezopasnosti pri rabote na gazo-
ballonnom avtomobile. Moskva, Nauchno-tehn. izd-vo avtotrans-
portnoi lit-ry, 1955. 45 p. (MLRA 8:8)
(Automobiles--Safety measures)

GOL'DBLAT, I.I., kandidat tekhnicheskikh nauk.

Engines for gas cylinder automobiles. Avt.i trakt.prom.no.11:8-11
N '56. (MIRA 10:1)

1. Nauchno-issledovatel'skiy avtomotornyy institut.
(Automobiles. Engines. (Compressed gas))

GOL'DELAT, I.I.

Isothermal tank for low-temperature storage of propane. Gaz.prom.
no.12:35-36 D '57. (MIRA 11:1)
(Propane--Storage)

SAMOL', Ivan Ivanovich, kand. tekhn. nauk; GOL'DBLAT, Il'ya Isaakovich,
kand. tekhn. nauk; DIVAKOV, N.V., red.; NIKOLAYEVA, L.N., tekhn.
red.

[Motortrucks operated with liquified gas; manual for drivers] Gazo-
ballonne avtomobili; posobie shoferu. Moskva, Nauchno-tekhn. izd-
vo M-va avtomobil'nogo transp. i shosseinykh dorog RSFSR, 1961. 141 p.
(MIRA 14:6)

(Motortrucks—Motors (Compressed gas))

SAMOL', G.I., kand. tekhn. nauk; GOL'DBLAT, I.I., kand. tekhn.
nauk; KISELEV, V.Z., inzh., retsevzent; VASIL'YEVA, I.A.,
red. izd-va; EL'KIND, V.D., tekhn. red.

[Gas cylinder automobiles] Gazoballonnye avtomobili. Izd.3.
Moskva, Mashgiz, 1963. 386 p. (MIRA 16:5)
(Automobiles--Engines (Compressed gas))

1. СИДОРЧУК, В. П.
2. УССР (СССР)
4. Тривалій період
7. Тривалій період від пульсальної туберкульозу. Пробл. мед., №. 1, 1953.

9. Monthly List of Russian Accessions, Library of Congress, Sept 1953, Unclassified.