

GOLDENVEZER, A. L.

"On the Application of the General Laws of the Theory of Elasticity to Thin Shells," Prik Mat i Mekh. Vol. 8, No. 1, 1944, pp. 3-14

GOLDENVEYER, A. L.

"Investigation of Spherical Shells under a State of Strain," Prikl Mat i Mekh
Vol. 8, No. 6, pp. 441-467, 1944

GOL'DENWEYZER, A. L.

"Qualitative Investigation of the State of Tension of Thin Shells," Prikl Mat i Mekh, Vol. 9, No. 6, 1945, pp. 464-473

GOL'DENVEYZER, A. I.

"On the Integration of a System of Differential Equations of the Theory of Thin Shells," Report at the Meeting on the Theory of Elasticity, Building Mechanics and Plasticity, 25-28 March 1946. Published in the Doklady of the Meeting.

GOLDENVEYER, N. L.

Gelevenweizer, A. I. [P]roblems of integration of equations of the theory of thin shells. *Appl. Math. Mech. [Akad. Nauk SSSR, Izd. Mat. Mosh.]* 10, 387-395 (1946). (Russian, English summary)

The author gives a method of determining the edge effects along those contours of the shell whose tangents do not coincide with the asymptotic lines of the middle surface. The integration of the system of equations of the above theory is reduced, in several cases, to the integration of inmomentless theory. This paper is a sequel to the author's earlier paper [same journal 9, 465-478 (1945); these Rev., 7, 351].
I. S. Sokolnikoff (Los Angeles, Calif.)

Source: Mathematical Reviews,

Vol. 8, No. 4

220

GOL'DEWSYER, A. L. D_v. Physicomath Sci.

Dissertation: "Qualitative Investigation of the Equations of the Theory of Thin Shells and Certain Methods of Their Integration." Inst. of Mechanics, Acad. Sci. USSR, 25 Feb 47.

SO: Vechernaya Moscow, Feb, 1947 (Project #17636)

GOLDENWEISER, A. L.

Goldenweiser, A. L. Momentless theory of shells whose middle surface is of a curve of the second order. *Appl. Math. Mech. [Akad. Nauk SSSR. Fiz.-Mat. Mekh.]* 11, 285-299 (1947). (Russian. English summary)

[A more accurate translation of the Russian title would omit the phrase "of a curve." This note indicates that the solution of the problem of momentless shell theory for shells whose middle surfaces are quadrics is reducible, by a suitable change of dependent and independent variables, to the integration of Poisson or wave equations.

I. S. Sosulinoff (Los Angeles, Calif.)

Source: Mathematical Reviews, 1948, Vol. 9, No. 2

GOLDENWERTHER, A.

Goldenwérther, A. L. Approximate calculation of thin shells of zero Gauss curvature. Akad. Nauk SSSR, Pril. Mat. Meh. 11, 409-422 (1947). (Russian. English summary)

The main object of this paper is a qualitative analysis of stressed states in thin elastic shells with developable middle surfaces. The paper also contains an outline of the methods of approximate calculation of stresses. The shell is covered by a net of lines of curvature α, β (the α -lines are of zero curvature) so that the first fundamental form for the surface is of the type $ds^2 = d\alpha^2 + B^2 d\beta^2$. In this case Love's general equations of the shell theory are reducible to two differential equations for the stress functions t and m from which the forces, moments, and deformations can be computed by differentiation. These equations are:

$$\frac{\lambda^2}{B^2} \frac{\partial}{\partial \alpha} \frac{\partial t}{\partial \alpha} - \frac{\lambda^2 h}{3(1-\sigma^2)} N(m, \sigma) = 0,$$

$$\frac{\lambda^2}{B^2} \frac{\partial}{\partial \alpha} \frac{\partial m}{\partial \alpha} + \lambda^2 N(t, -\sigma) = 0,$$

where λ^2 and N are introduced to make the terms of these equations have the dimensions of t ; for cylindrical shells ($\partial B/\partial \alpha = 0$),

$$N(F) = \frac{1}{B} \frac{\partial}{\partial \beta} \frac{\partial}{\partial B} \frac{\partial}{\partial \beta} \frac{1}{B} \frac{\partial}{\partial \beta} \frac{1}{B} \frac{\partial F}{\partial \beta} + \frac{1}{BR} \frac{\partial}{\partial \beta} \frac{1}{B} \frac{\partial F}{\partial \beta}$$

R being the radius of curvature of the β -line; for noncylindrical shells ($\partial B/\partial \alpha \neq 0$) $N(F)$, in addition to the terms

Source: Mathematical Reviews, 1948, Vol. 9, No. 4

given above, contains the term

$$\frac{1}{B} \frac{\partial}{\partial \beta} \frac{1}{B} \left(\frac{\partial}{\partial \alpha} \frac{1}{B} \frac{\partial}{\partial \alpha} \right) \frac{\partial F}{\partial \beta}.$$

For conical and cylindrical shells the system can be integrated approximately in the form of a series involving trigonometric and Bessel functions provided certain restrictions on the lengths of the shells and on the generatrix angle are imposed.

Several results obtained in the author's two earlier papers [same journal 9, 463-478 (1945); 10, 387-396 (1946); these Rev. 7, 391; 8, 241], dealing with thin shells of zero Gaussian curvature which are so stressed that the state of stress can be decomposed into a membrane state and into a state produced by moments and boundary effects, appear as special cases in this more general treatment.

I. S. SAVENKOFF (Los Angeles, Calif.).

GOLDENVEYER, A. L.

Gol'denveizer, A. L. and Lur'e, A. I. On the mathematical theory of the equilibrium of elastic shells. (Survey of the work published in the USSR.) Acad. Nauk SSSR. Prikl. Mat. Meh. 11, 565-592 (1947). (Russian)

This is a condensed survey of the research literature on the subject published in Russia during the past decade. Three distinct directions are discernible: (a) theoretical investigations based on the fundamental equations of the mathematical theory of elasticity; (b) works on stability and vibrations; (c) papers concerned with the engineering applications of the theory. This survey is concerned only with the first aspect. The development surveyed in this article falls into three categories: (a) formulation of the basic equations of the theory of thin shells, which extends the classical theory of Love with the aid of modern tools of differential geometry; (b) specialization of the general three-dimensional problem of elasticity to a two-dimensional one by introducing certain geometrical hypotheses; (c) physical assumptions. Papers in this category are concerned with the analysis of the nature of the simplifications and with the study of the magnitude of errors inherent in the interpretation of the equations formulated in category (a). This is accomplished by replacing the somewhat abstract equations by special forms yielding the pertinent information about the edge effect and the behavior of shells under various loads.

CIA-RDP86-00513R000515620015-6
CIA-RDP86-00513R000515620015-6

(1) interpretation of the equations formulated in category (a). This is accomplished by replacing the somewhat abstract equations by special forms. These include:

(2) interpretation of the equations of the theory of shells associated with specific geometrical forms. These include:
cylindrical shells, conical shells, and shells with vanishing Gaussian curvature. The survey concludes with a bibliography of 48 items. J. S. Shabotoff (Los Angeles, Calif.).

Source: Mathematical Reviews,

Vol.

No.

SAC

GOL'DENVEYER, A. I.

"The Influence of Border Fastening on the State of Stress of Thin Shells,"
Trudy of the Central Aero-Hydrodynamic Institute (ZAGI) 1948, No. 669

USSR/Engineering

Mechanics

Bibliography

Jan/Feb 49

"Review of V. V. Novozhilov's 'Theory of Thin Shells,'"
A. L. Gol'denveizer, 3 pp

"Priklad Matemat i Mekh" Vol XIII, No 1

Generally favorable review of subject book, which
attempts to classify and clarify accumulated data
on the theory of thin-walled shells.

39/49T43

USSR/Engineering - Rods
Shells

Nov/Dec 49

"On the Theory of Thin-Walled Rods," A. L. Gol'den-
veyzer, Moscow, 35 pp

"Prikl. Matemat. i Mekh." Vol. XIII, No 6 - p. 561-74

This work differs from others on thin-walled rods in that it does not make use of special hypotheses based on qualitative analysis of integrals of equations as found in the theory of shells. Purpose of its investigation is to determine approximately the basic stressed state in a rod span loaded by transverse load R and system of forces and moments T applied to terminal transverse section. It is

USSR/Engineering - Rods (Contd) Nov/Dec 49

assumed that terminal sections of the rod are fixed arbitrarily and longitudinal sides are free of bonds. Submitted 21 Jun 49.

APPROVED FOR RELEASE: Thursday, September 26, 2002
APPROVED FOR RELEASE: Thursday, September 26, 2002

CIA-RDP86-00513R000515620015-6
CIA-RDP86-00513R000515620015-6"

153755

GOL'DENVEYZER, A. I.

PA 153755

GOL

"APPROVED FOR RELEASE: Thursday, September 26, 2002
APPROVED FOR RELEASE: Thursday, September 26, 2002

CIA-RDP86-00513R000515620015-6

CIA-RDP86-00513R000515620015-6"

I. N. Vekua, and A. I. Lur'e.

The mathematical theory of the propagation of elastic waves. (A review of papers published in the USSR). New York, Pergamon. (American Mathematical Bulletin, 1958, v. 55.)

Bibliography: p. 11-12.

Pravo i vlastatel'stvennye teorii ravnopravija upravlenija obmenem.
DRASKOVICH, V. N. et al.

SO: Aeronautical Sciences and Aviation in the Soviet Union, Library of Congress, 1955.

177T47

USSR/Mathematics - Shells, Equilibrium of Mar/Apr 51

"Applying the Solution of the Riemann-Hilbert Problem
to Computation of Momentless Shells," A. L. Golden-
veiser, Moscow

"Prik Matemat i Mekh" Vol XV, No 2, pp 149-166

Applied to 2d-order surfaces of pos curvature in
cases where moments may be neglected. In this case
tangential forces are computed by integrating eq of
equil. Momentless shell is statistically detd only
in definite boundary cases.

177T47

PHASE I

TREASURE ISLAND BIBLIOGRAPHICAL REPORT

AID 378 - I

BOOK

Author: GOL'DENVEYZER, A. L. Call No.: QA935.G6

Full Title: THE THEORY OF ELASTIC THIN SHELLS

Transliterated Title: Teoriya uprugikh tonkikh obolochek

Publishing Data

Originating Agency: None

Publishing House: State Publishing House of Engineering and
Theoretical Literature

Date: 1953

No. pp.: 544

No. of copies: 4,000

Editorial Staff

Editor: None

Tech. Ed.: None

Editor-in-Chief: None

Appraiser: None

Text Data

Coverage: The theory of shells as based on the assumption of the inalterability of the normal element is considered in this book. It is further assumed that the materials are isotropic and obey Hook's law generalized, and that the second powers of deformations, displacements, and angles of return are sufficiently small to be neglected. The author made an effort to present as completely as possible the many existing approximate methods of calculation of shells. The book is the result of many years of the author's research. It is divided into five parts, each part being a complete entity which may be studied separately.

1/10

sections; 11. External loads; 12. equilibrium equations....

2/10

Journal of the American Ceramic
Society
Vol. 37 No. 4
Apr. 1, 1954
Cements, Limes, and Plastics

Autoclave method of making asbestos-cement shingles. T. M. BERKOVICH, I. L. RABINOV, AND V. L. GOT'DENZLER. *Cement*, 19 [4] 19-23 (1953).—In the existing method of making asbestos-cement shingles, high-grade Portland cement is used as the bond. The shingles are steamed at 50° to 60°C. for 8 to 16 hr. and then hardened in storage for 7 to 10 days. An improvement of this method involves the addition of not less than 50% finely ground quartz sand to the cement and steaming in an autoclave at 8 atm. pressure for 8 hr.

B.Z.K.

GOL'DENVEYZER, A. L.

Gol'denveizer, A. L. On the calculation of shells with concentrated forces. Akad. Nauk SSSR, Prikl. Mat. Meh. 18, 181-186 (1954). (Russian)

There are two methods of calculating shells on concentrated forces. The first one starts with a distributed load acting in a small region which is allowed to shrink to a point, the load accordingly increasing infinitely at the same time. The second method consists of constructing a function satisfying the elasticity differential equations which has a certain defined singularity in the neighborhood of the point of application of the concentrated force. The author considers the second method only, which is mathematically very convenient, but which can be used only if the nature of the singularity is known beforehand. The author uses the following singularity: $r^1 \ln r$.

T. Leser.

gyp

SUBJECT USSR/MATHEMATICS/Differential equations CARS 1/2 PG - 490
 AUTHOR GOL'DENVEYZER A.L.
 TITLE An improvement of the theory of the simple edge effect.
 PERIODICAL Priklad.Mat.Mech. 20, 335-348 (1956)
 reviewed 1/1957

Edge effects which arise in the near of a contour which nowhere touches the asymptotic lines of the medium surface of a shell, have been treated until now in first approximation only. In the case of axial symmetric shells only Lurje has proposed a method the exactness of which corresponds to that one of the theory of shells. In the general case the complex unknown function

$$W = \sqrt{\frac{h^2}{3(1-\sigma^2)}} 2 E h w + i c$$

(h - half thickness of the shell, σ - coefficient of Poisson, E - Young modulus, w - normal flexure of the shell, c - tension function) is obtained from the differential equation

$$L(W) + \frac{h}{\lambda} \frac{1}{\sqrt{3(1-\sigma^2)}} N(W) = 0 \quad \lambda - \text{characteristic radius of curvature of the shell}$$

by the set up

AUTHOR: J. L. Synge
VOLUME: 1953, Vol. 1, No. 1, pp. 1-12.
TOPICAL: The author has written a paper on the theory of the propagation of waves in the water-beds in the present paper in joint form. The basic equations for the propagation of waves in the usual tensor symbolism without simplifying it in the paper. It is supposed that the chief significance of the paper is in arbitrary coordinate systems, the basic equations given by the author are of service in writing it remains unclear under what circumstances the equations can be used. The author does not say whether the paper is supposed, nor the derivatives is differentiated in the equation and the possible errors are estimated. In addition to applying the fundamental equations, the following note the fundamental equations can be used for a wide range of applications. In these there are many

Card 1/2

The Equations of the Theory of Shells - Variational and Functional of the Type

1) *Equation:*

- A state of stress and strains of shells,
 - A shell's boundary,
 - The outer surface of the shell,
- Equations with the theory of shell's outer boundary.
- and 2), are discussed. The author mentions three possible methods for the approximate calculation of the state of stress in complicated sections of shells without discarding them in detail, according to different preferences.

DATE: April 15, 1977

NAME: Library of Congress

1. Shells-Theory

Card #72

FUKS, Boris Abramovich, prof.; BAKHSHIYAN, P.A., prof.; ANDRIYEVSKIY,
F.P., dotsent; MIROSHKOV, R.K., dotsent; NAGAYEVA, V.M., dotsent;
SOBOLEV, N.A., dotsent; SOKOLOV, A.M., dotsent; SHAPIRO, Z.Ya.,
dotsent; SHUSHARA, G.N., dotsent; KAPLAN, I.B., starshiy pre-
podavatel'; POLOZKOV, A.P., starshiy prepodavatel'; POLOZKOV,
D.P., starshiy prepodavatel'; TOPAZOV, N.G., starshiy prepoda-
vatel'; SHCHERBAKOV, S.S., starshiy prepodavatel'; Prinimali
uchastiye: GOL'DENVEYZER, A.L., prof.; BARANENKOV, G.S., dotsent;
BERMAN, Ya.R., dotsent; LUNTS, G.L., dotsent; SHESTAKOV, A.A.,
dotsent; GMURMAN, V.Ye., starshiy prepodavatel'; Rozental', M.I.,
assistant; SOKOLOVA, L.A., assistant. ROZANOVA, G.K., red.izd-vn;
KUZ'MINA, N.S., tekhn.red. (Continued on next card)

FUKS, Boris Abramovich--(continued) Card 2.

[Higher mathematics; methodological instructions and control assignments for the students of correspondence technical schools of university level] Vysshiaia matematika; metodicheskie ukazaniia i kontrol'nye zadaniia dlia studentov zaочnykh vysshikh tekhnicheskikh uchebnykh zavedenii. Izd.9. Pod red. B.A.Fuksa. Moskva, Gos.izd-vo "Sovetskaiia nauka," 1958. 179 p.

(MIRA 12:9)

1. Russia (1923- U.S.S.R.) Ministerstvo vyshego obrazovaniya.
Metodicheskoye upravleniye.

(Mathematics--Study and teaching)

AUTHOR: Gol'denveyzer, A.L. (Moscow) SCW/24-58-4-19/39

TITLE: On Reissner's Theory of the Bending of Plates (O teorii izgiba plastinok Rayssnera)

PERIODICAL: Izvestiya Akademii Nauk SSSR, Otdeleniye Tekhnicheskikh Nauk, 1958, Nr 4, pp 102 - 109 (USSR)

ABSTRACT: The author discusses Reissner's paper (Ref 1) in which a thin plate of constant thickness is subjected to normal forces of variable intensity at the upper and lower boundaries of the plate. Body forces are assumed to be absent. Reissner's theory is described and its generalisation discussed. The following question is proposed which has greater influence on the corrections introduced by the theory - the elastic phenomena at the boundary of the plate or those far from it? As an example an unclamped circular plate is considered at whose boundary are applied a bending moment, a transverse force and a twisting moment. It is shown that Reissner's theory gives corrections to the constants A_1 and A_2 corresponding to the classical theory and a new constant A_3 is defined. The stressed Cardl/2 state (called by Reissner the boundary effect) associated

307/24-38-4-19/39

On Reissner's Theory of the Bending of Plates

with this constant has a strongly local character. In a special case of the above example, the author finds that Reissner's theory can give wrong corrections to the classical theory. This is because the theory is based on a hypothesis concerning phenomena far from the boundary of the plate, while phenomena near the boundary play an important part. In conclusion, Vlasov's theory (Ref 4) is discussed. It gives the same law for the distribution of the bending stresses. The two theories are compared inconclusively. There are 4 references, 2 of which are Soviet and 2 English.

SUBMITTED: December 2, 1957

Card 2/2

S27/4-Sub-B3/34

AUTHOR: Panovko, Ya. G.
 TITLE: A Conference on Elastic Vibrations at the Institute of
 Mechanical Engineering of the Academy of Sciences of the
 Latvian SSR (Governmental Press of the Latvian Academy of Sciences)

PERIODICAL: Izvestiya Akademii Nauk SSSR, Otdeleniye tekhnicheskikh
 nauk, 1960, No. 10, pp. 156-159 (USSR)

ABSTRACT: This Conference took place on June 11-15, 1959, in Riga.
 Altogether over 100 reports were presented. Eleven papers were read;
 from these normally brief reports eleven papers were published.

1) The effect of vibration on the strength of structural elements by
 I. A. Bel'skii and O. V. Dzhobava. Discussion (Zvezdochkin).

2) Two papers on fatigue of structures by V. V. Buzdin and I. A. S. Val'air
 (Zvezdochkin and Zvezdochkin).

3) A qualitative study of the form and frequencies of
 natural vibrations of thin elastic shells by A. L. Golden-

vezers (Moscow).
 4) Two papers in connection with vibration of elastic
 rods in the case of large displacements by Yu. S. Sineev
 (Leningrad).

Card 1/2
 5) Coupled vibrations of waves and waves in turbulent fluid

"Resonance through resonance of a linear system with non-
 linearly varying frequency" by A. P. Prostakov (Khar'kov).

6) Some problems in the dynamics of an elastically damped
 structural element by V. A. Shchegolev (Khar'kov).

7) On the stability of dynamic processes in solid bodies,
 by A. S. Larin (Tver').

8) The problem of calculating and adjusting a flywheel by G. J.
 Put'yan (Riga).

9) On methods for calculating static stresses in shells absor-
 bing energy by G. I. Vlasov (Riga).

The conference was closed with a speech by Dr. A. P. Sologub-

Bordachev (Riga).

Cart. 172

GOL'DENVEYZER, A.I. (Moskva)

Asymptotic integration of partial differential equations with
parameter dependent boundary conditions. Prikl.mat. i mekh. 22
no.5:657-672 S-0 '58. (MIRA 11:11)
(Differential equations, Partial)

SOV 2660

PHASE I BOOK EXPLOITATION

16(1)

Vsesoyuznyj matematicheskiy s'ezd. 3rd. Moscow. 1956.

Trudy, 3rd. Kratkoye soderzhanie dokladov na vystavochno-tekhnicheskikh uchenykh (Transactions of the 3rd All-Union Mathematical Conference in Mechanics). Vol. 4: Summary of Scientific Reports of Foreign Scientists.) Moscow, 1956. In USSR, 1958. 217 p., 200 copies printed.

Sponsoring Agency: Akademija nauk SSSR. Matematicheskiy institut.
Tech. Ed.: G.N. Shevchenko, Editorial Board: A.I. Abramov, V.G. Boltyanskiy, A.M. Vasil'yev, S.V. Kudrjavcev, A.D. Myshkis, S.M. Nikol'skiy (Resp. Ed.), A.G. Piatnitskiy, V. Prudnikov, J. Ye. Rubnikov, P. L. Ul'yanov, V.A. Uspenskiy, N.O. Shil'nikov, and A.I. Shiranov.

PURPOSE: This book is intended for mathematicians and physicists.

COVERAGE: The book is Volume IV of the Transactions of the Third All-Union Mathematical Conference, held in June and July 1956. The book is divided into two main parts. The first part contains summaries of reports presented at the conference, and the second part contains the texts of reports submitted to the conference that were not included in the first volume. The second part contains the texts of reports submitted to the conference by non-Soviet scientists. It has been decided that this section did not deserve a separate volume, since the title of the paper did not correspond to the approximate volume. The paper, volume, reference to which it refers, and the author's name are given. Both Soviet and non-Soviet scientists presented papers on differential equations, function theory, topology, mathematical analysis, probability theory, foundations of mathematics, and the problems of mechanics and the foundations of mathematics, and the history of mathematics.

TRANSLATION: Yu. L. Kirev, and L.P. Mironik (Editor). The purpose of one new boundary value problem for a difference equation of parabolic type

Section on the Mathematical Problems of Mechanics

Abramov, N.P. (Moscow). On the plane problem of the theory of elasticity for a rectangular region

Vilenkin, V.Z. (Moscow). Method of initial functions in the theory of thick plates under static and dynamic conditions

Zelenchuk, A.D. (Moscow). Formal asymptotic representation of the solutions of partial differential equations with small parameter

Zelenchuk, A.D. (Moscow). Nonlinear vibrations of cylindrical panels in supersonic flow

Zelenchuk, A.D. (Moscow). The method of integral equations in problems of the theory of a thin wire in a compressible flow

Card 20/35

101

102

102

102

102

102

GOL'DENVEYZER, A.L. (Moskva)

Asymptotic integration of linear differential equations with partial derivatives having a small main part. Prikl. mat. i mekh. 23 no.1:
35-37 Ja-F '59. (MIRA L2:2)
(Differential equations, Partial)

Oct. 26 1952 - R. H.

- PROJECT PRESENTED AT THE 1ST ALL UNION CONVENTION OF THE INSTITUTE FOR PROBLEMS IN MECHANICS, BY DR. I. V. DANILOV
66. N. N. Gulyaev, Yu. N. Kuznetsov, Yu. N. Sushchik (Moscow). On problems of calculating strength of electrical power plants under a load of alternating currents of various frequencies and intensities. In view of the fact that the electric load fluctuates, it is of interest to find the corresponding stresses in the structures.
 67. G. A. Gulyaev (Moscow). Some problems concerning the theory of composite materials. Some problems concerning the theory of plastic deformation of plates.
 68. G. A. Gulyaev (Moscow). Strength of plates under a system of elastic loads.
 69. I. G. Gulyaev (Moscow). A dynamic problem for a twisted shell.
 70. N. V. Gulyaev (Moscow). Testimorphosis - a new method of application mechanics to nonlinear problems, simulation of processes of plastic deformation and rupture of solids with great variation of temperature.
 71. N. O. Gulyaev (Moscow). Development of a theory of pressure waves with the use of the method of continuation methods.
 72. I. T. Gulyaev (Moscow). Some generalizations of the basic equations of viscoelasticity.
 73. I. T. Gulyaev (Moscow). The propagation of longitudinal waves in a viscoelastic solid.
 74. A. M. Gulyaev, T. D. Lur'e (Institute of Applied Mathematics and Computational Methods of Design of the USSR Academy of Sciences) Power plants. A generalized theory of plastic flow.
 75. I. T. Gulyaev (Moscow). The theory of finite deformations of anisotropic elastic solids.
 76. I. T. Gulyaev, N. A. Moshkin (Moscow). A general theory of stability of shells.
 77. A. N. Gulyaev (Moscow). Development of the theory of thin-walled cylindrical shells.
 78. A. N. Gulyaev (Moscow). A geometric interpretation of the theory of thin-walled cylindrical shells.
 79. N. I. Gulyaev (Moscow). Generalization of the theory of shells over noncircular bases under the pressure of a liquid. The influence of the pressure of a liquid on the pressure of a liquid.
 80. A. N. Gulyaev (Moscow). On secondary effects in torsion and bending of nearly plastic shells.
 81. I. T. Gulyaev (Moscow). On critical values and the condition of plasticity of anisotropic shells under steady-state conditions of loading.
 82. A. N. Gulyaev, D. S. Zaitsev (Moscow). Continuum in the domain of inelastic mechanics. Continuum of variable length.
 83. A. N. Gulyaev (Moscow). On elastic-plastic deformation of nonhomogeneous plates and shells.
 84. A. N. Gulyaev (Moscow). Equilibrium of nonlinear shells under a system of statical loads and stresses.
 85. G. A. Gulyaev (Moscow). Group action of thin-walled shells.
 86. A. N. Gulyaev (Moscow). The general equations of壳:
 87. G. A. Gulyaev and his particular solution.
 88. D. V. Gulyaev (Moscow). Theory of shells under a load.
 89. A. N. Gulyaev (Moscow). Strong concentration in function spaces.
 90. V. A. Gulyaev, N. I. Gulyaev (Moscow). The presence of an irregularity on an elastic shell.
 91. I. T. Gulyaev (Moscow). Effects of shear stresses in the design of rectangular strips of arbitrary thickness under arbitrary loads.
 92. N. O. Gulyaev (Moscow). The bending of a shell under a load with a reticular base.
 93. N. O. Gulyaev (Moscow). The static equilibrium of an elliptic plate also took into account local loads.
 94. N. O. Gulyaev (Moscow). A plane with a central point reticulated to a hexagonal lattice. A point with a concentrated reaction.
 95. N. O. Gulyaev (Moscow). The effect of a concentrated reaction on the equilibrium of a rectangular plate.
 96. N. O. Gulyaev (Moscow). The effect of a concentrated reaction on the equilibrium of a rectangular plate.
 97. N. O. Gulyaev (Moscow). The effect of a concentrated reaction on the equilibrium of a rectangular plate.

GOLDENVEYSER, A. L. (Acad. Sci. USSR)

"The geometrical criterion of the momentlessness of the state of stress of a thin elastic shell."

Report presented at the 10th International Congress of Applied Mechanics, (ICSU)
Stresa, Italy, 31 August - 7 Sep 1960.

In the author's absence the paper was presented by Oniashvili.
Momentlessness means that nearly everywhere in the shell (except in zones of edge effects), the bending stresses are not significant. Quantitatively, this can be defined by the relative magnitude of the membrane strain energy W_m and the bending strain energy W_b . Let the characteristic of the middle surface K be defined by the equation.

REF ID: A6100

151-100

AUTHOR:

S. S. Devanayyer, A. L.

TITLE:

The asymptotic integration of differential equations with vanishing small main part and oscillating boundary conditions

PUBLICATION:

Referativnyj zhurnal, Matematika, no. 4, 1962, 45,
Bulvarnaya ulica. ("Izdat. Nauk. Sovetov na po differentia-
l'nye uravneniya, 1962". Yerevan, AN ARM SSR, 1960, 73-62)

TEXT:

The author generalizes results which he obtained for the
solution of some problems of the theory of thin elastic shells. The
equation

$$hN(\cdot, \cdot) + L(\cdot) = 0 \quad \dots \quad (1)$$

is considered, where $h > 0$ is a small parameter, L and N linear differential operators with the orders r and n , $r \leq n$, and two independent variables x and y . It is assumed that the coefficients of L and N are sufficiently smooth and that x and y form a coordinate system similar
to the polar system, i. e. the curve $x = x_0$ represents the boundary of
the finite simply connected domain

Card 1/2

37501

5/04/62/62/CCC/004/040/099

C111/0333

2/000/00/000/001/040/029
S101/0535

The asymptotic integration of ...

$$\cdot \quad 0 < x < x_0, \quad 0 < t < 2\pi.$$

Let λ be a large parameter, $\lambda = h^{-t}$, where t is a rational positive number which is smaller in absolute value than $\frac{1}{2}$. The author gives the asymptotic behavior of the solution of equation (1) with quickly oscillating boundary conditions for the Dirichlet and the Cauchy problem (the latter one in the half-disk neighborhood $|z| < x_0$) with conditions of the form

$$\frac{\psi^{(n)}(x)}{x^{\infty}} \rightarrow \psi^{(n)}(\infty) e^{i k \pi} \quad (n=0)$$

where $\psi^{(n)}(\infty)$ is complex, $\psi(\infty)$ is real function and $\psi^{(n)}(\infty) \neq 0$. In dependence on the numbers t and

$\frac{1}{h-1}$ the author considers three cases. The solution is sought in the complex domain. There are many misprints. Another method for solving

Card 2/3

3/044/62/000/004/046/039
0111/0303

The asymptotic interpretation of ...

similar problems is due to M. I. Vishik and L. A. Lyusternik (ZhMat, 1951, 78(6)).

Abstracter's note: Complete translation.

J

Card 5/5

MUSHTARI, Kh.M., red.; ALUMYAE, N.A., red.; BULGAKIN, V.V., red.; VOL'KIN, A.S., red.; GANTYEV, N.S., red.; GLUDENOVICH, A.L., red.; ISKAMBAYEVA, F.S., red.; KIL'CHEVSKIY, N.A., red.; KORNISHIN, M.S., red.; LUR'YE, A.I., red.; SAVIN, G.N., red.; SACHENKOV, A.V., red.; SVIRSKIY, I.V., red.; SURKIN, R.G., red.; FILIPPOV, A.I., red.; ALEKSEGIN, V.I., red.; SEMENOV, Yu.P., tekhn. red.

[Proceedings of the Conference on the Theory of Plates and Shells] Trudy Konferentsii po teorii plastin i obolochek, Kazan', 1960. Kazan', Akad. nauk SSSR, Kazanskii filial, 1960. (426 p.) (MIFI 15:7)

1. Konferentsiya po teorii plastin i obolochek, Kazan', 1960.
2. Moskovskiy energeticheskiy institut (for Fel'dtin). 3. Kazanskiy khimiko-tehnologicheskiy institut (for Janiyev).
4. Institut mekhaniki Akademii nauk USSR (for Kil'chevskiy).
5. Kazanskiy gosudarstvennyy universitet (for Sachenkov).
6. Kazanskiy filial Akademii nauk SSSR (for Svirskiy).
(Elastic plates and shells)

83225

S/042/60/015/005/001/005
C111/C222

10.7300

AUTHOR: Gol'denveizer, A L.

TITLE: Some Mathematical Problems in the Linear Theory of Elastic
Thin Shells

PERIODICAL: Uspekhi matematicheskikh nauk 1960 Vol.16 No.5 pp.3-75.

TEXT: The author has the aim to turn the attention of the mathematicians to the difficulties of the theory of shells and gives a representation of the corresponding mathematical problems. The contents of the paper is partially taken from the author's book (Ref.:) and partially from his numerous publications (Ref.6.9.11.12.14.23.26).

Contents: Introduction; chapter I: Asymptotic methods for the integration of partial differential equations; chapter II: Binding by boundary conditions; chapter III: Eigenvalue problems of the theory of shells; chapter IV: Theory of shells free of moments and its connection with the theory of infinitely small deformations; chapter V: Asymptotic integration of the differential equations of the theory of shells subject to moments; chapter VI: Influence of the conditions of clamping to

Card 1/2

8325
S/042/60/015/005/001/005
C111/C222

Some Mathematical Problems in the Linear Theory of Elastic Thin Shells;

the state of stress of the shell;

The author mentions I.N. Vekua. There are 26 references, 23 Soviet
2 American and 1 English.

SUBMITTED: November 5, 1959

Card 2/2

PHASE I BOOK EXPLOITATION SOV/6201

Vsesoyuznyy s"yezd po teoreticheskoy i prikladnoy mekhanike. 1st, Moscow, 1960.

Trudy Vsesoyuznogo s"yezda po teoreticheskoy i prikladnoy mekhanike,
27 yanvarya -- 3 fevralya 1960 g. Obzornyye doklady (Transactions of the
All-Union Congress on Theoretical and Applied Mechanics, 27 January to
3 February 1960. Summary Reports). Moscow, Izd-vo AN SSSR, 1962.
467 p. 3000 copies printed.

Sponsoring Agency: Akademiya nauk SSSR. Natsional'nyy komitet SSSR po
teoreticheskoy i prikladnoy mekhanike.

Editorial Board: L. I. Sedov, Chairman; V. V. Sokolovskiy, Deputy Chairman;
G. S. Shapiro, Scientific Secretary; G. Yu. Dzhanelidze, S. V. Kalinin,
L. G. Loytsyanskiy, A. I. Lurye, G. K. Mikhaylov, G. I. Petrov, and
V. V. Rumyantsev; Resp. Ed.: L. I. Sedov; Ed. of Publishing House:
A. G. Chakhirev; Tech. Ed.: R. A. Zamarayeva.

Card 1/ 6

Transactions of the All-Union Congress (Cont.)

SOV/6201

PURPOSE: This book is intended for scientific and engineering personnel who are interested in recent work in theoretical and applied mechanics.

COVERAGE: The articles included in these transactions are arranged by general subject matter under the following heads: general and applied mechanics (5 papers), fluid mechanics (10 papers), and the mechanics of rigid bodies (8 papers). Besides the organizational personnel of the congress, no personalities are mentioned. Six of the papers in the present collection have no references; the remaining 17 contain approximately 1400 references in Russian, Ukrainian, English, German, Czechoslovak, Rumanian, French, Italian, and Dutch.

TABLE OF CONTENTS:

SECTION I. GENERAL AND APPLIED MECHANICS

Artobolevskiy, I. I. Basic Problems of Modern Machine Dynamics	5
Bogolyubov, N. N., and Yu. A. Mitropol'skiy. Analytic Methods of the Theory of Nonlinear Oscillations	25

Card 2/6

Transactions of the All-Union Congress (Cont.) SOV/6201

Sretenskiy, L. N. Review of Reports on the Theory of Tides 213

Struminskii, V. V. Present State of the Problem of Supersonic
Gas Flow Past Bodies 225

SECTION III. MECHANICS OF RIGID BODIES

Berezantsev, V. G. The Theory of Limiting State of Stress in
Soil Mechanics and Its Applications 299

Vekua, I. N., and N. I. Muskhelishvili. Methods of the Theory
of Analytic Functions in the Theory of Elasticity 310

Gol'denveizer, A. L. Development of the Theory of Elastic
Thin Shells 339

Card 5/6

SAVIN, G.N., otv.red.; ABADUOV, A.A., red.; ALUMAE, N.A., red.;
AMBARTSUMIAN, S.A., red.; ANILO, I.Ye., red.; BULATIN, V.V., red.;
VOL'MIN, A.S., red.; GALENENKO, A.L., red.; GR.GOLYUK, E.I.,
red.; KAN, S.F., red.; KAMISHIN, A.V., red.; KIL'CHEVSKIY, N.A.,
red.; KISELEV, V.A., red.; KOVALENKO, A.D., red.; MUSHTARI, Kh.N.,
red.; MOVOZHILOV, V.V., red.; OMANSKIY, A.A., red.; FILIPOV, A.P.,
red.; LINGVETS, A.M., tekhn. red.

[Proceedings of the Second All-Union Conference on the Theory of
Plates and Shells] Trudy Vsesoyuznoi konferentsii po teorii plastin i
oklochek. Lvov, Pol.Kiev, Izd-vo Akad. Nauk U.S.S.R., 1961. (MRA 10:12)

I. Vsesoyuznaya konferentsiya po teorii plastin i oklochek. 2,
Lvov, 1961.
(Elastic plates and shells)

x. 13
3/046/01/025/004/012/021
D274/D306

244200

1321 also 2607 2807

AUTHOR: Gol'denveyzor, A.L. (Gloscov)

TITLE: Asymptotic properties of eigenvalues in the elastic-shell theory

PERIODICAL: Trudy Akademii Nauk Gruzinii, v. 21, no. 4,
1961, p. 19-74

TEXT: Linear problems are considered of free oscillations and the stability of thin elastic shells. Special attention being given to asymptotic properties of eigenvalues as a function of the density and configuration of the nodal lines of the eigenfunctions. It was shown by the author (R.F. 11-A. L. Gol'denveyzor, Teoriya uprugikh toroidov obolochek (Theory of thin elastic shells), GITRL, 1955) that in many cases the approximate description of the stress-strain state of elastic shells reduces to the following of equations:

$$\begin{aligned} L(C) - a^2 R^2 \cdot (\partial^2 u / \partial r^2) + K \cdot u &= 0 \\ L(2RM) - M(C) &= 0 \end{aligned} \quad a^2 = \frac{12}{3K + 4E^2} \quad (1.1)$$

where C is the shear modulus, M is the normal modulus, $K = a$
Carl 1/7

S/040/01/023/004/012/021
D276/D366

Asymptotic properties.

$$\frac{m}{2Eh} \omega^2 = k^{2\alpha-4} \frac{1}{v} \left(t + h^{1-\alpha} \frac{4\pi^2}{3G} \frac{R^2}{(1-\frac{\alpha}{2})^2} n \right) \quad (4.1)$$

k is a parameter; $\frac{1}{v} = \left(\frac{h\sqrt{R}}{E} \right)^{\alpha}$, $t > 0$.

t , n and v are random variables;

$$\begin{aligned} k^2 \rho_1 &= \int_{\mathbb{R}} \{ H(v) \text{ and } \psi(v) \} \{ 1 - \psi(v) \} v^{\alpha-1} dv \\ \kappa_{11}^0 &= \int_{\mathbb{R}} \{ H(v) \text{ and } \psi(v) \} \{ 1 - \psi(v) \} v^{\alpha-1} v^2 dv \\ k^4 \varphi &= \int_{\mathbb{R}} \{ H(v) \text{ and } \psi(v) \} \{ 1 - \psi(v) \} v^{\alpha-1} v^4 dv \end{aligned} \quad (4.2)$$

(φ has yet to be determined); for q_0 one obtains:

$$q_0 = k^{2\alpha-2} \frac{1}{v} \left(t + h^{1-\alpha} \frac{4\pi^2}{3G} \frac{R^2}{(1-\frac{\alpha}{2})^2} n \right) \quad (4.3)$$

where $\kappa_{11}^4 \varphi = \int_{\mathbb{R}} \{ H(v) \text{ and } \psi(v) \} \{ 1 - \psi(v) \} v^{\alpha-1} v^4 dv$ (4.4)

(φ has yet to be determined), v and n in Eq. (4.3) are given by

Card 5/7

4f.33
S/040/01/025/004/012/021
D274/0506

Asymptotic properties...

$$\begin{aligned} c &= c_* \left[r_1 \cos k(f_1 + f_2) + r_2 \sin k(f_1 + f_2) \right] \\ w &= w_* \left[\cos k(f_1 + f_2) + \sin k(f_1 + f_2) \right] \end{aligned} \quad (3.1)$$

where c_* , w_* , f_1 , f_2 , r_1 , r_2 are functions of θ and ρ which can be chosen. It is required that w_* and c_* be non-negative. The stress-strain state D is considered, determined by (3.1) and (1.1). The density of the nodal lines of D increases with k , i.e., with (for given h/R); the number N is termed the index variability. By appropriately choosing f_1 and f_2 it is possible that state D should have two (or one) system of nodal lines which belong to two (or one) pre-assigned families of curves, by the appropriate choice of V it is possible with fixed f_1 and f_2 to increase or to reduce the density of nodal lines. It is postulated that the functions f_1 , f_2 , w_* , c_* , r_1 , r_2 can be chosen in such a way that their variability should not be very large and that c and w should be sufficiently close to a solution of Eq. (1.1) - (3.1); with such a choice of these functions, formulas (3.1) and (4.6) give sufficiently exact values of ω^2 and ϵ_0 . This postulate is verified for all cases, except when the middle

Card 4/7

S/040/01/0017/004/012/021
D/77, D/006

Asymptotic properties

surface of the shell has negative curvature. Now the problem consists in constructing asymptotic (for $k \rightarrow \infty$) expressions for the integrals (4,3), (4,5) on the assumption that α and β have the form (5,1). Thus, the condition of compatibility is satisfied.

$$\iint_D \{L_0(r_1) + L_0^*(r_1)\} \cdot e_1 u_{111} dr_1 dr_2 \neq 0 \quad (10.3)$$

where L_0 is given by

$$L_0 = \frac{1}{\lambda^2 R_1^2} \frac{\partial}{\partial r_1} \left(\frac{\partial u_1}{\partial r_1} \right)^2 + \frac{1}{\lambda^2 R_1^2} \frac{\partial}{\partial r_1} \left(\frac{\partial u_1}{\partial r_2} \right)^2$$

this condition may be imposed at all points of the region under investigation

$$L_0'' + L_0^{**} = 0 \quad (10.4)$$

It follows that in (4,3) the quantities u_1 , v_1 and v_2 will remain finite when $k \rightarrow \infty$, if ρ is approached by choosing $\beta \neq 2$ if condition (10.3) is satisfied, and $\beta = 0$ if (10.4) holds. If both relations do not hold one should possibly assume that $\beta = 1$. For relatively flat shells one should probably assume that $\beta = 1$. For a shell of positive curvature, (10.3) always holds. For zero curvature, (10.3) is not valid, but a simple case when zero strain values of state D can only be assumed on the lines (the asymptotic lines of

Card 5/7

S/060/01/025/004/012/021
D.267/10366

Asymptotic properties

the middle prisms, and have extrema which are considered. Further, the asymptotic properties of the eigenfunctions are discussed. On the basis of the theory of stability it is shown that β is positive, hence λ is large, and the eigenvalues will take the characterisitic value λ_0 as β tends to zero. (12.2)

λ_0 depends on α , β , γ and δ only on one of the three values: 2, 1, 0; (hence λ_0 is a bimodality value too). A typical (for shell theory) reversal in respect to eigenvalue increase with increasing number of nodal lines in this takes place only up to a certain point - when it occurs, the characteristic value λ_0 ; thereupon the regular stability is established; increasing eigenvalues with an increasing number of nodal lines. In stability problems, the least value of the critical load is important. It is found, for which configuration of nodal lines, loss of stability occurs. For zero curvature, 2 cases may arise: in the first case, stability may be lost for one family of nodal lines which coincide with rectilinear generators; in the second case stability is not

Card 6/7

S/040/62/026/004/004/012
D400/2701

24.4200

AUTHOR: Col'denveyzer, A.L. (Moscow)

TITLE: Construction of an approximate theory of bending of plates by the method of asymptotic integration of the equations of elasticity theory

PERIODICAL: Prikladnaya matematika i mehanika, v. 26, no. 4,
1962, 668 - 686

1/3

TEXT: The possibility of rendering more exact the classical theory of bending of plates is considered. The bending problem is formulated as a three-dimensional problem of elasticity theory which is solved by the iteration method; thereby it is assumed that one of the dimensions is small as compared to the other two. The stressed state of the plate is sought in the form of a sum of a slowly-decreasing (with distance from the plate edge) stressed state which is constructed by means of the principal iteration process, and of fast-decreasing stressed states, constructed by means of auxiliary iteration processes. Such an approach is often used in the asymptotic integration of differential equations and corresponds to the

Card 1/4

S/040/62/026/004/004/013
D409/D501

Construction of an approximate ...

tions. The auxiliary iteration process is constructed in two different ways. In the first, the construction of the solution amounts to the integration of an harmonic equation, whereas in the second, the solution involves the integration of a biharmonic equation. Five types of boundary conditions are considered, and the corresponding equations are set up. These equations are used to determine the sought-for functions (the biharmonic function $\Psi(x, y)$, the harmonic function $\Phi(x, y)$, and the biharmonic function $\Psi_F(x, y)$). The function $\Psi(x, y)$ can be expressed in terms of $\Phi(x, y)$. The main consequence of the above results is as follows: the stressed state has 3 components (the principal stressed state, the stressed state of edge twisting, and the stressed state of plane edge deformation). The principal stressed state corresponds to the principal iteration process, whereas the other stressed states correspond to the auxiliary processes. With such an approach, classical theory can be considered as an approximate method, based on the principal iteration process only, for which only the first approximation is constructed. The fundamental difference between the proposed method and classical theory, consists in introducing the auxiliary iteration processes, i.e. the processes constructed by integration of diffe-

Card 5/4

S/040/62/C26/C04/C04/C13
D409/D501

Construction of an approximate ...

differential equations which contain ξ as an independent variable.

SUBMITTED: April 5, 1962

/E

Card 4/4

L 12946-63 EWP(r)/EWT(m)/BDS AFFTC
ACCESSION NR: AP3004108

S/0040/63/021/004/0593/0608

AUTHOR: Col'denveyzer, A. L. (Moscow)

5D

TITLE: Development of an approximate shell theory by the asymptotic integration
of the elasticity-theory equations

SOURCE: Prikladnaya matematika i mekhanika, v. 27, no. 4, 1963, 593-608

TOPIC TAGS: approximate shell theory, asymptotic integration, shell theory

ABSTRACT: An asymptotic method of integration of differential equations of the elasticity theory is proposed, by means of which an approximate theory of shells can be established with a desired degree of accuracy in a way analogous to that used earlier by the author to develop an approximate theory of flexure of plates (Postroyeniye priblizhennoy teorii izgiba plastinki metodom asimptoticheskogo integrirovaniya uravneniy teorii uprugosti, FMM, 1962, v. 26, no. 4). This is closely associated with the method of asymptotic integration of differential equations of the theory of shells discussed in the author's monograph Teoriya uprugikh tonkikh obolochek, Gostekhizdat, 1953. Tensor analysis is applied in the representation and solution of the initial system of differential equations

Card 1/2

L 12946-63
ACCESSION NR: AP3004108

(equilibrium equations, symmetry conditions, and elasticity relationships) for determining displacements and stresses. Iterative processes are formulated for determining the states of stress which are, in the zero approximation, equivalent to the membrane-stress state, the pure flexural-stress state, and the states with large indexes of variation, as well as the iterative processes corresponding to the states of torsion and of plane strain at the edges. Through the combination of these iterative processes, the boundary conditions of the three-dimensional elasticity theory can be satisfied with an arbitrary degree of accuracy. The physical interpretation of the equations of the iterative processes is given. Certain conditions ensuring the asymptotic convergence of these iterative processes and thus determining the region of application of results obtained are briefly discussed. Orig. art. has: 62 formulas.

ASSOCIATION: none

SUBMITTED: 15Jan63

DATE ACQ: 15Aug63

ENCL: 00

SUB CODE: AP

NO REF Sov: 007

OTHER: 008

Card 2/2

GOL'DENVEYZLR, A.L. (Moscow):

"Asymptotic methods of analysis of the spectrum of free vibration frequencies of shells".

report presented at the 2nd All-Union Congress on Theoretical and Applied Mechanics, Moscow, 29 Jan - 5 Feb 64.

GOL'DENVEYZER, A. L.

"The principles of reducing three-dimensional problems of elasticity to two-dimensional problems of elasticity to two-dimensional problems of the theory of plates and shells."

report submitted for 11th Intl Cong of Theoretical & Applied Mechanics & General Assembly, Munich, 30 Aug-1 Sep 64.

L 41657-65 EWT(d)/EWT(m)/EWA(d)/EMP(w) EM

S/0040/58/029/001/0141/0155

ACCESSION NR: AP5006263

27

AUTHOR: Gol'denveyzer, A. L. (Moscow); Kolas, A. V. (Moscow)

B

TITLE: Contribution to the construction of the two-dimensional equations of the theory of thin elastic plates

SOURCE: Prikladnaya matematika i mehanika, v. 29, no. 1, 1965, 141-155

TOPIC TAGS: elasticity theory, elastic shell, applied mathematics, mechanical stress, strain measurement, stress calculation

ABSTRACT: The authors discuss ways to construct an approximate theory governing the calculation of thin elastic plates without employing assumptions typified by Kirchhoff's hypothesis. Up to now the only method of solving this problem was the method based on the use of power series or series expansions in Legendre polynomials. In some recent papers such problems have been handled by asymptotic integration of the equations of elasticity theory. In the present work the authors discuss the features of these methods and derive equations to which the asymptotic method reduces in the problem of the general strain of thin plates whose mean surface is related to an arbitrary orthogonal system of curvilinear coordinates. Orig. art. has: 59 formulas.

Card 1/2

L 64122-55 EWT(d)/EWT(m)/EWP(w)/EWA(d)/EWP(k)/EWA(h)/EWP(r) WW/DC

UR/0040/65/025/004/0701/0715

ACCESSION NR: AP5021303

AUTHOR: Gol'denveyzer, A. L. (Moscow)

TITLE: On errors in the classical linear shell theory and on means of improving it

SOURCE: Prikladnaya matematika i mehanika, v. 29, no. 4, 1965, 701-715

TOPIC TAGS: linear shell theory, improved classical shell theory, classical shell theory, asymptotic method, three dimensional elasticity equation

ABSTRACT: An asymptotic method of integrating the three-dimensional equations of the theory of elasticity is proposed for determining the stresses and displacements in closed shells in which the effect of support conditions is eliminated (for example, in a complete sphere). It is assumed that the curvatures of the middle surface of the shell change smoothly, that its reduced length is not too large, and that the stress distribution sought for can be formally constructed by means of the membrane theory under an arbitrary set equilibrating

Card 1/3

L 64122-65

ACCESSION NR: AP5021303

system of stresses with components differentiable a sufficient number of times. The results obtained by this method are compared with data obtained by applying the classical (based on the Kirchhoff-Love hypotheses) theory of shells, and the effect of errors caused by assumptions made in its initial relationships on the Fifi's results is investigated. In order to compare both results, the final formulas obtained by the method proposed are expressed in terms of the classical shell theory. The error estimates given here take account of the index of variation t , and it is shown that these errors (which have in the classical theory an order of the nondimensional thickness h_*) can be essentially reduced (up to values of the order h_*^{2-2t}). The expressions for the elasticity relationships which must be used to achieve this improvement are derived. The comparison leads to the conclusion that a more exact classical shell theory can be proposed for the solution of the discussed problem in which the error (in the case when $t = 0$) will be of the order t^2 in comparison with unity. The effect of the variations in the state of stress on the values of errors in the classical theory is also discussed. Orig. [VK]

ASSOCIATION: none

Card 2/2

APPROVED FOR RELEASE: Thursday, September 26, 2002

CIA-RDP86-00513R000515620015-6

APPROVED FOR RELEASE: Thursday, September 26, 2002

CIA-RDP86-00513R000515620015-6

L 64122-65

ACCESSION NR: AP5021303

SUBMITTED: 22 Apr 65

UNCLAS: ENCLB: 00

SUB CODE: AS

NO REF Sov: 006

OTHER: 002

ATTN PRESS: 4030

Card 3/3

L 20626-66 EXP{z}/EXP{w}/EXP{w}/EXP{y}/EXP{y}/EXP{E}/EXP{r} Card 1/2
ACC NR: AF6000/141

AUTHOR: Gol'denveyzer, A. L. (Moscow)

9222

ORG: none

TITLE: Qualitative analysis of free vibration of elastic thin shells

SOURCE: Prikladnaya matematika i mehanika, v. 30, no. 1, 1966, 94-108

TOPIC TAGS: shell, thin shell, shell vibration, shell natural frequency, vibration mode

ABSTRACT: An asymptotic method of integrating dynamic equations associated with free-vibration problems of the classic linear theory of elastic thin shells is presented. Equations of equilibrium, elasticity, and strain-displacement relationships, containing the frequency and displacement parameters, are taken from the author's "Theory of elastic thin shells" and are used as initial equations in investigating the free vibration of an elastic thin shell, by a method which is a "dynamic" version of the asymptotic method developed by the author in the above-mentioned book for solving the static problem. The principal attention is paid to vibrations associated with a large index of variation in the states of stress and strain. The problem is solved in a rough approximation; the possibility of refinements is discussed. The asymptotic properties of expressions for determining the frequencies and the associated states of stress are analyzed in relation to the order of the magnitude of the nondimensional thickness of the shell, and to the density

Card 1/2

L 20606-66

ACC NR: AP6007581

and shape of nodal lines. The classification of free-vibration modes is established, simplified equations for determining them in the first approximation are derived, and qualitative analyses of their natural-frequency spectra are carried out. The characteristic features of the boundary conditions in problems not studied before are discussed only qualitatively. New concepts of "quasi-lateral" and "quasi-tangential" vibrations (characterized by the predominance of the lateral and tangential displacements, respectively) are introduced, as well as of the concepts their integrals, which are analogous to integrals with a large index of variation in the static problem where they describe the distributions of flexural and tangential stresses. Examples of examining the existence of certain modes of vibration, and the spectra of natural frequencies are given. Orig. art. has: 1 table and 38 formulas.

[VK]

SUB CODE: 20/ SUBM DATE: 23Sep65/ ORIG REF: 006/ OTH REF: 002/ ATD PRESS:

4225

Card 2/2

"C. HENKYSER, . I.

"Investigation of the action of sulfur dioxide on various metal solutions."
Sht. 2 Dec. 51, "Central Center of Lenin Radiotekhnicheskii Inst. (part 1)." M. Mandalevsky

Dissertation accepted for defense and with minor changes
Moscow inst. 1951.

Kh. - um. - o. 1951. 0. 1. 1951.

GOL'DENTSYAK, Ya.D.

Determination of carbon dioxide pressure in the blood in
clinical practice. Lab.delo 5 no.2:17-24 Mr.Ap '59.
(MIRA 12:5)

1. Iz kafedry propedevtiki vnutrennikh bolezney (zav. - dots.
Z.A.Gorbunkova) Smolenskogo meditsinskogo instituta.
(BLOOD--ANALYSIS AND CHEMISTRY) (CARBON DIOXIDE)

ARKHANGEL'SKIY, Ye.V., kand.tekhn.nauk; GOL'DENTUL, B.A., inzh.

Improvement in methods of determining load on switching throat-tracks.
Vest.TSMII MPS 18 no.1:61-63 F '59. (MIRA 12:3)
(Poland--Railroads--Switching)

the following information is contained in the document:

Information concerning the development of the Soviet space program, particularly the R-7 rocket and the R-7 missile.

Information concerning the development of the Soviet space program, particularly the R-7 rocket and the R-7 missile.

KROPACHEV, N.G., inzh.; GOL'DER, E.L., inzh.

Operational accounting and analysis of production cost in
steel foundries and rolling mills of the Kuznetsk Metallurgical
Combine. Stal' 25 no.10:953-955 O '65. (MIRA 18:11)

1. Kuznetskiy metallurgicheskiy kombinat.

ACCESSION NR: AP4037174

6/20/99/06/22/2001/290/0285

AUTHOR: Vaynshtok, V. V.; Kertishev, D. A.; Gol'der, G. A.

TITLE: The structure of soaps modified by addition of lead and lithium stearate.

SOURCE: *Kolloidnyy zhurnal*, v. 24, n. 1, p. 290-297, 1962, ausz. Russ. Soed. Akad. Nauk.

TOPIC CODES: soap oil dispersion structures, soap electrosorption, lithium stearate, lithium stearate, aluminum stearate, eutectic mixture, lithium stearate crystal, crystal aggregate, lithium lithium stearate aggregate, lithium dispersed particle

ABSTRACT: The authors studied the crystallization of lithium stearate added with other stearates, widely used in the manufacture of lubricating oil soap (soaps), dispersions), and conducted electro-sorption and soap electrosorption on modified soaps. Their melts and the soap-oil films that prepared by the method of the film method specimens were prepared in anhydrous ether and powdered on filter paper in diethyl ether. The results are photographs, considered as dispersed. Thus, the powdered soaps differed little in outer aspect, some of them had small crystalline structures depending upon the soap earlier, and will not be continued.

CONF 4/3

ACCESSION NR: AP4037174

thickener composition. The lead-stearate based greases showed a marked increase in thickening ability with a low degree of crosslinking, whereas the thickening ability and thickening ability of such systems aluminum and lithium stearates formed distinctly shaped combined curves (to 25% mol.-% of aluminum stearate). Increase of aluminum soap concentration caused a decrease of colloidal stability and rheologic indicators. No significant difference was observed in X-ray studies of these stearates, due to the relatively good crystalline structure diffraction, with the exception of the tri-n-butyl citrate. Diffraction line distances and line intensity are relatively weak, the size of the particles being less pronounced than those of the stearates. The softening effect of the lead-stearate on the structure seems to be more important, and the influence on formation of joint dispersed particles is at 15.00% mol.-% aluminum stearate formation of the structure at 30.00%. Other elements do not influence significantly lead or aluminum stearate ability to form joint dispersed particles. Dispersed particles did not form in presence of citrate, because the citrate was combined in the tri-n-butyl citrate. The mechanism of crystallization at room temperature could probably be explained by neutral lead stearate combined with citrate, which are concentrated in the core due

Page 2/3

ASSOC CODE: AP4037174

20-30 mol% concentration. Nickel, lead and aluminum steenraces...
separately from melts as eutectic mixtures. Cris. temp. 1300°C.

ASSOCIATION: Moskovskiy Instituts metallokhimicheskoy i gazovoy promstv. im. I. M. Gubkina (Moscow Institute of Petrochemical and Gas Industry)

SUBMITTED: 02Nov62

EWU: 100

SUB CODE: FP

NO PTF CODE: 001

STENOP: 0.7

Conf: 3/3

APPROVED FOR RELEASE: Thursday, September 26, 2002 : CITA-RDSSC04516R00516020910-6

APPROVED FOR RELEASE: Thursday, September 26, 2002 : CITA-RDSSC04516R00516020910-6

Some Results of the Application of Spectrographic Analysis to Non-Ferrous Light Alloys. G. Goldber (*Izv. Akad. Nauk S.S.R.*, 1940, *Fiz. Khim.*, 21S, 219). [In Russian.] A short account of the application of the technique to the smelting of Duralumin (17S and 24S), Al-Mg, and Al-Si alloys. N. A.

ASA-SLA METALLURGICAL LITERATURE CLASSIFICATION

12

APPROVED FOR RELEASE: Thursday, September 26, 2002

CIA-RDP86-00513R000515620015-6

APPROVED FOR RELEASE: Thursday, September 26, 2002

CIA-RDP86-00513R000515620015-6"

GOL'DER, G. A.

Grad. Technical Sci.

"Energy and Stability of Crystal Lattices." Sub 6 Mar 47, Moscow
Aviation Technological Inst

Dissertations presented for degrees in science and engineering in Moscow
in 1947

SO: Sum No. 457, 1^o Apr 55

GOL'DER, G.A.; UMANSKIY, M.M.

Goniometric and X-ray analysis of crystals of 1,3,8-trinitronaphthalene. Zhur. Fiz. Khim. 25, 555-6 '51.
(MLRA 4:5)
(CA 47 no.17:8457 '53)

I. L.Ya.Karpov Phys.-Chem. Inst., Moscow.

X-ray study of crystals of some nitro and halogen derivatives of benzene and naphthalene. G. A. Gol'der, G. S. Zhdanov, M. M. Umanskii, and V. P. Glushkova (L. Ya. Karpov Phys.-Chem. Inst., Moscow). Zhur. Fiz. Khim. 26, 1250-65 (1952).—The 1,8-dichloronaphthalene crystallizes from hexane in the form of elongated transparent plates, $m. 87^\circ$, $d = 1.51$. Each plate has a 110° angle between the edges of rhombic prisms $c[001]$ and $m[100]$. The unit cell has $a = 11.5$, $b = 10.8$, $c = 7.9\text{kX}$, $d(\text{x-ray}) = 1.53$; the space group $C_{2h}^6 = P2_1/C$, 4 mols. per cell. It was detd. that $\{h0l\}$ is present only when $l = 2n$, and $\{0k0\}$ when $k = 2n$. Colorless crystals of 2,6-dichloro-1-nitrobenzene (from cyclohexane) have $a[100]$, $b[010]$, $c[001]$, $\bar{a}[101]$. It crystallizes with 4 mols. in a monoclinic cell with $a = 5.82$, $b = 9.33$, $c = 14.2\text{kX}$, $\beta = 91^\circ$, $d = 1.46$, $d(\text{x-ray}) = 1.61$, its space group $C_{2h}^6 = P2_1/m$ or $C_2^6 = P2_1$. Monoclinic crystals of 2,4,6-tribromo-1-nitrobenzene crystallize from chloroform. The unit cell has $a = 9.3$, $b = 12.4$, $c = 9.8\text{kX}$, $\beta = 127^\circ 20'$, $d = 2.40$, $d(\text{x-ray}) = 2.54$, and contains 4 formula units. It was estd. that $\{hkl\}$ is present only when $k + l = 2n$, $\{h0l\}$ when $k = 2n$ and $l = 2n$, and the $\{0k0\}$ is present only when $k = 2n$. The crystal has space group $C_{2h}^6 = A2/a$ or $C_2^6 = Aa$. The benzophenone crystals from hexane have well-defined facets of rhombic prisms: $a[100]$, $b[010]$, $c[001]$, $m[110]$, $d[101]$, $\bar{a}[201]$, and rhombic dipyramids $\{111\}$. Its unit cell has $a = 8.0$, $b = 10.2$, $c = 12\text{kX}$, d . (by flotation method) = 1.1, $d(\text{x-ray}) = 1.05$; 4 mols. per cell with space group $D_4^6 = P2_12_12_1$. The $\{h00\}$ is present when $h = 2n$; $\{0k0\}$ when $k = 2n$; $\{00l\}$ only when $l = 2n$. Rhombic crystals of 1,3,5-trinitrobenzene have the following dimensions of a

unit cell: $a = 12.8$, $b = 27.0$, $c = 9.8\text{A}_1$, with 10 formula units in each. The space group $D_{3h}^6 = P_{na2}$. The golden-colored needles of 1,3,6,8-tetrinitronaphthalene (I) (from EtOH) gave complicated x-ray diffraction probably owing to "regular polysynthetic ferrantion." X-ray study of these crystals at -110° eliminated the possibility of interferences due to thermal vibrations. Crystals obtained from other solvents (e.g. AmOAc , ligroin, AcOH) gave similar interferences in x-ray diagrams. Crystn. from the mixts. of acetone with benzene or with toluene led to formation of new compds., which were very unstable in the air. By choosing planes without diffuse spots these investigators were able to show that the unit cell of I has $a = 26.3$, $b = 7.75$, $c = 5.54\text{kX}$, and when $d = 1.64$ there are 4 mols. in a cell. For such a cell the $\{h0l\}$ was estd. to be present only at $h = 2n$, $\{0kl\}$ when $k + l = 2n$. On these bases the space group can be assigned: $D_{3h}^6 = Pna2$ or $C_{2h}^6 = Pna2$. The x-ray study of 2,4,6-trinitrotoluene (II), with interferences analogous to I, is in disagreement with E. Hertel's expts. (C.A. 27, 5228). By choosing only well-defined diffraction patterns it was possible to det. that the unit cell of II has 4 mols. with $a = 20.2$, $b = 6.2$, $c = 7.7\text{kX}$, and the space group $C_{2h}^6 = P_{n1}/m$ or $C_2^6 = P_{n1}$. It is concluded that in II, as in I, no true monoclinic crystals are formed.

Anatole P. Kotloby

10
15/54

GOL'DER, G.A.

The crystal structure of trichlorobenzonitrile. A. Gol'der, G. S. Zhuravlov, and M. M. Umnanskii (USSR). Zhur. Fiz. Khim. Rast. 24, 2427. (1950). — The mol. structure of 2,4,6-trichlorobenzonitrile was detd. by Fourier analysis. It is monoclinic; the space group is C_{2h}^1 . The unit cell has the dimensions $a = 4.10$, $b = 10.87$, $c = 11.81$ kX, and $\beta = 91^\circ 30'$; and contains 4 mols. The plane in which the mols. lie forms a 21° angle with the (100) plane of the crystal. The line through C₁ and C₄ in the ring forms a $54^\circ 30'$ angle with the (010) direction. The bond distances C-C (nitrile), C₁-C₂, C₂-C₃, C₃-C₄, C₄-Cl, C₆-Cl, and N-C (nitrile) are 1.47, 1.39, 1.41, 1.38, 1.76, 1.77, and 1.18 kX, resp. The C₁=C₂—Cl angle is distorted about 2-3°.

J. W. Lovberg, Jr.

GOL'DER, G. A.

USSR/Physics - Dislocations

Feb 52

"Translation of A. H. Cottrell's 'Theory of Dislocations in Crystalline Lattice,'" by G. A. Gol'-der

"Uspekhi Fiz Nauk" Vol XLVI, No 2, pp 179-230

Russian translation of English-language article, which appeared in "Progress in Metal Phys," edited by B. Chalmer, 1949, p 77. Translation made under editorship of Prof G. S. Zhdanov. Editor discusses differences in the following tech terms that are otherwise synonyms: "Zatsepleniye" (meshing), "dislokatsiya" (dislocation), "smeshcheniye" (shift), "stsepleniye" (gripping, cohesion). 2107100

Röntgenographic determination of structure of σ -nitro- p -chloroaniline. G. A. Gol'dur, G. S. Shustov, and N. M. Umnovskii. Zh. struk. soed. 1982, No. 2, p. 123-126.

(Abstract, cf. Kitagurodskii, T. I., 1981, 10, No. 2, p. 92.)

The bulk cell of picryl chloride (I) is $a = 11.10$, $b = 0.83$, $c = 12.20$ Å, $\beta = 102^{\circ}30'$, space group C_{2h}^1 , $P2_1/c$. Each unit cell contains 4 mols. The C—Cl bond length is 1.71 Å, C—C bond lengths in the ring are normal (1.37-1.41 Å), C—N bond lengths are 1.31 Å, from sum of covalent radii, but in the σ -NO₂ groups the C—N bond length is 1.36 Å, whereas in p -position it was 1.38 Å. The σ -groups are thus nearly perpendicular to the ring, but the p -group is paraEcl.

G. M. Kosobutskii /

GOL'DER, G. A.

V 14669* Physical-Chemical Study of Lithium Peroxide. Fiziko-khimicheskoe iuchenie perekisi litia. (Russian.) T. V. Rode,
T. A. Dobrynina, and G. A. Gol'der. Izvestia akademii nauk
SSSR, otdelenie khimicheskikh nauk, 1955, no. 4, July-Aug.,
p. 611-621.
Includes graphs, tables, diagram. 29 ref.

(2)

Z-ray investigation of the structure of 4-nitro-N-(
dimethylaminomethyl)benzonitrile. G. S. Zeldecker and G. M. Hartman
Karpov Inst. Pure Chem. Moscow, USSR
Received October 6, 1977
The title compound belongs to the monoclinic system; $a = 7.51$, $b = 14.8$, $c = 7.25$ Å, $\beta = 64^\circ$, the no. of mol. in the elementary cell is 4, and $d_{\text{c}} = 1.08$. The x-ray $d = 1.72$ Å/C. The angle $\text{C}_6-\text{C}_7-\text{C}_8$: The nitro group is at an angle of 61° to the benzene ring, the length of the C-N bond is 1.44 ± 0.03 Å. In the azido group the C-N spacing is 1.37 ± 0.03 Å, which permits the assumption of a coplanar arrangement of this group with the benzene ring. W. M. Stremberg

M. Hartman

GOL'DER, G. A.

✓ Physicochemical investigation of the system sodium superoxide-sodium oxide. T. V. Rode and G. A. Gol'der.
Zavod. Akad. Nauk S.S.R., Odz. Khim. Fiz. No. 1, p. 299-308. — Na₂O in an atm. of dry O₂ decomps. at approx. 120°; in dry, CO₂-free air it decomps. at 80-90°. On thermal decompn. Na₂O gives a succession of solid solns. with the limiting compd., Na₂O₂. Na₂O₂ decomps. endothermically with the production of Na₂O, at 250° in O₂ and at 215° in CO₂-free air. Na₂O₂ decomps. slowly, beginning at approx. 350°; it melts at 510°. At 545° Na₂O₂ decomps. to the limiting compd., Na₂O. The existence of a sequence of continuous compds. was detd. in the system Na₂O-Na₂O with limits varying from Na₂O to Na₂O₂. Under the conditions the compd. Na₂O was not detd., but a limiting solid soln. Na₂O₂ in which decomps. immediately to Na₂O was found. J. M. Widén.

Inst.-Gen. & Inorg. Chem. m. N. S. Kurnakov, A.S.U.S.S.R.

✓ X-ray determination of the structure of indigo and thindigo. B. A. Gafurov, G. S. Zhdanov, and O. A. Grigor'eva. *Zh. struk. soed.* 1960, 1, 13-16 (1960). -- The Baeyer-Villiger formula of indigo does not explain the absence of geometric isomers, the deep color, and the high chain and physical reactivity of this mol. Many hypotheses were advanced (Kuhn, C.A. 16, 6, 42; Hodgson, C.A. 40, 8449) to give an improved formulation. The authors submitted indigo and obtained good crystals. The elementary cell dimensions of indigo are: $a = 2.08 \pm 0.01$, $b = 5.74 \pm 0.05$ Å, $c = 11.18 \pm 0.01$. In thindigo they are: $a = 2.02 \pm 0.02$, $b = 3.92 \pm 0.02$, $c = 5.9 \pm 0.04$ Å. The space group is $P\bar{1}$. The electronic d. projection on (010) of the indigo cell, the atomic approximation method delivered the following coordinates: C₁: $x = 0.404, y = 0.235, z = 0.003$; C₂: $x = 0.870, y = 0.128$; C₃: $x = 0.370, y = 0.188$; C₄: $x = 0.226, y = 0.214$; C₅: $x = 0.056, y = 0.177$; C₆: $x = 0.071, y = 0.115$; C₇: $x = 0.217, y = 0.091$; S: $x = 0.229, y = 0.008$; O: $x = 0.036, y = 0.108$ (C₄ number); N: $x = 0.103, y = 0.703, z = 0.142$. The mol. includes with (010) angle of 37°. In thindigo the inclinations to all 3 planes of the elementary cell are considerably: the planes of the benzene ring in indigo parallel to (210). A 3-dimensional Fourier synthesis gave the following coordinates in indigo. For C₁: $x = 0.049, y = 0.907, z = 0.020$; C₂: $x = 0.103, y = 0.703, z = 0.142$; C₃: $x = 0.107, y = 0.903, z = 0.138$; C₄: $x = 0.277, y = 0.431, z = 0.219$; C₅: $x = 0.054, y = 0.305, z = 0.205$; C₆: $x = 0.373, y = 0.201, z = 0.037$; C₇: $x = 0.304, y = 0.403, z = 0.115$; C₈: $x = 0.146, y = 0.612, z = 0.039$; N: $x = 0.112, y = 0.700, z = 0.039$; O: $x = 0.050, y = 0.847, z = 0.219$. In indigo complex bonding occurs, also no classic double bonds. The bond angles are intermediate between single and double valence bonds, and

Gribova EA, Zhidkov GS and Gol'den G A

only in the benzene rings but also in the heterocyclic part of the mol. A system of single + carbon alternating is accomplished; an equivalent inner-side constancy is real. The distance N-O in the indigo mol is practically sum of the interatomic radii of both elements. No other bonds occur, but between the CO and NH groups there is a bond of the mol-hydrogen-bond type, and every ring is bonded to 4 surrounding mols by such H bonds. The thermal stability of Indigo is thus explained. The electronic distribution shows a distinctly decreasing tendency from the center to the periphery of the mol.; this illustrates the thermal oscillations of the whole mol around its center of gravity. The absence of cis-trans isomers is also explained by the results of the structure built in the present paper.

25
26

Golder, G. A.

Distr: 44.1/4E3d

✓ Compounds of constant and of variable composition in the sodium superperoxide-sodium chloride system. T. V. S. and G. A. Golder. Proc. Acad. Sci. U.S.S.R., Chem. 116, 635-8 (1966) (English translation). Ser. C, 51, 14480c.

RODE, T.V.; GOL'DER, G.A.

Compounds of constant and variable composition in the NaO_{1-x}-Na₂O system. Dokl. AN SSSR 110 no.6:1001-1004 O '56. (MLRA 10:2)

1. Institut obshchey i neorganicheskoy khimii imeni
N.S. Kurnakova Akademii nauk SSSR. Predstavлено akademikom
I.I. Chernyayevym.
(Sodium oxides)

AUTHOR: Ozarov, A.P., Gol'der, G.A. and Zhdanov, T.S. 70-2-3/24

TITLE: An X-ray structural investigation of the oxygen vanadium
bronzes of sodium and potassium $\text{Me}_{0.55}\text{V}_2\text{O}_5$. (Rontgenograf-
icheskoye issledovaniye struktury kislotosavnih vanadiyevykh
bronz natriya i kaliya $\text{Me}_{0.55}\text{V}_2\text{O}_5$.)

PUBLICATION: "Kristallografiya" (crystallography), 1977, Vol. 22,
No. 2, pp. 217-225 (U.S.S.R.)

ABSTRACT: Experimental. The valency state of V in broches and in
vanadium-sulfur-oxygen catalysts is particularly of interest.
Crystals of composition $(\text{K},\text{Na})_2\text{O}\cdot\text{V}_2\text{O}_4\cdot5\text{V}_2\text{O}_5$ were obtained as
black laths having a blue metallic lustre. They showed a
large number of faces including the simple forms 102, 101,
100, 001 variously developed. X-ray photographs assigned
them to the space class C_{2h}^+ . Weissenberg and oscill-
ation photographs (11.456 cm diameter camera) with Fe radia-
tion give unit cell dimensions $a = 10.039$, $b = 3.605$, $c = 15.355$ Å
(all ± 0.005 Å) and $\beta = 109^\circ 12' \pm 5'$, for the sodium compound
 $\text{Na}_{0.55}\text{V}_2\text{O}_5$. This gives $V = 524.2$ Å³. The compound $\text{K}_2\text{V}_{12}\text{O}_{30}$
had $d_{\text{obs.}} = 5.67$ g/cm³ giving $\rho = 1$ (0.97). $d_{\text{calc.}}$ is then 5.60.

Card 1/3

Available: Library of Congress

70-2-3/24

An X-ray diffraction investigation of the oxygen tungsten
bronze of sodium was carried out by $\text{Ko}_{0.55}\text{W}_{0.45}\text{O}_3$. (Cs^+ .)

The possible space groups (from the extinction) were $\text{A}2/\text{n}$, $\text{A}/2$ and Am . On the basis of a knowledge of the crystal chemistry of the oxides of V, Mo and W and of the W bronzes the group $\text{A}2/\text{n}$ was chosen. This is confirmed by the dimension t which leads to the expectation of octahedra or trigonal bipyramids (see V.P. Ozerov - Zap. Akad. Nauk., 951, 1955). Using Mo radiation 500 reflections were measured from reticulum pictures using comparison scales. No extinction corrections were applied. $\text{P}(h, k)$ was constructed. A.D. Wadsley's determination of the structure of $\text{Na}_2\text{V}_5\text{O}_13$ helped in solving this Patterson projection. Projections for both Na and K bronzes were constructed.

Several atoms overlap and this was the reason for repeating Wadsley's work. The Fourier section at $y = 0$ was calculated giving interatomic distances for the K bronze very close to those found by Wadsley (Acta Cryst. 10, 615, 1957) for the Na bronze. A table of interatomic distances is given. Slight differences naturally occur in the Mo-O distances (as observed for K bronze first followed by Wadsley's value for the Na bronze); for K bronze first followed by Wadsley's value for the Na bronze); Mo-O₁ (2.70, 2.46); Mo-O₂ (2.49, 2.75); Mo-O₃ (2.56, 2.51);

Card 2/3

70-2-3/24

An X-ray structural investigation of the oxygen valence bounces
of sodium and potassium $\text{Na}_0.5\text{V}_2\text{O}_5$. (cont.)

$\text{Na}-\text{O}_3$ (1.77; 1.51); $\text{Na}-\text{Na}$ (1.65, 2.22). The geometry of the
structures is discussed. The structure is built from strongly
distorted VO_6 octahedra. The distortion is so great that cer-
tain groupings are better regarded as trigonal bipyramids. The
polyhedra differ greatly from themselves V-C distances oscillat-
ing to 1.77, 1.77, 1.77, 2.00 and 2.35 Å. There is a strong
correspondence with the structure of the V oxides. The alkali
atoms lie in canals between the octahedra each surrounded by
seven oxygens. Seven-fold co-ordination is rare but is also
found in the ion $(\text{LiF}_6)^{2-}$ and in Si_3N_4 .

Card 3/3 There are 5 figures, 2 tables and 20 references, 10 of which are
Slavic.

ASSOCIATION: Ya.V. Samoylov Scientific Institute for Fertilisers and
Insecto-fungicides. (Nauchnyy Institut po Dobrobytym i Insekt-
fungisidam im Ya.V. Samoylova)

SUBMITTED: September 21, 1976.

AVAILABLE: Library of Congress

A Radiographic Structural Examination of
Naphthazarine

20-118-6-23/43

in the elementary mesh conform the assumption (reference 1) that a center of symmetry exists in the molecule of the crystals of the 1st modification. The introduction of an inner hydrogen compound O . . . H-O in the conjugated bond-system must have caused an essential change of the π -electronic interaction in the whole molecule. This must, in return, lead to a redistribution of the electronic density in the molecule. A complete radiographic analysis of the crystals of this modification was interesting therefore. The lengths of the bonds between the atoms in the molecule were computed (II) from the atomic coordinates computed from ρ (Okl)(table 2). The computations of the distances between the atoms were carried out under the assumption that the molecule of the surface yz lies parallel. The angle formed by the bond-line $C_9 - C_{10}$ with the y-axis of the mesh, is 50° . The smallest distance between the carbon- and oxygen-atoms in various molecules is 3,10 Å. The results of the radiographic structural analysis confirm the presence of a center of symmetry in the 1st modification of naphthazarine. As mentioned above, all 3 modifications precipitate simultaneously with the crystallization of the solution: 2 centro-symmetrical ones (A), and a none-centro-symmetrical one (B). The

Card 2/4

A Radiographic Structural Examination of Naphthazarine 20-118.4-23/43

recrystallization of each of these modifications leads in return to the formation of all these 3 modifications, though one of them prevails largely. It may thus be presumed that the transition of an isomer of an A-structure into an isomer of a B-structure (and viceversa) takes place. This transition is explained with scheme III and was presumed in reference 4. The orientation in space of the molecule in the yz-surface achieved by the authors, is very similar to that for the centro-symmetrical modification 2) given in reference 3. A three-dimensional synthesis is required for determining the 3rd coordinate x and for defining precisely the obtained results.
There are 1 figure, 2 tables, and 4 references, 1 of which is Slavic.

ASSOCIATION: Physico-Chemical Institute imeni L. Ya. Karpov
(Fiziko-khimicheskiy institut im. L. Ya. Karpova)

PRESENTED: November 20, 1957, by N. V. Belov, Academician.

SUBMITTED: August 16, 1957.
Card 3/4

2p-119-1-23/54

AUTHORS: Dokunikhin, N. S., Gol'der, G. A., Mhdanov, G. S.

TITLE: The Radiographic Investigation of 1,4-di-Anilido-Anthraquinone and 1,4-Dimesido-Anthraquinone (Рентгенографическое исследование 1,4-дianilidoантрахинона и 1,4-dimesidoантрахинона)

PERIODICAL: Doklady Akademii Nauk SSSR, 1958, Vol. 111, Nr 1,
pp. 87 - 89 (USSR)

ABSTRACT: Sulfo acids of 1,4-di-(aryl amino)-anthraquinone form an important group of solid dyes for wool. The majority of the 1,4-di-(aryl amino)-substitutes of anthraquinone are green. An exception is made by the derivatives in which all hydrogen atoms, in an ortho-position, of the aryl-residues are substituted. Such compounds as well as the corresponding alkyl-amino-and hydro-aryl-amino-derivatives have an intensive bright-blue color. In the presence of methyl-ethyl-groups or of bromine atoms in all ortho-positions of the phenyl residues or in the position of 2,3-anthraquinone respectively

Card 1/6

The Radiographic Investigation of 1,4-di-Anilido-Anthraquinone and
1,4-Dimesido-Anthraquinone

cycles by hydrogen and is caused by the interaction of the conjugation-system (Reference 4). It would be desirable to find a direct proof of the flat structure of the molecules of 1,4-di-(arylamino)-anthraquinone in the absence of spatial difficulties. For the purpose of deciding the problem of coplanarity of the benzene nuclei with the plane of the basic part of the molecule, crystals of both compounds mentioned in the title were radiographically measured. The results are given in table 1. From the dimensions of the elementary cell of the first compound can be assumed that the basic part of the molecule is here entirely or almost parallel with the ac-plane, as axis b is the shortest one (a,73 Å). From the conditions of symmetry of the spatial group

$$C_{2h}^5 = P\bar{2}_1/c$$

Card 3/6

70-110-1-23/52

The Radiographic Investigation of 1,4-Di-Anilido-Anthraquinone and
1,4-Dimesido-Anthraquinone

follows that a slip plane with a displacement along axis c runs vertical to axis b. thereby the molecules occurring in the unit cell are orientated in layers which are perpendicular to axis b. A variant of this orientation is shown by figure 4. It admits a slight turn of the benzene nucleus in relation to the other part of the molecule as well as a certain possible turn of the entire molecule in relation to the plane ac. Thus the packing of the molecules in the crystal does not require an additional change of the angle of rotation of the benzene nucleus as compared to the free molecule. The shortest axis in the crystal of the second compound is the a-axis (7,75 Å). Its length corresponds to the dimensions of the benzene nucleus up to the CH_3 -groups connected with it (9,9 Å). A solid packing of molecules in the crystal and the fulfilment of the conditions of symmetry of the spatial group for molecules of the second compound

Card 4/6

144-14-23,52

The Radiographic Investigation of 1,4-Di-Anilido-Anthraquinone and
1,4-Dimesido-Anthraquinone

of the methyl groups to all meta-positions of the benzene
nucleus creates so great steric difficulties that the
coming out with the interquinone cycles from the coplanarity
amounts to almost 90°. Thereby the inner-molecular linkage
is considerably weakened. There are 7 figures, 1 table,
and 5 references, all of which are quoted.

ASSOCIATION:Nauchno-issledovatel'skiy in-titut or organicheskikh polupro-
duktov i krasiteley im. K. Ye. Voroshilova (The Scientific Re-
search Institute of Organic Semiproducts and Dyes imeni
K. Ye. Voroshilova). Nauchno-issledovatel'skiy Fiziko-
khimicheskiy institut im.L. Ya. Karpova (Scientific Phys-
ical-Chemical Research Institute named L. Ya. Karpov)

PRESENTED: November 2c, 1957, by N. V. Belov, Member, Academy of

Sciences, USSR

SUBMITTED: August 16, 1957

Card 6/6

PAGE 1 BOOK SPECIFICATION 39-439

Moscow, Vsesoyuznnoye Khimicheskoye Institut

Problemy Fizicheskoy Kinetiki i Trifazovoy Fiziki (problems in Physics

Chemical Kinetics and Three-phase Physics), Moscow, Naukova Dumka,

Osnovnyi Redaktor: Yu. M. Vaynshteyn, Doctor of Chemical Sciences.

Akhiezer, Ya. M., Kolobratin, D. I., L'vov, G. A., and others.

Redaktsionnyy Tsentralnyy Uchebnyy Komitet po Osnovnym Sistemam

Doctor of Chemical Sciences; V. M. Gromov, Institute of General Chemistry,

Candidate of Chemical Sciences; V. S. Chibisova, Institute of General Chemistry;

Candidate of Chemical Sciences; E. A. Moshkov, Institute of

Ekon. V. S. Sipkin.

REMARKS: This collection of articles is intended for physical

chemistry. The collection is the second issue of the "Trifazovaya

Kinetika i Trifazovaya Fizika" series. It contains 17 articles which involve

discussions of the kinetics of reversible reactions which involve

reactions of the formation of hydrogen sulfide and the formation of

hydrogen sulfide exchange systems and systems of electro-

chemical transformations.

14

Editor-in-Chief: S. V. A. Kostomarov, Yu. I. Pecherskiy,

V. M. Gromov, Institute of General Chemistry, Academy of Sciences of the USSR, and the Department of General

Chemical Kinetics (Izhevsk). Redaktsionnyy Sipkin.

REMARKS: This collection of articles is intended for physical

chemistry. It contains 17 articles which involve discussions of

the kinetics of reversible reactions which involve the formation of

hydrogen sulfide exchange systems and systems of electro-

chemical transformations.

15

Editor-in-Chief: S. V. A. Kostomarov, Yu. I. Pecherskiy, Yu. I.

V. M. Gromov, Institute of General Chemistry, Academy of Sciences of the USSR, and the Department of General

Chemical Kinetics (Izhevsk). Redaktsionnyy Sipkin.

REMARKS: This collection of articles is intended for physical

chemistry. It contains 17 articles which involve discussions of

the kinetics of reversible reactions which involve the formation of

hydrogen sulfide exchange systems and systems of electro-

chemical transformations.

16

Editor-in-Chief: S. V. A. Kostomarov, Yu. I. Pecherskiy, Yu. I.

V. M. Gromov, Institute of General Chemistry, Academy of Sciences of the USSR, and the Department of General

Chemical Kinetics (Izhevsk). Redaktsionnyy Sipkin.

17

Editor-in-Chief: S. V. A. Kostomarov, Yu. I. Pecherskiy, Yu. I.

V. M. Gromov, Institute of General Chemistry, Academy of Sciences of the USSR, and the Department of General

Chemical Kinetics (Izhevsk). Redaktsionnyy Sipkin.

18

Editor-in-Chief: S. V. A. Kostomarov, Yu. I. Pecherskiy, Yu. I.

V. M. Gromov, Institute of General Chemistry, Academy of Sciences of the USSR, and the Department of General

Chemical Kinetics (Izhevsk). Redaktsionnyy Sipkin.

19

Editor-in-Chief: S. V. A. Kostomarov, Yu. I. Pecherskiy, Yu. I.

V. M. Gromov, Institute of General Chemistry, Academy of Sciences of the USSR, and the Department of General

Chemical Kinetics (Izhevsk). Redaktsionnyy Sipkin.

20

Editor-in-Chief: S. V. A. Kostomarov, Yu. I. Pecherskiy, Yu. I.

V. M. Gromov, Institute of General Chemistry, Academy of Sciences of the USSR, and the Department of General

Chemical Kinetics (Izhevsk). Redaktsionnyy Sipkin.

21

5(2)

AUTHORS: Kost, M. Ye., Gol'der, G. A.

SOV/78-4-7-4/44

TITLE: The Crystal Structure and Density of Cerium Hydrides (Kristallicheskaya struktura i plotnost' gidridov tseriya)

PERIODICAL: Zhurnal neorganicheskoy khimii, 1959, Vol 4, Nr 7,
pp 1488-1490 (USSR)

ABSTRACT: Cerium hydrides with a composition of from $\text{CeH}_{0.2}$ to CeH_3 were investigated. The trihydride was produced in an apparatus described in an earlier paper (Ref 5). The samples poor in hydrogen were obtained by heating and by sucking off the liberated hydrogen. The composition of the hydrides was determined by measuring the hydrogen liberated in a solution of hydrochloric acid. The Debye powder patterns were recorded by means of the camera RKD. The values of the lattice periods are given by table 1. Up to the composition $\text{CeH}_{1.5}$ two cubical face-centered lattices exist, which correspond to the metal Ce and to the dihydride. The sample $\text{CeH}_{1.97}$ shows a phase in the period 5.55 Å. A further increase of the hydrogen content leads to a reduction of the period to 5.53 Å at $\text{CeH}_{2.73}$. If the com-

Card 1/2

SOV/78-4-7-4/44

The Crystal Structure and Density of Cerium Hydrides

position CeH_3 is approached, the lines widen, so that exact calculation of the lattice period is rendered difficult. Because of the great sensitivity of cerium hydrides to vestiges of water, density was determined in an apparatus (Fig. 1), in which argon was used as a pyknometric substance, and in which the volume of the sample was determined on the basis of a variation of pressure according to the Boyle-Mariotte law. The density of the various hydrides is given by table 2. It decreases up to the compound CeH_2 , after which it rises somewhat up to CeH_3 .

Figure 2 gives a graphical comparison of density variations with the X-ray pictures, the curve of which shows the presence of two phases (metallic cerium and CeH_2) up to the compound CeH_2 .

The lines of the metallic Ce then vanish. The phase with the periods 5.645 - 5.612 Å, which was observed by M. C. Auphas-sorho (Refs 3,4) could not be found. There are 2 figures, 2 tables, and 8 references, 2 of which are Soviet.

SUBMITTED: April 4, 1958

Card 2/2

34(4), 5(1)

347/70-21-27/71

UDC#S:

Gol'dier, G. A., Mil'kov, G. S., Ivanov, V. M., Firsov, L. N.,
G. N., Chugun, Ye. A.

TITLE:

The Use of X-Ray Phase Analysis in Spray Technology (Primenenie rentgenovskogo fazychnogo analiza v tekhnike volnoklyazhnoi)

PERIODICAL:

Zavodskaya Laboratoriya, 1971, Vol. 10, No. 1,
pp. 101 - 107 (USSR)

ABSTRACT:

The present paper lists the results of investigations carried out by the laboratories of the plant "Vzvodnyy zavod", Yaroslavl', GIFI-4, IRF-1, "Krasnyy Shchit", Leningrad, Fiziko khimicheskiy institut im. I. Ya. Karpeva (Leningrad), Chemical Institute imeni I. Ya. Karpeva) and others. A standard isometric X-ray apparatus was used. Since the X-ray phase analysis has a low sensitivity for impurities, it should not be used for jet spraying of dilute mixtures (less than 1-3%). Analyses of titanium dioxide are described: 1) A study of titanium dioxide in water; 2) the optimum production conditions of utilized form of a lead oxide if one found by X-ray to consist of the

Card 1/2

The Use of X-Ray Phase Analysis in Chemical Technology J V (O-2)-2-27/71

yellow substance did not correspond to the usual red tetragonal modification of PbO, but to the yellow rhombic modification, and that the color was due to a polymorphous change.
3) By means of X-ray analysis it was possible to simplify the production control of active pyrolusite of the Gdr. 4) Examinations of domestic and foreign refracting types were carried out to determine the dispersion degree of the iron oxide. 5) Moreover, the production of thiocure was controlled with regard to dicyan-diamide. 6) The X-ray analysis was also successfully used in the examination of luminophores, which can also be applied for the examination of other substances (e.g., catalysts).

ASSOCIATION: Nauchno-issledovatel'skiy fiziko-khimicheskiy institut im L. Ya. Karpova (Scientific Research Institute of Physical Chemistry imeni L. Ya. Karpov)

Card 3, 2

GOL'DEN, G.A. [translator]; DUDAREV, V.Ya.[translator]; SOLOV'YEV,
S.P.[translator]; ZHDANOV, G.S., red.; LATIN, S.I., red.;
BELEVVA, M.A., tekhn. red.

[Annihilation of positrons in solids] Annigiliatsiia po-
zitronov v tverdykh telakh; sbornik statei. Moskva, Izd-vo
inostr. lit-ry, 1960. 228 p. (MIRA 15:3)
(Positrons)

RODE, T.V.; GOL'DER, G.A.; ZACHATSKAYA, A.V.

Interaction of sodium peroxide and sodium superoxide with sodium bicarbonate. Zhur. neorg. khim. 5 no.3:535-539 Mr'60.

(MIRA 14:6)

(Sodium peroxide)
(Sodium superoxide)
(Sodium carbonate)

MIRKIN, Lev Iosifovich; UMANSKIY, Ya.S., prof.. red.; GOL'DER, G.A., red.;
MAKAROV, Ye.F., red.; MURASHOVA, N.Ya., tekhn. red.; TUMARKINA, N.A.,
tekhn. red.

[Manual on X-ray diffraction analysis of polycrystals] Spravochnik po
rentgenostrukturnomu analizu polikristallov. Pod red. IA.S.Umanskogo.
Moskva, Gos. izd-vo fiziko-matem. lit-ry, 1961. 863 p. (MIRA 14:8)
(X-ray crystallography)

GOL'DER, G.A.; TODES-SELEKTOR, Z.V.; BOGDANOV, S.V.

Structure of benzofuroxan. Zhur.strukt.khim. 2 no.4:478-479
Jl-Ag '61.
(MIRA 14:9)

1. Nauchno-issledovatel'skiy fiziko-khimicheskiy institut imeni
L.Ya. Karpeva i Gosudarstvennyy nauchno-issledovatel'skiy insti-
tut organicheskikh poliproduktov i krasiteley imeni L.Ye.
Voroshilova.

(Benzofuroxan)

CHETKINA, L.A.; GOL'DER, G.A.; ZHDANOV, G.S.

X-ray diffraction study of dihalogen derivatives of anthraquinones. Kristallografia 6 no.4:628-629 Jl-Ag '61. (MIRA 14:8)

1. Fiziko-khimicheskiy institut imeni L.Ya.Karpova i Moskovskiy gosudarstvenny universitet imeni M.V.Lomonosova,

(Anthraquinone) (X-ray crystallography)
(Halogen compounds)

S/192/62/003/002/003/004
5267/5301

AUTHORS: Chamova, V.N. and Gol'der, G.A.

TITLE: X-ray investigation of the potassium carbonate peroxyhydrate $K_2CO_3 \cdot 5H_2O_2$

PERIODICAL: Zhurnal strukturnoy khimii, v. 5, no. 2, 1962,
215 - 216

TEXT: One of the authors (Ref.2: Makarov, S.Z., Chamova, V.N., Izv. Akad. Nauk SSSR, Otd. khim. nauk, v. 3, 1958, 103) discovered a static solid phase of the above composition. X-ray analysis of this substance was carried out by the powder and monocrystal method, and the crystal was found to belong to the orthorhombic system. The parameters of the elementary cell are : $a = 5.50$, $b = 6.04$, $c = 17.8$ Å. The density of the peroxyhydrate was measured ($\rho = 2.02$). There are four molecules in the elementary cell, and the calculated density is $\rho = 2.01$.

Card 1/2