

MIDOYAN, A. A.

Midoyan, A. A. - "On the problem of a fractured ankle bone," Sbornik nauch. trudov (In-t gematologii i perelivaniya krovi. Fak. khirurg. klinika Yerevansk. med. in-ta), III, 1948, p. 177-95

SO: U-4355, 14 August 53, (Letopis 'Zhurnal 'nykh Statey, No. 15, 1949)

BELOTSKIY, D.P.; NOVAL'KOVSKIY, N.P.; MIDONOVA, N.N.

Study of the interaction of sulfuric acid and glycerol in aqueous solution by methods of physicochemical analysis. Izv.vys.ucheb. zav.; khim.i khim.tekh. 4 no.6:1035-1037 '61. (MIRA 15:3)

1. Chernovitskiy gosudarstvennyy universitet, kafedra neorganicheskoy khimii.
(Sulfuric acid) (Glycerol)

MIDLER, T.L.; GONCHARUK, A.N.

Mud therapy in the early stages in poliomyelitis in children. Report
No.2. Vop. kur., fizioter. i lesh. fiz. kul't. 25 no.2:142-145 Mr-
Ap '60. (MIRA 13:9)

1. Iz kafedry detskikh bolezney lechebnogo fakul'teta (zav. -
dotsent V.P. Chernyuk) i infektsionnykh zabolevaniy detskogo
vzrastu (zav. - dotsent N.G. Stepina) Odesskogo meditsinskogo
instituta imeni N.I. Pirogova (dir. - prof. I.Ya. Dayneka) i Odesskoy
gorodskoy infektsionnoy bol'nitsy (glavnyy vrach S.T. Kolesnikov).
(POLIOMYELITIS) (BATHS, MOOR AND MUD)

GONCHARUK, G.M. [Honcharuk, H.M.]; MIDLER, T.L.

Importance of taking the skin temperature in the recognition of various forms of poliomyelitis. Ped., akush. i gin. 22
no.3:26 '60. (MIRA 14:4)

1. Kafedra infektsiynikh zakhvoryuvan' dityachogo viku (zav. - dotsent N.G. Stepina [N.H.Stepina]) ta kafedra dityachikh khvorob likulval'nogo fakul'tetu (zav. - dotsent V.P.Chernyuk) Odes'kogo derzhavnogo medichnogo instituttu im. M.I.Pirogova (direktor zasluzh.diyach nauki prof. I.Ya.Deyneka) i mis'ka klinichna infektsiyna likarnya (golovniy likar L.Zhidovlenko).

(POLIOMYELITIS) (SKIN)

MIDLER, T. L.: Master Med Sci (diss) -- "The effectiveness of mid therapy for children suffering from poliomyelitis (as a function of the time of infection)". Odessa, 1958. 13 pp (Odessa State Med Inst im N. I. Pirogov), 220 copies (KL, No 16, 1959, 110)

STEPINA, N.G. [Stepina, N.H.], dots.; MIDLER, T.L.; GONCHARUK, G.M.
[Honcharuk, H.M.]

Fangoththerapy for poliomyelitis patients during initial hospitalization.
Ped., akush. i gin. 19 no.4:11-14 '57. (MIRA 13:1)

1. Kafedra infektsionnykh bolezney detskogo vozrasta (zav. - dots.
N.G. Stepina), Klinika detskikh bolezney lechebnogo fakul'teta (zav. -
dots. V.P. Chernyuk) Odesskogo gosudarstvennogo meditsinskogo insti-
tuta im. M.I. Pirogova (direktor - prof. I.Ya. Deyneka) i gorodskaya
infektsionnaya bol'nitsa (glavnyy vrach - S.T. Kolesnikov).
(BATHS, MOOR AND MUD) (POLIOMYELITIS)

MIDLER, T.L.

SOSNOVA, P.S.; MIDLER, T.L.

Burn reaction as an index of reactivity in children. *Pediatrics*,
Moskva no. 4:71 July-Aug. 1952. (CJML 22:5)

1. Of the Department of Children's Diseases of the Therapeutic
Faculty of Odessa Medical Institute.

MIDIVANI, A.G.; SADYKHOV, Yu.V.

Stuck pipes and their causes. Azerb. neft. Khoz. 41 no.1:
17-18 Ja '62. (MIRA 16:7)

(Oil well drilling)

MINITESCU, E.

The Iron, Manganese, and Argillaceous Refractory Clay Mining Sectors.
Revista Minelor (Mining Journal), 48:247: Aug 55

GVELESIANI, G.G.; MIDELADZE, I.S., red.; BOKUCHAVA, T.P., red. izd-va;
BOKERIYA, E.B., tekhn. red.

[Principles of the socialist distribution of production] Ob
osnovakh sotsialisticheskogo razmeshchenia proizvodstva.
Tbilisi, Izd-vo Akad. nauk Gruzinskoi SSR, 1961. 135 p.
(MIRA 15:3)

(Russia--Economic policy) (Industries, Location of)

RACV.MNU, V. prof.; LAZARUS, N., dr.; HODMAN, H., dr.; POE, V., dr.;
LUTTENBERG, H., dr.; ANSCHUTZ, R., dr.; HIRSH, EDWIN, dr.

Considerations on the principal methods of prosthetic denture
in children. *Otolaryngologic (Basel)* 10 no.1/2:20
Jan-1965.

RUMANIA

MIDAN, Florica, Dr.

O.R.L. Clinic, II, "Colțea" Hospital (Clinica O.R.L., II,
Spitalul "Colțea"), Bucharest. . .

Bucharest, Viata Militara, No 12, 15 Jun 63, pp 861-864

"Antistreptolysin O Determinations."

(1)

ORLOV, D.S.; GLEBOVA, G.I.; MIDAKOVA, K.Ye.

Analysis of the distribution of ferric compounds and humus in
the soil profile based on the curves of spectral brightness.
Nauch. dokl. vys. shkoly; biol. nauki no.1:217-222 '66.
(MIRA 19:1)

1. Rekomendovana kafedroy pochvovedeniya Moskovskogo
gosudarstvennogo universiteta. Submitted December 2, 1964.

MIDAK, Edmund; ZIELINSKI, Marian

Results of rapid pregnancy immunological test with progesterone.
Wiad. lek. 18 no.15:1251-1253 1 Ag 1966.

1. Z Oddz. Pol.-Gin. Centr. Szpitala Klin. Ministerstwa Spraw
Wewnętrznych w Warszawie (Kierownik: dr. med. M. Zielinski).

MIDAK, Edmund; KROCHMAL, Andrzej

Treatment of sample of the wax is stated after comparison with
a note on the use of organic glass. *Wied. Roz. 1965. 11. 15. 1965*
15 Je '65.

1. Z Oddzialu Pol.-Gin. Szpil. Min. Wn. Sprac. Swan. w Warszawie
(Kierownik: dr. nauk M. Wielinski).

MIDAK, Edmund

A case of early cervical pregnancy. Pol. tyg. lek. 19 no.45:
1742-1743 N 9'64

1. Z Oddziału Położniczo-Ginekologicznego Centralnego Szpitala
Klinicznego MSW w Warszawie (Kierownik: dr. med. Marian Zieliński).

MIDAK, Edmund

Analysis of indications for cesarean section in the era of antibiotics according to observations on 14,000 cases. Pol. tyg. lek. 19 no.39:1492-1495 28 S '64

1. Z Oddziału Położniczo-ginekologicznego Centralnego Szpitala Klinicznego MSW w Warszawie (Kierownik dr. med. Marian Zielinski).

ILLEGIBLE

MICZYNSKI, Kazimierz

The Rothamsted Experiment Station. Postepy nauk roln 7 no.4:109-121
Jl-Ag '60. (EEAI 10:2)

1. Zaklad Fizjologii Roslin Polskiej Akademii Nauk. Pracownia
Wirusologii, Krakow.
(Great Britain--Agriculture) (Poles in Great Britain)

MICZYNSKI, Kazimierz

Agricultural studies at Cornell University in the United States.
Postepy nauk roln 7 no.2:131-142 Mr/Ap '60. (EEAI 9:10)

1. Katedra Botaniki Wyzszej Szkoły Rolniczej, Krakow.
(United States--Agriculture)

MICZYNSKI, K.

P O L . E

✓ Susceptibility to bunt of wheat varieties cultivated in Poland.
K. Mieczynski (*Roczn. Nauk Rol.*, 1953, 67, A, [2], 89-90).--The
majority of the 89 winter varieties of wheat grown in Poland were
susceptible to bunt. Four of the varieties were highly resistant.
Some varieties which showed no to slight infection in the first
year, were moderately to severely infected in subsequent years.
Most spring wheat varieties were highly susceptible to infection.
The American variety Hope showed the least infection.

A. H. CORNFIELD

MICZYNSKI, JAN

Sugar-Beets

DECEASED

C. 62

1964

0
CATEGORY : PLANT DISEASES. Diseases of Cultivated Plants.
ABS. JOUR: Ref Znan-Biologiya, No. 5, 1959, No. 20621
AUTHOR : Miczynska, Zofia; Wnekowski, Stanislaw
INST. : Inst. of Plant Protection
TITLE : Preliminary Research on Species Composition
in Fusarium Infection of Corn.
ORIG. PUB.: Roszn. nauk rolniczych, 1957, A77, No.3,
357-371
ABSTRACT : It has been demonstrated by research made at
the laboratories of physiography and statis-
tics of the Institute of Plant Protection
(in the city of Pulava) th t stunting of corn
plants caused by the species of the genus
Fusarium (F. culmorum, F. sporotrichiella v.
poal, F. moniliforme v. subglutimans) are a
serious danger in cultivating this corn in
Poland.

CARD: 1/1

COUNTRY : Poland
CATEGORY : Plant Diseases. Diseases of Cultivated
Plants.
ABS. JOUR. : RZbiol., No. 12, 1958, No. 54000
AUTHOR : Nieczyńska, Zofia
INSE. : Not given
TITLE : Fusarium Infection in Corn
ORIG. PUB. : Postępy nauk. roln., 1957, 4, No. 1, 111-118
ABSTRACT : Information is given on fusarium diseases
which attack corn in various lands, with
description presented of the symptoms of
these infections. The occurrence of Fusa-
rium poae, F. graminearum, F. culmorum has
been determined in Poland on corn suffering
from fusarium disease. This study was made
at the Institute for Plant Protection in
Pulawy. --V.T. Vergovskiy

CARD: 1/1

POLAND/General and special Zoology. Insects

2-2

Obs Jour : Pr. Zwier. - Zool., 1957, 10, 15, 195, 20, 204-211

Author : Nieczynska-Zonia
Inst : "
Title : Observations of Insect Mycoses

Orig Pub : Polskie pismo entomol., 1957, 8, No 1, 57-59

Abstract : During the 1954-1955 periods of mass appearance of certain caterpillars, observations indicated that development of epiphytes of certain fungi. The fungus, Beauveria sp., developed upon the hibernating caterpillars of the brown-tailed moth, but the fatality rate did not exceed 1-5%. Judging from materials collected in the following summer, the activity of the fungus never reached epidemic proportions. The disease caused by the aquatic calicoid fungus, which developed in the summer of 1955 among caterpillars of the brown-tailed moth, did have the character of an epidemic.

Card : 1/2

Country : Poland
CATEGORY : Plant Diseases. Diseases of Cultivated
Plants.
ABS. JOUR. : RZBiol., No. 12, 1958, No. 54023
AUTHOR : Mieczynska, Zofia
INST. : Not given
TITLE : Gummosis in Cotton, Caused by Pseudomonas
malvaearum Sm.
ORIG. PUB. : Acta agrobot., 1956 (1957), 5, 169-173
ABSTRACT : This disease agent was introduced into Poland
on seeds, and the disease is seen at all
locations of cotton sample plantings. The
disease's developmental conditions, damage,
means of spreading, symptoms in all stages
of the plant's development are described.
Agricultural management and seed disinfection
with H₂SO₄ and formalin are recommended. This
study was made at the Institute for Plant
Protection in Pultavy.--V.I. Vergovskiy

CARD: 1/1

MICZULSKI, Stanislaw, mgr

Beginnings and development of the first blast furnace center
in Poland. Wlad hut 18 no.7/8:227-229 J1-Ag '62.

L 00109-66
ACCESSION NR: AP5013188

Orig. art. has: 4 figures, 4 tables, and 5 formulas.

ASSOCIATION: Ustav termomechaniky CSAV, Prague (Institute of Thermomechanics, CSAV)

SUBMITTED: 05Oct64

ENCL: 00

SUB CODE: ME

NO REF SOV: 000

OTHER: 002

3

44.55

Card

dg
2/2

L. 00109-66 EPF(c)/EWP(j)/ETC(m) RPL RM/WW/JW
ACCESSION NR: AP5013188 CZ/0041/65/000/002/0206/0211

AUTHOR: Miczek, Gerhard (Mitsek, G.) (Engineer)

TITLE: Thermophoresis of solid and liquid particles

SOURCE: Strojnický časopis, no. 2, 1965, 206-211

TOPIC TAGS: aerosol, temperature gradient, high temperature research

ABSTRACT: The importance of thermophoresis, or motion of solid or liquid particles floating in a gaseous medium caused by a temperature gradient, in devices used for energy and chemical conversions at high temperatures is discussed. The practical significance of thermophoresis is due to the fact that the particles tend to deposit on the walls of the device, which are cooler than the gaseous medium. When the gas cools, this aerosol deposit may change the surface conditions and eventually obstruct the orifices of the equipment. Although the motion of particles in a gaseous medium with a temperature gradient is explained by several theories, certain values can be determined by experiment only. The principle of thermophoresis and its properties are discussed on the basis of the work of G. Stetter and K. H. Schmitt, and equipment is described by means of which the thermophoretic velocity at temperatures up to 1500C will be measured.

Card 1/2

MICZEK, Gerhard, inz.

Heat transmission and the transfer of substance. Stroj cas
16 no.2:190-191 '65.

Thermophoresis of solid and liquid particles. Ibid:206-211

1. Institute of Thermomechanics of the Czechoslovak Academy of
Sciences, Prague. Submitted October 5, 1964.

MICZEK, G.

"Quality of cyclone dustbins." Strojirenstvi, Praha, Vol. 4, No. 7, July 1954, p. 500.

SO: Eastern European Accessions List, Vol. 3, No. 11, Nov. 1954, i.O.

HUNGARY

MICZEAN, Izabella, Dr; Janos Hospital-Ambulatory Service (director: TAKO, Jozsef, Dr), Central Laboratory (chief physician: HAMMER, Sarolta, Dr) (Janos Korhaz-Rendelointezet, Kozponti Laboratorium), Budapest.

"Gastrocytological Examinations."

Budapest, Orvosi Hetilap, Vol 107, No 18, 1 May 66, pages 837-838.

Abstract: [Author's Hungarian summary] Based partly on his own experiences and partly on literature data, a brief summary report is given by the author on methods and techniques related to clinical gastrocytological diagnostics including the techniques of obtaining of the samples, their processing as well as the evaluation of the results. 1 Hungarian, 14 Western references.

1/1

KOKAS, Eszter; MICZBAN, Izabella

Effects of guanyl nucleotides on the working capacity, heart and adrenal hypertrophy in rats. Kiserletes orvostud. 10 no.2-3:229-235 Apr-June 58.

1. Orszagos Koranyi Tbc. Intezet klinikai laboratoriuma es Janos Korhaz kozponti laboratoriuma.

(NUCLEOSIDES AND NUCLEOTIDES, eff.

guanylic acid deriv. on musc. work capacity, adrenal & heart hypertrophy in rats (Hun))

(MUSCLES, eff. of drugs on

guanylic acid deriv. on musc. work capacity in rats Hun))

(HEART, eff. of drugs on

guanylic acid deriv. on hypertrophy due to musc. work in rats (Hun))

(ADRENAL GLANDS, eff. of drugs on same)

MICZBAN, IZABELLA

LUDANY, Gyorgy, Dr.; GATI, Tibor, Dr.; MICZBAN, Izabella, Dr.; RIGO, Janos, Dr.;
FOLDVARI, Peter

Novocaine and the pituitary-adrenal system. Orv. hetil. 98 no.37:1018
15 Sept 57.

1. A Budapesti Orvostudományi Egyetem Korleletani Intezete (igazgató:
Sós József egyet. tanár) közleménye.

(PROCAINE, eff.

on adrenal-pituitary stress mechanism in rats (Hun))

(PITUITARY GLAND, physiol.

adrenal-pituitary stress mechanism, eff. of procaine in
rats (Hun))

(ADRENAL GLANDS, physiol.

same)

BANGA, L.,; KOKAS, E.,; MICZBAN, I.,; TORO, I.

Effect of purine nucleotides on the nucleic acid content of the heart during work. Kiserletes orvostud. 7 no.5:511-516 Sept 55.

1. Budapesti Orvostudományi Egyetem Szövet- és Fejlődéstan Intézete és I. sz. Kóronctani és Kísérleti Raktató Intézete.

(NUCLEOTIDES, effects

purine nucleotides, guanylic acid deriv., on nucleic acid in heart during work in rats)

(TISSUE EXTRACTS, effects

heart, embryonic, guanylic acid deriv., on nucleic acid in heart during work in rats)

(ADENYLPHOSPHATE, effects

on nucleic acid in heart during work in rats)

(NUCLEIC ACIDS, metabolism

heart, during work in rats, eff. of adenylypyrophosphate, and guanylic acid deriv. in purine nucleotides & corhormone)

KOKAS, E.; MICZBAN, I.; TORO, I.

Effect of purine nucleotides on working capacity, on the heart and the adrenal glands. Kiserletes orvostud. 7 no.5:501-511 Sept 55.

1. Budapesti Orvostudományi Egyetem Szövet- és Fejlődéstan Intézete.

(NUCLEOTIDES, effects

purine nucleotides, guanylic acid deriv., on working capacity, heart & adrenal in swimming test in rats)

(TISSUE EXTRACTS, effects

heart, embryonic, guanylic acid deriv., on working capacity, heart & adrenal in swimming test in rats)

(ADENYLPYROPHOSPHATE, effects

on working capacity, heart & adrenal in swimming test in rats)

(WORK

musc. work, eff. on heart & adrenal in swimming test in rats)

MICZBAN, I.

3460. Effect of purine nucleotides on working capacity, on the heart and adrenal glands. E. Kokus, I. Miczban, and I. Tord *Acta Physiol. Acad. Sci. Hung.*, 1955, 7, 409-420 (Inst. of Histology and Embryology, Med. Univ., Budapest, Hungary).—250 g. rats in 4 groups as in following paper were forced to swim for 23 days. Swimming performance and body wt. decreased in the saline injected rats whereas it increased in the other 3 groups in the following ascending order: Corhormone, ATP, GAD. Cardiac wt. increased with exercise in the saline group; this increase was prevented in increasing degrees by Corhormone, ATP, and GAD. Adrenal wt. behaved similarly to cardiac wt. It is concluded that GAD is one of the active principles of Corhormone. A. B. L. BAZAN.

3

MICZBAN, I

8489: Action of purine nucleotides on nucleic acid content of heart during exercise. I. Banga, R. Kokas, I. Miczban, and I. Tóro. *Acta physiol. Acad. Sci. hung.*, 1955, 7, 401-407 (Inst. of Histology and Embryology, Medical Univ., Budapest, Hungary).—160 240. Rats were loaded with lead wt. and made to swim in 4 groups. Each group was injected with one of the following, saline, Cor-hormone (trade name of an embryonic heart extract), ATP, or GAD a guanylic acid nucleotide isolated as Na salt from Cor-hormone. (Kokas and Banga, *ibid.*, 1950, Suppl. 1, 24). Swimming took place daily for 10-16 and 17-23 days and the total nucleic acid, RNA, and DNA content of their hearts determined. The 2 latter determinations gave variable results. Total nucleic acid diminished during exercise in the saline group. The diminution was proportional to the severity of the exercise. The nucleic acid content increased or in strenuous overwork returned to normal in the groups injected with the other 3 substances. In the hearts of these latter groups great number of nuclei appeared of the type of young animals.

A. B. L. BEZNAK.

MICZBAN, I.

Recent experimental trends in histology; roentgenspectrography. *Orr.*
hetil. 91 no.29:918-920 16 July 50. (CLML 20:7)

MICZBAN
MICZBAN, I.

Recent experimental developments in histology; optical analysis by
means of the polarization microscope. Orv. hetil. 91:26, 25 June 50.
p. 823-5

CLML 19, 5, Nov., 1950

MICYK, Albin

Talc schists in the Sobotka Mountain region. *Przełgl geolog*
10 no.7:365-366 J1 '62.

MACAROVICI, Const. Gh.; MICU-SEBENIUC, Rodica

Study of complex combinations of diphenyl derivatives. Pt. II.
Rev. chimie Roum 9 no.2:127-130 F '64

1. Laboratory of General Chemistry, University of Cluj.

MACAROVIGI, Const. Gh.; MICU-SMENIUC, Rodica

Study of complex combinations with diphenylic derivatives. Pt. 3.
Studia Univ B-B S Chem 8 no. 1:151-161 '63

1. "Babes-Bolyai" University, Cluj.

POPOVICI, D.Gh.; MICUSAN, V.V.

Research on protein fractions in maternal blood, newborn blood,
and in goat colostrum. Studii cerc biochimie 6 no.1:39-48
'63.


1. Laboratorul de biochimie, Institutul de cercetari zootehnice,
Bucuresti.

*

MICUSAN, V.V.; RUZILA, Lucia

Antigenicity relations between the maternal seric proteins, the colostrum immunoglobulins, and the seric proteins of the new born calf. *Studii cerc biochimie* 7 no.2:213-219 '64.

1. Laboratory of Biochemistry of the Zootechnical Research Institute, and the Institute of Biochemistry, Rumanian Academy, Bucharest. Submitted February 17, 1964.



(1) The purpose of the present work is to determine the explosion limits of
hydrogen-oxygen mixtures at various pressures. The apparatus used for
this purpose is described in detail in the following section. The
mixture is prepared by passing hydrogen and oxygen through a
series of wash bottles and then into the explosion chamber. The mixture
is ignited by a spark gap between two electrodes. The whole apparatus is
maintained at a given pressure. The results of the experiments are described and the
results compared with values taken from the literature.

4
J. S.

R.M.T.

BANKOVSKIY, Yu.A.; MIGULOVINA, Z.V.; TSIRULE, Ya.A.; IYEVIN'SH, A.F.
[Ievins, A.]

8-Chloro-8-mercaptoquinoline and its salts. Metod.poluch.khim.reak.i
prepar. no.4/5:79-85 '62. (MIRA 17:4)

1. Institut khimii AN Latvviyakoy SSR.

MICULKA, Bretislav, inz.

Refractometric dry matter in strawberries of world assortment.
Rost výroba li no. 3:287-302 M-165.

I: Cultivation Station, Velehrad. Submitted February 5, 1964.

MICULKS, Bretislav, inz.

Evaluation of strawberry varieties for canning. Prax. rosta 16
16 no.1:38-42 Ja 1965.

1. Plant cultivation station, Velhrad. Submitted August 5,
1964.

CZECHOSLOVAKIA / Weeds and Weed Control

N

Abs Jour: Ref Zhur-Biol., 1958, No 17, 77967

Author : Miculka, B.

Inst : ~~Not given~~

Title : Use of 2.4-D for Control of Weeds in Berry Patches.

Orig Pub: Ovoenar. a zelinar., 1956, 4, No 6, 166-168

Abstract: When spraying berry patches with herbicides, it is necessary to take into account the stage of development of the plants. With 2.4-D spraying of soil in a dose of 3 kg/ha of 100 m² through 6 weeks immediately before transplanting strawberries, 12 kg of weeds were counted; on the control plot, there were 555 kg. But, in addition on the

Card 1/2

HAUPTMANN, Erik, dr.; MICHLINIC, Rudolf, dr.

Comparative studies on the IGT, chlorpropamide and tolbutamide test. *Liječn. vjesn.* 86 no.4:427-432 Ap '64

1. Iz Internog odjela Bolnice "Dr. O. Novosel" , Medicinskog fakulteta u Zagrebu.

MICULESCU, R.

"Improvement of the technological process in manufacturing tires in the
Rosita Metallurgic Combine."

p. 351 (Studii Si Cercetari De Metalurgie) Vol. 2, no. 3, 1957
Bucharest, Rumania

SO: Monthly Index of East European Accessions (FEAI) LC. Vol. 7, no. 4,
April 1958

MICULESCU, R.; BERCU, S.; DRAGAN, I.

"Determination of the free expansion of copper and brass, laminated with noncalibrated cylinders. In French."

p. 107 (Revue De Metallurgie. Journal of Metallurgy) Vol. 1, 1956
Bucharest, Rumania

SO: Monthly Index of East European Accessions (EEAI) LC. Vol. 7, no. 4,
April 1958

MICU, Viorel

Putting the thematic plan of innovations into practice. Munca
sindic 6 no.7:38-40 J1 '62.

MIȚU, S.; VLAD, I.

"Degree of mechanization in model nurseries and machines used in various technical phases". p. 226, (REVISTA PADURILOR, Vol. 69, No. 5, May 1954, Bucuresi, Rumania)

SO: Monthly List of East European Accessions, (EMAL), LC, Vol. 4, No. 12, Dec. 1954, Uncl.

MICU, S. ; TUDOSIU, P.

"System of technical maintenance of tractors in forest protective stations." p. 37.
(REVISTA PADURILOR, Vol. 68, no. 4, April 1953, Bucuresti, Rumania)

SO: Monthly List of East European Accessions, L. C., Vol. 3, No. 4, April 1954, Uncl.

MICU, M.

Systematic development of labor productivity. Munca sindic 7 no.4:
12-14 Ap '63.

1. Presedinte al comitetului sindicatului uzinele "Infratirea",
Oradea.

MICU, M.

Compound nucleus influence on Coulomb excitation. Acta physica Pol
23 no.3:305-320 Mr '63.

1. Institute for Atomic Physics, Bucharest.

21306

P/045/61/020/002/004/006

B108/B209

Angular correlation ...

formula is the following: If θ_1, φ_1 determine the direction of the momentum \vec{k}_1 in the initial (center of mass) frame, the angles θ, φ determine the direction of the momentum \vec{k}_2 in the frame obtained by rotation and subsequent Lorentz transformation in the direction of the new z-axis. The Lorentz transformation was chosen so that in the final frame the nucleus B was at rest. If the masses of the nuclei A, B, and C are greater than the masses of the particles d_1 and d_2 , the Lorentz transformation is not necessary and θ is the angle between the momenta of the emitted particles d_1 and d_2 .

There are 2 non-Soviet-bloc references.

ASSOCIATION: Institute for Atomic Physics, Bucharest

SUBMITTED: August 5, 1960

Card 5/5

P/045/61/020/002/004/006
 B108/B209

Angular correlation ...

$$W = \sum_{\substack{\lambda_1, \lambda_2, \lambda_B \\ \lambda_C^k}} (-1)^{\lambda_B + \lambda_C + \lambda_1} |\langle \lambda_2 \lambda_C | S_2^{j_B} | b \rangle \langle \lambda_1 \lambda_B | S_1^{j_A} | a \rangle|^2 C_{\lambda_B - \lambda_B^0}^{j_B \lambda_B^k} C_{\lambda_1 - \lambda_C \lambda_C - \lambda_0}^{j_B \lambda_B^k} P_k(\cos \theta) \quad (10)$$

For any polarization detection the author introduces the efficiency matrix ϵ and the density matrix ρ of the final state (5), and from the formula $W = \text{Spur}(\rho \epsilon)$ he obtains

$$W = \sum_{\lambda_1, \lambda_2} B_{\lambda_1}^k(1) B_{\lambda_2}^k(2) D_{\lambda_1, \lambda_2}^k(\varphi \theta - \varphi) \quad (11)$$

where

$$B_{\lambda_1}^k(1) = \sum_{\lambda_1', \lambda_B^k} (-1)^{j_B - \lambda_B} \langle \lambda_1 \lambda_B | S_1^{j_A} | a \rangle \langle \lambda_1' \lambda_B^k | S_1^{j_A} | a \rangle^* C_{\lambda_B - \lambda_B^k}^{j_B \lambda_B^k} \langle \lambda_1' | \epsilon_1 | \lambda_1 \rangle \delta_{\lambda_1 - \lambda_B \lambda_1 - \lambda_B^k} \quad (12)$$

$$B_{\lambda_2}^k(2) = \sum_{\lambda_2', \lambda_C^k} (-1)^{\lambda_C - \lambda_2 - j_B} \langle \lambda_2 \lambda_C | S_2^{j_B} | b \rangle \langle \lambda_2' \lambda_C^k | S_2^{j_B} | b \rangle^* C_{\lambda_C - \lambda_2 \lambda_2 - \lambda_C^k}^{j_B \lambda_C^k} \langle \lambda_2' | \epsilon_2 | \lambda_2 \rangle \quad (13)$$

with ϵ_1 (ϵ_2) referring to the polarization detection of the first (second) radiation. The meaning of the angles θ , φ used in the angular correlation

Card 4/5

21306

Angular correlation ...

P/045/61/020/002/004/006
B108/B209

in the form

$$|\vec{p} \lambda_1 \lambda_2\rangle = \sum_{JM} |JM \lambda_1 \lambda_2\rangle \langle JM \lambda_1 \lambda_2 | \vec{p} \lambda_1 \lambda_2 \rangle \quad (1) \quad (1)$$

where \vec{p} denotes the relative momentum. By using Eq. (1), one obtains

$$\langle \vec{k}_1 \lambda_1 \lambda_B | S_1 | a; j_A m_A \rangle = \langle \vec{k}_1 \lambda_1 \lambda_B | j_A m_A \lambda_1 \lambda_B \rangle \langle \lambda_1 \lambda_B | S_1^j | a \rangle \quad (7) \quad (7)$$

where $S_1^{j_A}$ represents the submatrix of S_1 belonging to a given angular momentum j_A . If parity is conserved in the decay $A \rightarrow B + d_1$, one finds

$$\langle \lambda_1 \lambda_B | S_1^j | a \rangle = \frac{\eta_1 \eta_B}{\eta_A} \langle -\lambda_1 - \lambda_B | S_1^j | a \rangle \quad (8) \quad (8)$$

the quantities η being the parity factors. If the initial nucleus is unpolarized,

$$W = \sum_{\lambda_B \lambda_C} |D_{\lambda_B \lambda_C}^{j_B}(\varphi_0 - \varphi)|^2 |\langle \lambda_2 \lambda_C | S_2^{j_B} | b \rangle \langle \lambda_1 \lambda_B | S_1^j | a \rangle|^2 \quad (9)$$

which, for detectors insensitive to polarization, becomes

Card 3/5

X

21306

P/045/61/020/002/004/006
B108/B209

Angular correlation ...

and, since the S matrix is Lorentz-invariant, the final state becomes

$$\psi = \sum_{\substack{\vec{k}_1, \lambda_1 \\ \vec{k}_2, \lambda_2}} (-1)^{j_B - \lambda_B} |\vec{k}_2, \lambda_2; \vec{k}_1, \lambda_1\rangle \langle \vec{k}_2, \lambda_2 | S_2 | b; j_B - \lambda_B \rangle \langle \vec{k}_1, \lambda_1 | S_1 | a; j_A m_A \rangle \quad (5)$$

The ket vector $|\vec{k}_2, \lambda_2; \vec{k}_1, \lambda_1\rangle$ describes the following state: The free particle d_1 with momentum \vec{k}_1 and helicity λ_1 , the free particle d_2 with momentum \vec{k}_2 and helicity λ_2 , and the nucleus C with momentum $-\vec{k}_2$ and helicity λ_C . The first momentum is referred to a frame, in which the nucleus A is at rest while both the second and the third momentum are referred to a frame, in which the nucleus B is at rest. The probability to find this state in the final state is

$$W = \left| \sum_{\lambda_B} \langle \vec{k}_2, \lambda_2 | S_2 | b; j_B - \lambda_B \rangle \langle \vec{k}_1, \lambda_1 | S_1 | a; j_A m_A \rangle \right|^2 \quad (6)$$

The two wave functions used by Jacob and Wick in their paper are interrelated by the matrix

(2)

21306

P/045/61/020/002/004/006
B100/B209

24.4500

AUTHOR:

Micu, M.

TITLE:

Angular correlation theory with the Jacob-Wick method

PERIODICAL:

Acta Physica Polonica, v. 20, no. 2, 1961, 157-159

TEXT: In the present paper, some simplifications in the angular correlation theory are made following the procedure applied by M. Jacob and C. C. Wick (Ref. 2: Ann. Phys., 7, 404 (1959)) in the collision theory. The state of the initial nucleus A is $|a; j_A m_A\rangle$. a stands for all the quantum numbers of this nucleus except spin (j_A) and its projection on the z-axis (m_A). If $S_1(S_2)$ is the S matrix which effects the first (second) transition in the cascade decay $A \rightarrow B + d_1$, $B \rightarrow C + d_2$, the final state is $\psi = S_2 S_1 |a; j_A m_A\rangle$ (3). In the momentum-helicity representation for the relative motion this reads

$$\psi = \sum_{\lambda_1 \lambda_2} \delta_{\lambda_1 \lambda_2} |\vec{k}_1 \lambda_1 \lambda_2\rangle \langle \vec{k}_1 \lambda_1 \lambda_2 | S_1 | a; j_A m_A \rangle \quad (4)$$

Card 1/5

X

MICU, M.

Stability of a plasma contracted by an electromagnetic field.
Studii cerc fiz ll no.4:827-836 '60. (EEAI 10:8)

1. Institutul de fizica atomica, Bucuresti.
(Fluids) (Electromagnetic fields) (Mathieu functions)
(Plasma(Ionized gases))

MICU, M.

On the transition intensities in disintegration β . Studii cerc fiz 11
no.2:437-439 '60. (EEAI 10:1)
(Beta rays) (Electric moment) (Radioactivity)

MICU, M.

The Coulomb excitation of particles, due to the quadrupole moment
of the projectile. Studii cer.fiz. 10 no.4:659-671 '59.

(EEAI 9:5)

(Particles) (Nuclear moments) (Electric moment)

MICU, M.

17

Reduced widths for nuclear emission, according to Nilsson's model. M. Micu and A. Sandulescu. Acad. rep. populare Romine, Inst. fis. atomica si inst. fis. Studi cercetari no. 10, 651-7 (1959) (English summary).—The reduced widths for a single nucleon emission were calcd. and it was found that the nuclear wave function is separately antisym. with respect to neutrons and protons, the individual

5

..
j
|||
dw
wave functions being those given by N. (CA 50, 7005h). Theoretical results were compared with exptl. data for a few light nuclei — N¹², N¹⁴, F¹⁷, B¹², C¹², O¹⁷.

M. Ben Eliezer

MICU, M.

19

✓ Reduced widths for deuteron emission calculated by means of the nuclear shell model. M. Micu and A. Săndulescu. *Acad. rep. populare Romîne, Inst. fiz. atomica si Inst. fiz., Studiul cercetării fiz.* 10, 257-76 (1969); cf. Lane, *C.A.* 48, 6289a.—A study is presented on reduced widths in reactions with deuterons produced by complex nuclei, calcd. by means of the nuclear shell model. Anal. expressions of reduced widths have been obtained for the case of the complex system of n equiv. nucleons, on the outside of a closed shell, as well as in the particular case when there is only one single nucleon. Each expression is obtained in the case of the 2 extreme couplings LS and JJ of the nuclear shell model. Results are compared with exptl. results in the case of the reactions $He^4 \rightarrow T^3 + D^2$ and $Li^6 \rightarrow He^4 + D^2$. M. Laphire

5

CFK

CMR

MICU,

M

RUMANIA/Electronics - Electrical Discharges of Gases and Gas Discharge Apparatus. H

Abs Jour : Ref Zhur Fizika, No 1, 1960, 1528

Author : Ciulli, S., Micu, M.

Inst : -

Title : Plasma Oscillations in a Static External Magnetic Field.

Orig Pub : Studii si cercetari fiz - Acad. RPR, 1958, 9, No 4, 489-496

Abstract : By using the Boltzmann equation, the author finds the distribution function of a beam of electrons in a plasma with cylindrical symmetry, located in an axial magnetic field. The perturbations of the stationary distributions are investigated by calculating the deviations of the first-order moments from their Maxwellian value. The values of the perturbations of the magnetic field and of the distribution

Card 1/2

- 79 -

PA - 2048

The Statistical Method for the Study of the Behavior of a
Totality of Charged Particles under the Effect of an Own
Magnetic Field (Russian).

equations is specialized for this case. By separation of the
variable the authors endeavor to find a solution which
satisfies the condition $e_1 f_1 + e_2 f_2$. On the occasion of transi-
tion to a system of reference in which electrons and ions
have the same absolute velocity the equations for f_1 and f_2
become identical. In the case of equal initial conditions
($f_1(0) = f_2(0)$) both functions thus become identical. The
necessary operations of computation are discussed in short.
An equation for $f(r)$ is given and several times transformed,
its solution is written down in implicit form and is also
transformed. The results obtained here are correct only if
the density of the ions (f) becomes very low near the interior
walls of the tube (where temperature must not exceed a certain
limit). In conclusion the definite form of the function of
statistical distribution is given.

ASSOCIATION: Institute for Nuclear Physics, Bucarest, Roumania
PRESENTED BY:
SUBMITTED:
AVAILABLE: Library of Congress
Date: 3/7

PA - 2048

The Statistical Method for the Study of the Behavior of a
Totality of Charged Particles under the Effect of an Own
Magnetic Field (Russian).

$B_{\mu\nu} = \partial A_\nu / \partial x^\mu - \partial A_\mu / \partial x^\nu$, $\square A^\mu = -j^\mu$. Here ξ^μ denotes
the four-vector of velocity, F_1 - the distribution function
of the ions, F_2 - the distribution function of the electrons,
 j^μ - the four-vector of the flux of charged particles. The
present work solves the integrodifferential equations for
the steady isothermal distribution with axial symmetry, all
computations being carried out in nonrelativistic approxima-
tion.

The field $B^{\mu\nu}$ can be expressed by the components of the
fourdimensional vector A^μ . The solution of the wave equa-
tion for A^μ is given, and also the herefrom resulting com-
ponents of B_{12} . In the case of cylinder symmetry the functions

F_1 and F_2 do not depend on φ and the components B_{12} , B_{23} ,
 B_{24} , B_{34} vanish in this case. The expressions for B^{13} , and
 B^{14} are explicitly given. The first of the above mentioned

Card 2/3

MICU, M.

PA - 2048

AUTHOR: ČJULILI, S., MIKU, M.
TITLE: The Statistical Method for the Study of the Behavior of a
 Totality of Charged Particles under the Effect of an Own
 Magnetic Field (Russian).
PERIODICAL: Atomnaia Energiia, 1957, Vol 2, Nr 1, pp 5-9 (U.S.S.R.)
 Received: 3 / 1957 Reviewed: 3 / 1957

ABSTRACT: The present work shows the existence of steady solutions for
 the function of the statistical distribution of particles in
 a gas discharge. On this occasion the discharge is held back
 round the symmetry axis only under the influence of its own
 magnetic field. The gas is assumed to be sufficiently warm
 and to be fully ionized. Therefore the influence exercised
 by the diffusion of neutral atoms into the plasma is not
 taken into account. The equations of relativistic statistics
 for the totality of the charged particles which are under
 the influence of their own field are written down in the form
 given by S. CICEJKA. If the rationalized system of GAUSS units
 and of the antisymmetric tensor

$B^{\mu\nu}$ for the electromagnetic field is used, the following
 equations are obtained: $\frac{\partial}{\partial x^\mu} (\xi^\mu F_{1,2}) + \frac{e_{1,2}}{m_{1,2}} B^{\mu\nu} \left\{ \frac{\partial F_{1,2}}{\partial \xi^\mu} \right\}_\nu = 0$

Card 1/3

MICU, M. ; CIULLI, S.

Diffusive and dispersive propagation of gravitational waves in a vacuum; also, remarks by c. Aretin, p. 507. Academia Republicii Populare Romine. Institutul de Fizica. STUDII SI CERCETARI DE FIZICA. Bucuresti. Vol. 6, no. 3, July/Sept. 1955.

So. East European Accessions List Vol. 5, No. 9 September, 1956

MICU, L., ing.

Some considerations connected with the use of the Wheatstone
bridge. Metrologia apl 10 no.9:411-417 S '63.

MIGU, L., ing.

Problems on the magnetoelectric galvanometers and their selection
for various mountings. Metrologia apl 10 no.4:162-168 Ap '63.

ROSU, E.; MICU, L.; BAN, I.

Preparation of injection solutions of novocaine with pyramidon
and sodium amytal. Apt. delo 10 no. 1:94 Ja-F '61. (MIRA 14:2)
(NOVOCAINE)

EXCERPTA MEDICA Sec 10 Vol 12/12 Obstetrics DEC 59

2105. A CASE OF CONGENITAL TUBERCULOSIS - Velezületett tuberkulózis esete - Deutsch M. and Micu J. Szatmár Egyesített Kórház Tüdőosztál. Közl. - ORV. SZLE 1958, 476 (538-539)

Mention is made of the low frequency of these cases and a description is given of a 19-day-old infant admitted with a 3-day history of influenza and bronchopneumonia. Birth weight was 2,650 g. BCG vaccination was given on the 8th day. Body weight was only 2,500 g. on admission; there was insomnia, diarrhoea and pre-dystrophy. Perioral cyanosis was also seen. No X-ray film was made. Fluoroscopic examination showed densification at the hilus. Enlarged lymph nodes were palpable in the left axillary region. The CSF was Pandy-positive. The tuberculin test (1:10,000) was negative. Blood counts: erythrocytes 3.9 million, leucocytes 20,600, Hb 75%. Streptomycin and isoniazid were given. On the 26th day of illness the subject died after further loss of weight. Autopsy revealed miliary tb. No tb was elicited in the home environment. Eight days after the child's admission the mother developed fever, cough and abdominal pain, and likewise showed miliary tb. It is suggested that infection of the infant had probably been intraplacental, being effected during the last few months of pregnancy.

Schaich - Luisenheim (XV, 7, 10)

ZAVATE, Olga; CONSTANTINESCU, N.; DOGARU, Maria; MORARU, Aneta; FRANCHE,
Maria; MICU, I., BALTEV, Aradna

Research on recurrences of exanthematous typhus. IV.
State of rickettsemia in sporadic typhus primo-infection
and in recurrences. Stud. cercet. infirmicobiol. 15
no.3:255-267 '64.

FRANCHE, Maria; MICU, I.; BALTIEV, Ariadna; DUMITRIU, St.; FELLER, H.;
APOSTOL, A.; BRAUNER, E.; CONSTANTINESCU, N.; ZAVATE, Olga;
DOGARU, Maria; NICA, V.

Research on recurrences of exanthematous typhus. II. Comparative
clinical aspects of typhus recurrences and primary infections.
Stud. cercet. inframicrobiol. 15: no. 3: 211-224 '64.

Mico, I.

ROMANIA

MICU, I.; OGANA, C.; MANTA, I.; IOAN, Elena; CUCIOREANU, Georgeta;
MIHUL, Valentina; VINTU, C.; GRADINARU, Lilliana; GRADINARU, I.;
IOSEFSOHN, Judith; MINASCURTA, S.; MOSANU, P.; COTAE, Gh.

Clinic of Contagious Diseases Iasi, Iasi Regional Sanepid.
(Clinica de boli contagioase Iasi, Sanepidul regional Iasi.)
- (for all)

Bucharest, Viata Medicala, No 7, 1 Apr 63, pp 457-460.

"Epidemic of Ornithosis in a Rural Locality."

(13)

BALDOVIN-AGAPI, Corălia, dr.; FRANCHE, Maria, prof.; BELOIU, Irina, dr.;
MICU, I., dr.; OVANESCU, A., dr.; ANDRONOVICI, G., dr.; BRAUNER, E.,
dr.; RADULESCU, A., dr.; DIMITRIU, St., dr.; DIMITRIU, A., dr.;
RUGINA, N., dr.; BLINDU, P., dr.

Receptivity to scarlet fever assessed by Dick's reaction with
fractional doses of purified toxin. Microbiologia (Bucur) 6
no. 1:69-76 Ja-F '62.

1. Institutul "Dr. I. Cantacuzino" si Spitalul "Izolarea" din
Iasi.

NICOLAU, S.St, academician; ZAVATE, O.; CONSTANTINESCU, N.; MICU, I.;
BIRZU, N.; RUSU, Florica; OVANESCU, Al.

Research on viral infectious hepatitis (V.I.H.) transmitted by
parenteral route. Stud. cercet. inframicrobiol. 12 no.4:421-435
'61.

(HEPATITIS, INFECTIOUS transmission)
(INJECTIONS complications) (HOSPITALS)

CONSTANTINESCU, N.; CELEANU, M.; MICU, I.; BIRZU, N.; ZAVATE, O.;
MORARU, A.

Strains of the modified rabic virus, isolated from human meningo-
encephalomyelitis with a course atypical of rabies. Studii cerc
inframicrobiol Special issue-supplement to 12:167-174 '61.

1. Institutul de inframicrobiologie al Academiei R.P.R. si Institutul
de igiena, Iasi.

(HYDROPHOBIA) (ENCEPHALOMYELITIS)

NICOLAU, St. S., acad.; CONSTANTINESCU, N.; BIRZU, N.; ZAVATE, O.; MICU, I.
TEODOROVICI, Gr.

Evolution of human rabies comparatively studied in both treated and untreated patients with antirabic vaccine. Consecutive therapeutic directives. Studii cerc inframicrobiol Special issue-supplement to 12:133-143 '61.

1. Institutul de inframicrobiologie al Academiei R.P.R. si Institutul de igiena, Iasi. 2. Membru al Comitetului de redactie si redactor responsabil, "Studii si cercetari de inframicrobiologie"(for Nicolau).

(HYDROPHOBIA)

MICU, I., dr.; OVANESCU, Al., dr.; CUCIUREANU, Georgeta, dr.; BEJENARIU, C.,
dr.

Pyocyanic and staphylococcal septicopyohemia with pleural focus.
Med. intern., Bucur 12 no.10:1531-1536 0 '60.

1. Lucrare efectuata in Clinica de boli contagioase Iasi (director:
prof. Maria Franche).
(SEPTICEMIA etiol.) (STAPHYLOCOCCAL INFECTIONS case reports)
(PLEURA dis.)

RUMANIA/General Problems of Pathology. Allergy

U-2

Abs Jour : Ref Zhur - Biol., No 14, 1958, No 65911

Author : Constantinescu N., Micu, I., Munteanu G., Gheorghiu M.,
Birzu, N., Blindu P., Vacs L.

Inst : Rumanian Academy

Title : Preliminary Data on the Allergic Intradermal Test in Epi-
demic Hepatitis

Orig Pub : Comun. Acad. RPR, 1957, 7, No 2, 273-277

Abstract : By means of the Vladimirova intradermal test (Clinic. med., 1951, 7), the authors detected the presence of an auto-allergen (AA) in a filtrate of gastric juice from a patient with epidemic hepatitis throughout the course of the disease. The AA reaction was negative in well persons and in patients with gastric and hepatic diseases.

Card : 1/1

Country : M
Category : CULTIVATED PLANTS, FRUITS

Ann. Jour. : SÉPTEMBRE-BIOL, 21, 1968, NO-96187

Author :
Instit. :
Title :

Orig. Pub. :

Abstract : conditions was 33.9% in the Italian Riesling, 41.2%
in the Cal'dan from Odobesti, and 81.9% in the
Muscat Hamburg.---Ye.T. Zhukovskaya

Card: 3/3

Country : M
 Category : CULTIVATED PLANTS, FRUITS
 Abs. Jour. : REF ZHUR-BIOL., 21, 1958, NO-96167
 Author :
 Distribut. :
 Title :
 Orig. Pub. :
 Abstract : half closed normally hermaphroditic varieties in the field is explained by the lack of access to the pollen grains from one variety to another (the shattering of blossoms in this case averaged 35.8%). Amongst varieties with normally hermaphroditic open flowers the shedding was 28.5% higher than in varieties with the half-open blossoms which were protected from the injurious effects of external factors (lowered temperature, strong rain). The degree of flower shedding under field

Card: 2/3

MTCU 1.

Country : Rumania
 Category : CULTIVATED PLANTS. FRUITS. Berries.

M

Abstr. Jour. : REF ZHUR-BIOL., 21, 1958, NO-96167

Author : Nicu, Ilie
 Institut. : N. Balcescu Agronomic Inst.; Agric. Inst. Bucuresti.
 Title : Observation on Blossom Dropping in Certain Grape Varieties Distributed to the RPR in Hothouses and under the Field Conditions of Bucuresti.
 Orig. Pub. : Gradina, via si livada, 1957, 6, No.5, 37-40

Abstract : It has been established by experiments made by the N. Balcescu Agronomic Institute and the Agricultural Institute in Bucuresti that when foreign pollen is absent, needed for cross-pollination in grape varieties having functionally female flowers, there is total flower dropping (100% in the hothouses and on the fields) in the case where the flowers are half-open, and partial (85% in the hothouses and 76.4% under field conditions) when the flowers are open. Flower shattering in the

Card: 1/3

MICU, Gh.

Organization of construction material transportation. Probleme
econ 17 no.1:102-117 Ja '64.

MICU, D.; ZAMFIRESCU-GHEORGHIU, Marcela; MIHAILESCU, Eugenia; MAXIMILIAN, Stefania; VLADESCU, C.

The value of investigating some leukocyte phosphatases in various malignant blood diseases. Stud. cercet. med. intern. 6 no.3: 279-285 '65.

MICH, B.

Blood and bone marrow cultures of a patient with acute leukemia.
Stud. cancer. med. intern. 5 no. 6: 537-6 (1974).

MICU, D.; SAMPHIRESCU-GHEORGHIU, Marcela; MAXIMILIAN, Stefania; MIHAILESCU,
Eugenia; DANCESCU, Ileana

Cyto-enzymatic anomalies in some acute leukoses. Stud. cercet.
med. intern. 5 no.4:419-421 '64.

MICU, D.

The use of lymph-node punch biopsy for the diagnosis of
benign and malignant adenomegaly. Stud. cercet. med. intern.
5 no.2:131-144 '64

MICU, D.; ZAMFIRESCU-GHEORGHIU, Marcela; MAXIMILIAN, Stefania; MIHAILESCU, Eugenia; LAZAROVICI, Miriam.

Contributions to the study of lipid disorders appearing in the lymph nodes in some systematic diseases. Stud. cercet. med. intern. 5 no.1:55-57 '64

1964.

MICO, D., MD; MURAI, V., Lecturer; MIHAILANOU, Eugenia, MD;
MIXIMILIAN, Stania, MD; MIROU, G., MD; MURIC, G., MD;
SALCA, Isadora, Technical Assistant.

Institute of Internal Medicine of the Academy of the Romanian
People's Republic and the Ministry of Health and Public
Welfare (Institutul de medicina interna al Academiei R.P.R.
si M.S.P.S.) - (for all); Director: Academician H. M. Ivan.

Bucharest, Medicina Interna, No 12, Dec 65, pp 1457-1464

"The Value of Certain New Methods in the Study of Immunoglobulins
in Chronic Hepatitis."

MICU, D.; JIPA, Gh.; MIRON, C.; RUNCAN, V.

Contribution to the knowledge of the circulating lymphocytes
in chronic hepatitis. (Fluorescence microscopic study). Stud.
cercet. med. intern. 4 no.6:795-801 '63

*

JIPA, G.; MICU, D.

Study of the bone marrow and blood cells in leucoses with the aid of the fluorescent microscope. Stud. cercet. med. intern. 4 no.4:497-502 '63.

(LEUKEMIA) (LEUKEMIA, MYELOCYTIC)
(LEUKEMIA, LYMPHOCYTIC)
(BONE MARROW EXAMINATION)
(BLOOD CELLS) (MICROSCOPY, FLUORESCENCE)
(DNA) (RNA)

MICU, D., dr.; MAXIMILIAN, StoŃania, dr.; MIHAILESCU, Eugenia, dr.

The value of the blood catalase index in the orientation of radiotherapy of the leukemias. Med. intern. 15 no.1:101-106 Ja '63.

1. Lucrare efectuata la Institutul de medicina interna al Academiei R.F.R. si M.S.P.S. Bucuresti (director: acad. N. Gh. Lupu).
(LEUKEMIA) (RADIOTHERAPY) (CATALASE)
(BLOOD CHEMICAL ANALYSIS)