

The Soviet Far East; Economic and Geographic Features (Cont.) 245

Ch. I. Physical and Geographic Features and Natural Resources	9
Topography	9
Minerals	17
Lakes and rivers	22
Climate	28
Permafrost	32
Soils	33
Flora	36
Fauna	45

Card 3/5

The Soviet Far East; Economic and Geographic Features (Cont.)	245
Ch. II. Stages of Colonization	48
Ch. III. Population	67
Ch. IV. Economy	77
Industries	78
Agriculture and animal husbandry	102
Transportation and interregional communications	121
Ch. V. Features of Individual Krays and Oblasts	134
Khabarovskiy Kray	134
Primorskiy Kray	154

Card 4/5

The Soviet Far East; Economic and Geographic Features (Cont.)	245
Amurskaya Oblast	175
Sakhalinskaya Oblast	192
Kamchatskaya Oblast	211
Magadanskaya Oblast	221
Bibliography	235
Appendix	237

AVAILABLE: Library of Congress

GC/bmd
May 28, 1958

Card 5/5

KLYUCHEROV, A.P.; KONDRAT'YEV, S.N.; Prinimali uchastiye: GUSAROV, F.V.;
UDOVENKO, V.G.; PETROV, G.A.; BURKSER, V.Ye.; SHMONIN, I.A.;
KUDRIN, Ye.A.; GALAKHMETOV, S.H.; ZIMINA, L.P.; SHISHARIN, B.N.;
KONDYURINA, R.V.; BURMISTROV, K.A.; SHIRNIN, I.A.; SIMONENKO, F.N.;
GORSHILOV, Yu.V.; KOLPAKOV, B.V.; GUSAROV, A.K.; BOLOTOV, P.G.

Heat insulation of open-hearth furnace crowns. Metallurg 5 no.11;
(MIRA 13:10)
14-17 N '60.

1. Nizhe-Tagil'skiy metallurgicheskiy kombinat.
(Open-hearth furnaces--Design and construction)
(Insulation (Heat))

KONDRAT'YEV, S.N.; KLYUCHEROV, A.P.; UDOVENKO, V.G.; SHIRNIN, I.A.;
VYDRINA, Zh.A.

Rapid methods of repair and the fritting of new hearth bottoms.
Metallurg 6 no.9+10+13 S '61. (MIRA 14:9)

1. Nizhne-Tagil'skiy metallurgicheskiy kombinat.
(Open-hearth furnaces--Maintenance and repair)

BARANOV, V.M.; DONSKOY, S.A.; TORSHILOV, Yu.V.; TRET'YAKOV, M.A.; UDovenko,
V.G.; FREYDENZON, Ye.Z.

Blowing of cast iron in high-capacity converters. Metallurg 10 no.9:
15-18 S '65. (MIRA 18:9)

1. Nizhne-Tagil'skiy metallurgicheskiy kombinat.

ALAMPIYEV, P.M.; OMAROVSKIY, A.G.; UDOVENKO, V.4

New features of the literature on the economic regions of the
U.S.S.R. Izv. AN SSSR. Ser. geog. no.4:147-151 Jl-Ag '63.

(Bibliography--Geography, Economic) (MTRA 16:8)

DUBROV, N.F.; KITAYEV, B.I.; KOKAREV, N.I.; UDOVENKO, V.G.; KONDRAT'YEV, S.N.;
ZATULOVSKAYA, Ye.Z.; KLYUCHEROV, A.P.

Review of the book by N.A.Vecher "Highly efficient operation of
open-hearth furnaces." Stal' 24 no.7:613-614 Jl '64.

(MIRA 18:1)

1. Ural'skiy nauchno-issledovatel'skiy institut chernykh metallov,
Ural'skiy politekhnicheskiy institut i Nizhne-Tagil'skiy metallurgi-
cheskiy kombinat.

ARNAUTOV, V.T.; BARANOV, V.M.; DONSKOY, S.A.; PASTUKHOV, A.I.; SMIRNOV, I.A.; TORSHILOV, Yu.V.; TRET'YAKOV, M.A.; UDovenko, V.G.; FREYDENZON, Ye.Z.; SHCHEKALEV, Yu.S.; Prinimali uchastiye: MAKAYEV, S.V.; KOMPAKIETS, G.M.; NAGOVITSYN, D.F.; NOVOLODSKIY, P.I.; VARSHAVSKIY, V.L.; KOROGCDSKIY, V.G.; KLIBANOV, Ye.L.: MEDVEDEVSKIKH, Yu.; TALANTSEVA, T.I.; DUBROV, N.F.; DZEMYAN, S.K.; TOPYCHKANOV, B.I.; CHARUSHNIKOV, O.A.; KHARITONOV, Yu.A.

Developing and mastering the technology of converting vanadium cast iron in oxygen-blown converters with a 100 ton (Mg) capacity.
Stal' 25 no.6:50/-508 Je '65. (MIRA 18:6)

1. Nizhne-Tagil'skiy metallurgicheskiy kombinat (for Makayev, Kompiets, Nagovitsyn, Novolodskiy, Varshavskiy, Korogodskiy, Klivanov, Medvedevskikh, Talantseva). 2. Ural'skiy nauchno-issledovatel'skiy institut chenykh metallov (for Dubrov, Dzemyan, Topychkanov, Charushnikov, Kharitonov).

FREYDENZON, Yu.Z.; UDENKO, V.G.; TVERDILOV, Yu.V.; ROMANOVSKIY, G.M.;
TRET'YAKOV, M.A.; BARANOV, V.M.; RAGOVITSIN, L.F.; MONGOV, S.A.;
PASTUKHOV, A.I.

Mastering the operation of the oxygen-blown converter plant
of the Nizhniy Tagil metallurgical combine. Stat' 25 no. 6:
534-537 Je '65.
(MIRA (Ed.))

I. Nizhne-Tagil'skiy metallurgicheskiy kombinat i Ural'skiy
nauchno-issledovatel'skiy institut chernykh metallov.


VARLAMOV, V.S., kand.tekhn.nauk; IL'INA, A.I.; KUDRYASHOV, A.I., inzh.;
UDOVENKO, V.S., inzh.; KOGAN, G.A., inzh.

Continuous oxidation of paraffins under industrial con-
ditions. Masl.-zhir.prom. 25 no.10:39-41 '59.

(MIRA 13:2)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut zhirov
(for Varlamov, Il'ina). 2. Shebekinskiy kombinat sintetiche-
skikh zhirnykh kislot i zhirnykh spirtov (for Kudryashov,
Udovenko, Kogan).

(Shebekino--Paraffins)



2

Adiabatic expansion of saturated vapors and fog formation. N. A. Korosovskii
AND V. V. Usovskii. *J. Gen. Chem. (U. S. S. R.)* 1, 1245-8(1931); cf. C. I., 36, 307N.
The validity of the rule $C_v'/(\bar{n}-10) > 0$, for sp. heats of satd. vapors, where C_v' is sp. heat
and \bar{n} no. of atoms in the mol., was established in a general way by allowing a satd. vapor
to expand adiabatically from a bulb. The appearance of a fog is an indication of a neg.
to expand adiabatically from a bulb. The appearance of a fog is an indication of a neg.
 C_v' , but when fog does not appear, the reverse is true. Fog formation was aided by
ultra violet rays. Fog was observed in the following compds. in which $n < 10$: CS,
CHCl₃, CCl₄, MeOH, MeCHO, nitromethane and EtOH. There was no fog formation
in the following compds. in which $n > 10$: Me₂CO, EtCOCl, CH₃OEt, PrOH, C₆H₆,
MeCOEt, AcOEt, Et₂O, Et₂CO, C₆H₅ and Pr₂O. S. L. MADDEN


ASIN-SEA METALLURGICAL LITERATURE CLASSIFICATION


Specific heat of liquids. II. N. A. Kolesovskii and V. V. Udrovenco, *J. Gen. Chem. (U. S. S. R.)* 4, 1027-33 (1934); cf. *C. A.* 30, 3649-. C_p, in cal., per g. mol. at 15-30°, and temp. coeff. of entropy (DS/DT), were detd. for a number of liquids. The following figures give benzene 31.39, 0.1090, C_p (DS/DT), and temp., resp. Benzene 47.84, 14.70; C₆H₅Br 42.73, 0.1411, 29.55; cumene 47.84, 0.1574, 28.87; limonene 59.82, 0.2032, 20.20; sabellene 0.0.32, 0.2031, 23.84; CCl₄ 30.21, 0.1047, 15.11; C₆H₅Br 40.03, 0.1354, 20.19; 34.77, 0.1149, 29.46; C₆H₅CHO 40.03, 0.1354, 20.19; methyl ethyl ketone 59.46, 0.1205, 23.81; heptanediol 59.05, 0.1390, 21.14; 1,3-dimethyl-1,3-dihydro-2H-1,3-dioxole 59.19, 0.1390, 21.14; formic acid 33.37, 0.1200, 21.82; ethyl butyrate 59.19, 0.1709, 24.06; butyl acetate 57.8; 0.1978, 19.30; benzyl acetate 50.78, 0.2042, 19.51; ethyl benzoate 57.84, 0.1980, 20.00; diethyl carbonate 50.28, 0.1710, 21.50; diethyl succinate 80.77, 0.2760, 19.42; ethyl acetoacetate 59.19, 0.2009, 24.39; m-nitrotoluene 48.28, 0.1600, 29.19; o-toluidine 50.09, 0.1636, 20.23; m-toluidine 51.84, 0.1712, 29.52; dimethylaniline 51.29, 0.1696, 20.15; diethyl aniline 55.58, 0.2171, 28.83; quinoline 46.13, 0.1525, 20.2. A table shows comparison of these results with those of other investigators. S. L. Mado.


S. L. Mado.

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7"









1. USANOVICH, M., STMAROKOVA, T., UDOVENKO, V.


2. USSR (600)


"Electrical Conductivity, Viscosity, and Surface Tension of the System $H_2SO_4-HClO_4$,"
Zhur. Obshch., 9, No. 21, 1939. Lab. of Physical Chem., Central Asiatic State
Univ. Received 9 June 1939.


9. [REDACTED] Report U-1626, 11 Jan 1952.











Mol. wt. determinations as a means of physico-chemical analysis.
V. V. Udoenko (*Acta Physicochim. U.R.S.S.*, 1942, 18, 336-347).—
The variation of mol. wt. with composition is determined by the
cryoscopic method and plotted in the case of MeOH-EtOH, and
PhMe-PhEt mixtures in C_6H_6 , to represent mixtures of similar
mols. where there is no chemical interaction, and for piperidine-
PhNCS and NHMePh-PhNCS mixtures, where there is interaction.
The mol. wt.-composition curve is a straight line where there is
no interaction, but the curve shows a max. coinciding with the com-
position of the compound where there is interaction. A. J. M.

(a)

The maximum of viscosity of irrational systems. V. V. Ulyanova (Phys. chem. Lab., Middle Asiatic State Univ.), Akad. Nauk S.S.R., Odz. Tekh. Nauk, Inst. Mashinostroyeniya, Sovetskoe po Vysokotekhnicheskim Kolloid. Rastvorov (Conf. on Viscosity of Liquids and Colloid. Solns.) 2, 89-93 (1944).—Comparison with melting diagrams has shown that in "irrational" binary systems the max. of the viscosity curves in terms of compn. does not occur at the compn. corresponding to that of the compn. formed. The usual interpretation in terms of dissoen. of the compn. is subject to caution. A diagram of a system A-B involving a compn. C can be constructed from the part diagram A-C, with n max., and C-B. The displacement of the max. in the resulting A-B curve is due to interaction of the compn. C and the component A and not to dissoen. This is further borne out by the observation that the max. does not tend to draw closer to the stoichiometric compn. as the temp. approaches the m.p., which would have to be the case if the displacement were due to dissoen. in liquid phase. The shift sometimes is found to persist below the m.p. in under-cooled liquid. In the system $\text{NaCl}-\text{CH}_3\text{COOH}$, the max. of viscosity lies at about 20% NaCl, when the compn. has the compn. 1:2 (m.p. 18.7°). In $\text{HClO}_4-\text{CH}_3\text{COOH}$, the max. lies at 67 mol. % CH_3COOH , and its position is unaffected by temp., whereas the compn. (m.p. 41°) is equimol. The same is true for the system phenol-aniline; the compn. (m.p. 30.5°) is equimol., but the max. of viscosity is at 66.07 mol. % phenol, independently of

temp. A further example of a shifted max. is $\text{HClO}_4-\text{H}_2\text{O}$. Comprn. and stability of binary compns. were further investigated by cryoscopic measurements on benzene solns. of various concns. of both the pure components and their mixts., and the results represented graphically. The curves of cryoscopic mol. wt. for the system chloro-EtOH show the compn. to be stable at all dilns. and dissoen. to be absent. The max. on the curve corresponding to infinite diln. is singular (discontinuous); on the other hand, with increasing concn. of the benzene soln., the max. of mol. wt. shifts gradually towards the EtOH; this evidences interaction between that component and the chloro ethylate. With rising temp., the max. moves towards the compn. In the 2 systems phenol-aniline and phenol-pyridine, the compn. is dissoed. at high dilns.; at a concn. of about 1 mole/1000 g. benzene, dissoen. is practically zero. At higher concns., the cryoscopic mol. wt. shows a max., shifted towards phenol; this again indicates interaction between the phenol and the equimol. compn. Some cases are also known where the compn. exists in the liquid phase only and therefore is not expressed in the melting diagram but only in the viscosity curve (example the system $(\text{C}_6\text{H}_5)_2\text{CH}-\text{SbBr}_3$). N. Thom

ASR-SLA METALLURGICAL LITERATURE CLASSIFICATION

SECTION 1: GENERAL SUBJECTS

SECTION 2: INDUSTRIAL METALS

SECTION 3: ALLOYS AND COMPOUNDS

SECTION 4: PROCESSING AND PROPERTIES

SECTION 5: EQUIPMENT AND APPARATUS

SECTION 6: RESEARCH AND DEVELOPMENT


SECTION 7: STANDARDS AND TEST METHODS

SECTION 8: PRACTICAL APPLICATIONS

SECTION 9: SPECIAL TOPICS

SECTION 10: INDEXES AND BIBLIOGRAPHIES

SECTION 11: OTHER



COMMON ELEMENTS		PROCESSES AND PROPERTIES INDEX																3RD AND 4TH ORDERS																	
		1ST AND 2ND ORDERS								3RD AND 4TH ORDERS																									
COMMON ELEMENTS																																			
MATERIALS NOTES																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS																																			
COMMON ELEMENTS				</td																															

Investigation of systems with formic acid. II. V. V. Udrovskaya and N. P. Alstraietova. *J. Gen. Chem. (U.S.S.R.)* 17, 605-8 (1947) (in Russian); cf. *C.A.* 43, 2302d. Viscosity, elec. cond., and d. were studied in binary systems for formic acid (I) with Rx formate (II) and with Rx ether (III). No compounds are formed in either system, but there is a weak interaction between I and II. In the I-II system the viscosity decreases smoothly from I to II, the curve being concave towards the concn. axis. Values of the viscosity at 0°, 25, and 50° for I are 2.8210, 1.5372, and 0.9767; and for II, 0.5226, 0.2072, and 0.0300 centipoise. The cond.-concn. curves are similar, except that there are breaks in the curves at about 25 molts of II. Values of the cond. are: I, 0.74, 1.24, 1.76; and II, 0.003, 0.003, and $0.003 \times 10^{-3} \text{ ohm}^{-1} \text{cm.}^{-1}$. Values of the d. are:

1.2375, 1.3088, 1.1846; and II 0.9474, 0.9186, 0.8818 g./cc. No breaks or max. were observed in the curves of d. viscosity, or cond. of I-III series. The viscosity-concn. curve was similar to the I-II system. At 0 and 25°, III has values of: viscosity 0.2368 and 0.2461 centipoises; cond. less than 10^{-7} ohm $^{-1}$ cm. $^{-1}$ at both temps.; d. 0.7223 and 0.7048 g./cc.

ASA-SEA REVALLURGICAL LITERATURE CLASSIFICATION												S-2-107-1000																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
SHELF NUMBER												ALPHABETICAL INDEX																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
SERIAL NO.												ITEM NO. ONE ONLY 101																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
SUBJEC ^T MAP ONE SHE												SECTION																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466	467	468	469	470	471	472	473	474	475	476	477	478	479	480	481	482	483	484	485	486	487	488	489	490	491	492	493	494	495	496	497	498	499	500	501	502	503	504	505	506	507	508	509	510	511	512	513	514	515	516	517	518	519	520	521	522	523	524	525	526	527	528	529	530	531	532	533	534	535	536	537	538	539	540	541	542	543	544	545	546	547	548	549	550	551	552	553	554	555	556	557	558	559	560	561	562	563	564	565	566	567	568	569	570	571	572	573	574	575	576	577	578	579	580	581	582	583	584	585	586	587	588	589	590	591	592	593	594	595	596	597	598	599	600	601	602	603	604	605	606	607	608	609	610	611	612	613	614	615	616	617	618	619	620	621	622	623	624	625	626	627	628	629	630	631	632	633	634	635	636	637	638	639	640	641	642	643	644	645	646	647	648	649	650	651	652	653	654	655	656	657	658	659	660	661	662	663	664	665	666	667	668	669	670	671	672	673	674	675	676	677	678	679	680	681	682	683	684	685	686	687	688	689	690	691	692	693	694	695	696	697	698	699	700	701	702	703	704	705	706	707	708	709	710	711	712	713	714	715	716	717	718	719	720	721	722	723	724	725	726	727	728	729	730	731	732	733	734	735	736	737	738	739	740	741	742	743	744	745	746	747	748	749	750	751	752	753	754	755	756	757	758	759	760	761	762	763	764	765	766	767	768	769	770	771	772	773	774	775	776	777	778	779	780	781	782	783	784	785	786	787	788	789	790	791	792	793	794	795	796	797	798	799	800	801	802	803	804	805	806	807	808	809	810	811	812	813	814	815	816	817	818	819	820	821	822	823	824	825	826	827	828	829	830	831	832	833	834	835	836	837	838	839	840	841	842	843	844	845	846	847	848	849	850	851	852	853	854	855	856	857	858	859	860	861	862	863	864	865	866	867	868	869	870	871	872	873	874	875	876	877	878	879	880	881	882	883	884	885	886	887	888	889	889	890	891	892	893	894	895	896	897	898	899	900	901	902	903	904	905	906	907	908	909	910	911	912	913	914	915	916	917	918	919	920	921	922	923	924	925	926	927	928	929	930	931	932	933	934	935	936	937	938	939	940	941	942	943	944	945	946	947	948	949	950	951	952	953	954	955	956	957	958	959	960	961	962	963	964	965	966	967	968	969	970	971	972	973	974	975	976	977	978	979	980	981	982	983	984	985	986	987	988	989	989	990	991	992	993	994	995	996	997	998	999	1000

UDOVENKO, V. V.

Determination of the molecular weight as a method of physicochemical analysis. II. V. V. Udoenko and S. V. Bubak. *J. Gen. Chem. (U.S.S.R.)* 17, 655-64 (1947) (in Russian); cf. *C.A.* 35, 61754.—The apparent mol. wt. M of a few binary systems of noninteracting components one of which is subject to assocn. was detd. by cryoscopy in C_6H_6 soln. and was plotted against the compn. for the total molalities $m = 0$ (infinite diln.), 0.5, 1.0, and 1.5 moles/1000 g. C_6H_6 . In the system PhMe-MeOH, the $m = 0$ isoconcentrate is a straight line; the $m = 0.5$ curve is convex to the axis of compn., the $m = 1.0$ and 1.5 curves pass through a min. shifting to the PhMe side with rising m . In PhMe-EtOH, $m = 0$ is linear, the $m = 0.5$, 1.0, and 1.5 curves pass through a min. increasingly shifted to PhMe with rising m . In the PhMe- $PbNO_3$, the $m = 0$ isoconcentrate is again rectilinear, the other curves convex to the axis of compn., merging with the $m = 0$ line at 72 mol. % PhMe. The system PhMe- $PhNH_2$ shows the same behavior; extrapolation to $m = 0$ gives for $PhNH_2$ a value of M somewhat higher than the theoretical; this was reproduced in repeated detns. irrespective of addnl. purification. The linearity of M at infinite diln. indicates absence of assocn. The min., and its shift, is the result, on the one hand, of the decrease of M due to decreasing assocn. of the alc. with its increasing diln. by PhMe, and, on the other hand, to the increase of M with increasing PhMe content. The absence of a min. in the case of $PhNO_3$ and of $PhNH_2$ means not only lower assocn. but mainly a lesser tendency to dissoen. on diln. with PhMe. — N. Thon

Determination of the molecular weight as a method of physicochemical analysis. III. V. V. Udrovenco and B. F. Babuk (Mid-Asian State Univ., Tashkent). *J. Gen. Chem. (U.S.S.R.)*, 17, 1583-9 (1947) (in Russian); cf. *C.A.*, 42, 4582; 45, 81785.—Apparent mol. wts. M (determined by cryoscopy in C_6H_6 soln.) of a series of binary mixts. with CHCl_3 are tabulated and plotted as a function of the compn. of the mixt., for four concns.: infinite diln., 0.5, 1.0, and 1.8 mole/1000 g. C_6H_6 (isoconcentrate lines I-IV). (1) In $\text{ROH}-\text{CHCl}_3$, I is rectilinear, II convex to the axis of abscissas, III and IV pass through a min.; the lines draw close with increasing CHCl_3 and merge with I at 93 mole % CHCl_3 . The diagram indicates disso. of assoc. ROH mols., which is borne out by the neg. thermal effect in mixing ROH with a large amt. of CHCl_3 . The chem. interaction indicated by the viscosity isotherm does not appear on the M diagram. (2) In $\text{PhNH}_2-\text{CHCl}_3$, lines I-IV have a shape similar to the corresponding lines of the preceding system; the lines merge with I at a 68 mole % CHCl_3 . The presence of the min. on III and IV , compared with its absence in the system aniline-toluene, indicates increased disso. of assoc. aniline mols. in the presence of CHCl_3 as compared with toluene. (3) In $\text{PAN}_3-\text{CHCl}_3$, all 4 lines are convex and merge at 80 mole % CHCl_3 ; this indicates only disso. (4) $\text{Me}_2\text{CO}-\text{CHCl}_3$: I is rectilinear, III - IV are slightly concave; this indicates some chem. interaction. N. Thom

六

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7"

UDOVENKO, V. V., PROF.

Solution (Chemistry)

Some problems of investigating liquid binary systems. Biul.Sredneaz.un.,
no. 25, 1947.

9. Monthly List of Russian Accessions, Library of Congress, November 1952. Unclassified.

UDOVENKO V. V.

PA 21/49T13

USSR/Chemistry - Heat of Vaporization Oct 48
Chemistry - Mixture

"On Heats of Vaporization of Binary Mixtures, III,"
V. V. Udrovenco, Ts. B. Frid, Cen Asia State U,
Lab Phys Chem, Tashkent, 8 pp

"Zhur Fiz Khim" No 10

Studies vapor pressure of five systems formed by
dichlorethane with alcohols: methyl and ethyl at
40, 50 and 60°, and propyl, isobutyl, and isooctyl
at 50, 60, 70 and 80. Calculates heats of
vaporization of these systems at one temperature
by own formula. Results agree with calculations
using Clapeyron-Clausius equation. Shows that
IC 21/49T13

USSR/Chemistry - Heat of Vaporization Oct 48
(Contd)

formula can also be used to calculate heats of
vaporization of real binary liquid systems.
Submitted 6 Dec 47.

LC

21/49T13

PA 8/49T29

UDEVENKO, V. V.

USSR/Chemistry - Analysis, Physicochemical Apr 48
Chemistry - Molecular Weight, Deter-
mination of

"Determination of Molecular Weight as One Method of
Physicochemical Analysis, IV," V. V. Udevenko,
S. F. Babak, Lab Phys Chem, Cen Asiatic State U,
6¹ pp

"Zhur Obshch Khim" Vol XVIII (LXXX), No 4

Studies following systems by molecular weight method:
methyl alcohol-aniline, methyl alcohol-nitrobenzene,
ethyl alcohol-aniline and ethyl alcohol-nitrobenzene.
Analyzes molecular weight diagrams of systems with
an associated component. Submitted 2 Dec 1946.

8/49T29

RA 8/49T30

USSR/Chemistry - Analysis, Physicochemical Apr 48
Chemistry - Molecular Weight, Determination
of

"Determination of the Molecular Weight as One Method
of Physicochemical Analysis, V," V. V. Udoerenko,
S. F. Babak, Lab Phys Chem, Cen Asiatic State U,
52 pp

"Zhur Obshch Khim" Vol XVIII (LXX), No 4

Studies following systems by the molecular weight
method: ethyl alcohol-acetone, aniline-acetone,
aniline-nitrobenzene and ethyl alcohol-acetic acid.
Shows how molecular weight diagram alters when
interaction of the components occurs at the same time

8/49T30

USSR/Chemistry - Analysis, Physicochemical Apr 48
(Contd)

as dissociation of the associated molecules. Sub-
mitted 2 Dec 1946.

UDOVENKO, V. V.

PA 8/49T31

USSR/Chemistry - Analysis, Physicochemical Apr 48
Chemistry - Molecular Weight, Determination
of

"Determination of Molecular Weight as One Method of
Physicochemical Analysis, VI," V. V. Udrovko, Lab
Phys. Chem., Cen. Asiatic State U, 74 pp

"Zhur. Obschch. Khim." Vol. XVIII (LXXX), No 4

Studies following systems by the molecular weight
method: Chloral-methyl alcohol, chloral-ethyl alcohol,
chloral-¹-acetyl alcohol and chloral-n-butyl alcohol.
Analyzes molecular weight diagrams of systems with
sharply defined chemical reaction of the components.
Shows that irrational maximum is explained, not by
[redacted] 8/49T31

USSR/Chemistry - Analysis, Physicochemical Apr 48
(Contd)

dissociation of the chloralcoholate, but by its
reaction with the alcohol. Submitted 21 Dec 1946.

PA
8/49T31

UDOVENKO, V. V.

V. V. Udoenko and Tz. B. Fried, The heats of evaporation of binary mixtures. I. P. 1136.

An equation has been derived for the heat of evaporation of binary liquid mixtures. For the calculation of this value, one has to know the heats of evaporation and the vapor pressures of the pure components and also their activity coefficients in the mixtures. The vapor pressure of three ideal systems have been studied at 50°, 60°, 70° and 80°: propyl alcohol - isobutyl alcohol, propyl alcohol - isoamyl alcohol and isobutyl alcohol - isoamyl alcohol.

Lab. of Physical Chemistry of the
Central Asiatic State University,
Tashkent
September 24, 1947


SO: Journal of Physical Chemistry (USSR) 22, No. 9, 1948

UDOVENKO, V. D.

V. D. Udoventko and Tz. B. Fried, Heats of evaporation of binary mixtures. II. P. 1135.
This work includes the results of investigation of the vapor pressure and heat of
evaporation of five systems composed of methyl, ethyl, propyl, isobutyl and isoamyl alcohols.
The vapor pressure was studied by the dynamic method. The composition of the liquid and the
vapor were determined refractometrically.

Central Asiatic State University,
Tashkent
December 1, 1947

SO: Journal of Physical Chemistry (USSR) 22, No. 9, 1948



UDOVENKO, V.V., TOROPOV, A.P. and OSININA, M.Ye.

Udovenko, V.V. - "Conductometric titration of anabasine," Doklady Akad. nauk UzSSR, 1949,
No. 1, p. 7-10 --- Summary in Uzbek

SO: U-3566, 15 March, 53, (Letopis 'Zhurnal 'nykh Statey, No. 14, 1949).

"APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7

UDOVENKO, V. V.

Ayrapetova, R. P., Granitova, C. I., ani Ulovenk, V. V. - "Physical-chemical investigation of a formic acid-phenol system", Doklady Akad. nauk UzSSR, 1949, No. 2, p. 13-20, (Resume in Azerbaijani), - Biblio: 14 iters.

SO: U-4392, 19 August 53, (Letopis 'Zhurnal 'nykh Statey, No. 21, 1949).

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7"

UDOVENKO, V. V.

"Research on a system with formic acid IV." Udovenko, V. V. and Airapetova, R. P.
(p.632)

SO: Journal of General Chemistry, (Zhurnal Obshchei Khimii) 1949, Vol. 19, No. 4.

UDOVENKO, V. V.

Chemical Abst.
Vol. 48 No. 4
Feb. 25, 1954
Inorganic Chemistry

Complex compounds of anabasine with cobalt salts.
T. I. Sushkevich and V. V. Udoenko (Mid-Asiatic State
Univ. Tashkent). Doklady Akad. Nauk UzSSR. 3, 57-60,
1949, No. 3, 18-20 (in Russian).—Addn. of anabasine to aq.
 CoCl_4 , with cooling, followed by addn. of HCl to dissolve the
greenish ppt. of hydrated Co, gave blue $\text{CoCl}_4 \cdot \text{C}_8\text{H}_{11}\text{N}_3 \cdot 2\text{HCl}$, m. 283°, when an excess of HCl was used, or the cor-
responding $\text{CoCl}_4 \cdot \text{C}_8\text{H}_{11}\text{N}_3 \cdot \text{HCl}$, m. 303°, with the corre-
spondingly smaller amount of HCl. Both are sol. in H_2O ,
insol. in usual org. solvents, except for hot HCO_2H ; both
are electrolytes in aq. soln., and the cond. indicates complete
dissocn. in soln. of the di-HCl salt with 0 ions being formed;
the mono-HCl salt is a weak electrolyte. Anabasine also
forms complexes with CuCl_4 and MnCl_4 , which are not de-
scribed.

O. M. Kosolapoff

1-27-54

LA

Determination of molecular weight as a method of physicochemical analysis. V. V. Ulyanov. (Central Asian State Univ., Tashkent, U.S.S.R.). Izvest. Sistem. Nauch. Khim. Anal., Inst. Obrabotki Nefti, Khim., Akad. Nauk S.S.R., 17, 101-18 (1949).—The use of the method is illustrated on MeOH and EtOH in C₆H₆. On the abscissa is plotted the concn. of the alc. in C₆H₆ expressed as mol. of alc. per 100 g. of C₆H₆ and on the ordinate the apparent mol. wt. A family of curves is obtained of which the lowest curve is that of MeOH and the upper one of EtOH at zero concns. in C₆H₆. Between these two are the curves of alc. mixts. taken in various proportions but of the same total concn. in C₆H₆. Extrapolation to the ordinate, i.e. at zero concn., gives the mol. wt. of infinitely dil. solns. On the basis of this diagram is plotted a diagram of compn.-mol. wt. on which MeOH-EtOH compn. is the abscissa and the mol. wt. the ordinate. The family of curves (straight lines) thus obtained are isoconcentrations in which the lowest (at zero concn.) and the others at increasingly higher concns. by this method were obtained compn.-mol. wt. curves for a no. of systems. In the system toluene-ethylbenzene, the components do not react with each other and the mol. wt. of the pure components as well as of their mixt. does not change with their concn. in C₆H₆. Therefore, all the isoconcentrations coincide and on the diagram appears one straight line. In the system toluene-MeOH, the latter is an associ. compd. and its degree of association depends on its concn. in C₆H₆ or in its mixt. with toluene. Thus, at zero concn. of MeOH, the isoconcentration is a straight line and as its quantity increases the curves bend toward the abscissa. Similarly are discussed the diagrams of toluene-aniline, nitrobenzene-EtOH, acetone-EtOH, phenol-pyridine, formic acid-Et₂O, methylaniline-phenylisothiocyanate, and chloral-EtOH.

M. Hoss

2

CA

Systems with formic acid. V. V. Uderman and N. P. Akrapetova. Zhur. Otschtech. Khim., 19, 187-44; J. Gen. Chem. U.S.S.R. 19, 142-6 (1949) (English translation); cf. C.A. 42, 3000a. Mol. wt. were determined for various mixts. of HCOOH (I) with PhNO_2 (II), AcOH (III), EtOOCH (IV), and Et_2O (V). The solvent used in the mol. wt. determinations was CaH_2 . Graphs are given showing concns. of dissolved substances, viz., 0.0, 0.5, 1.0, and 1.5 moles per 1000 g. of CaH_2 . The values graphed were extrapolated or interpolated from the exptl. results. In all cases the curves are displaced towards higher mol. wt. by increasing the total concn. In the system I-II, the mol. wt.-concen. curve for the 0.5 molal case is linear, extending from (approx.) 90 for I to 320 for II. In the other three cases the curves are linear in the range 0-80 mole % I. From 80 to 100 mole % I, the 0.0 molal curve is concave, and the 1.0 and 1.5 molal curves are convex, toward the concen. axis, giving values for the mol. wt. of I of, resp.: 94, 95, and 116. In the system I-III, all the mol. wt.-concen. curves are linear, the 0.5 molal curve, e.g., extending from a mol. wt. of about 97 for I to about 128 for III. With increasing total concn. the slopes of the lines decrease. In the systems I-IV and I-V the curves all rises from a mol. wt. of 98 for I to a max. of 100 at 80 mole % I, and then drops to 78 for IV. The sharpness of the max. increases with increasing total concn. In the I-V case, the 0.5 molal curve rises from a mol. wt. of 98 for I to a max. of 106 at 80 mole % I and then drops to 78.2 for V. The mol. wt. of pure V changes less with concn. than that of any other pure substance or soln. investigated, and there is no consistent trend, whereas the mol. wt. of all other substances increased with increasing total concn.

Arlid J. Miller

UDOVENKO, V. V.

58/49T24

USSR/Chemistry - Chloral
Chemistry - Viscometric Analysis Jan 49

"Viscosity of Systems of Chloral With Alcohols,"
V. V. Udroenko, Ye. I. Kalavanskaya, M. F.
Propkopyeva, Lab of Physicochem, Cen Asiatc
State U, 3 1/4 pp

"Zhur Obshch Khim" Vol XIX, No 1

Tabular studies of viscosities and densities
of the system of chloral and methyl alcohol
at 40, 60, and 75° and the system of chloral
and isooctyl alcohol at 40, 60, and 80° show
maximum viscosity of the systems is invariably

58/49T24

USSR/Chemistry - Chloral (Contd) Jan 49

affected by the alcohol, regardless of the
value for the viscosity of the chloral. Sub-
mitted 29 Sep 47.

58/49T24

UDOVENKO, V. V.

PA 67/49T44

USSR/Chemistry - Anabasine May 49
Acids, Mineral

"Reaction of Anabasine With Mineral Acids," V. V.
Udovenko, L. A. Vvedenskaya, Lab of Physicochem,
Cen Asia State U, 1 3/4 pp

"Zhur Obshch Khim" Vol XIX, No 5

Derives the crystalline compounds of anabasine with
hydrogen chloride and nitric acid, respectively:
 $C_{10}H_{14}N_2 \cdot HCl$ and $C_{10}H_{14}N_2 \cdot HNO_3$. Submitted
28 Mar 48.

67/49T44

CA

Systems with formic acid. IV. V. V. Ushatinskii and R. P. Alimpetova. Zhur. Otschitel Khim. 19, 632-6; J. Gen. Chem. U.S.S.R. 19, 885-9 (1949) (English translation); cf. preceding note.—Mol. wts. were determined for various mixtures of formic acid (I) with acetone (II), 2-pentanone (III), and 2-pentanone (IV). As in the previous article, curves are shown for mol. wt. vs. concn. for total concn. of dissolved substances equal to 0.0, 0.6, 1.0, and 1.6 moles per 1000 g. benzene. In the system I-II, concn. and, whereas the 0.0 and 0.6 molal curves are concave towards the e.g., values of concn. of I (mole %) and mol. wt. are, resp.: 100, 107; 60, 108; 65, 102; 32, 98; 0.0, 66. In the system I-III, the curves at all total concns. go through a max. at about 75 mole % I, the 1.0 molal curve going from a mol. wt. of 107 for I, through a max. of 110, then dropping to 74 for III. The system I-IV is similar, the 1.0 molal curve rising from a mol. wt. of 107 for I to a max. of 118 at 70 mole % I, and dropping to 98 for IV. The max. in these curves are attributed to chem. interaction among the components in the various systems. A. J. M.

"APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7

UDOVENKO, V.V.

Viscosimetry in physicochemical analysis. Trudy SAGU no.15:3-16
'50.
(Viscosimetry) (Chemistry, Analytical)

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7"

2A

Viscosity and density of binary systems with nicotine.^a
III. S. P. Babak and V. V. Ulovenko (Samarkand Med. Inst.), Zhur. Obshchel Khim. (J. Gen. Chem.) 20, 1508-71 (1950); cf. C.A. 44, 6230. The nicotine-*o*-nitrophenol system shows a compn., nicotine- $2\text{C}_6\text{H}_4(\text{OH})\text{NO}_2$. Viscosity of the system shows a max. at the above compn.; the highest max. is at 35° 8.0113 centipoises, lower max. at 60° 4.3888, and at 75° 2.0000. Density-compn. curves are smooth curves of linear type going from 0.9090 at 35° and 100 mol.-% nicotine to 1.2615 at 15.63 mol.-% nicotine, at 60° from 0.9090 to 1.2851 at 0.0% nicotine, and at 75° from 0.9070 to 1.2571 at 0.0% nicotine. The nicotine- PhNO_2 system shows a linear viscosity relation: at 25° ranging from 1.7687 (0.0% nicotine) to 3.8042, at 50° from 1.1925 to 2.0376, at 75° from 0.8490 to 1.2620; values of d. similarly range from 1.2003 to 1.0068, from 1.1718 to 0.9366, and from 1.1480 to 0.9070, resp. The system with *o*- $\text{MeC}_6\text{H}_4\text{NO}_2$ has linear viscosity relations: at 25° from 2.0897 (0% nicotine) to 3.8042 (100%), at 50° from 1.3477 to 2.0376, at 75° from 0.9730 to 1.2620; d. varies similarly: from 1.1578 to 1.0098, from 1.1313 to 0.9806, from 1.1000 to 0.9679, resp. The system with *p*- $\text{MeC}_6\text{H}_4\text{NO}_2$ has a linear viscosity relation: at 25° from 2.8922 (41.08 mol.-% nicotine) to 3.8042 (100%), at 50° from 1.2008 (0%) to 2.0376 (100%), at 75° from 0.9381 (0%) to 1.2620 (100%); d. varies similarly: from 1.0793 to 1.0098, from 1.1223 to 0.9806, from 1.1013 to 0.9679, resp.

G. M. K.

CA

2

The viscosity and density of binary systems with nico-
tine. III. S P. Bubuk and V. V. Ulyanenko (Samarkand
Med. Inst.), *J. Gen. Chem. U.S.S.R.* 20, 1029-32 (1950).
See C.A. 43, 3104. R. M. S.
(Engl. translation).

CA

Viscosity of binary solutions with nicotine. IV. S. F. Babak and V. V. Udoenko (Sumarikard Med. Inst.), *Zhur. Obrabot. Khim.* 20, 2121-3 (1980); *J. Gen. Chem. U.S.S.R.* 20, 2190-2201, 2203-5 (Engl. translation); cf. *C.A.* 90, 49724.—*a* and *d*, were detd. for soins. of nicotine (*I*) with CCl_4 at temps. of 25, 35, and 50°, and with 1,2-dichloroethane, PhCl , and PhBr at temps. of 25, 35, and 75°. There is no indication of any chem. interaction between the components in any case. The η -concn. curves in each of the 4 systems is straight line at the highest temp. studied, but becomes increasingly convex towards the concn. axis with decreasing temp. In the system I- CCl_4 , e.g., values of mole % *I*, *d*, and *a*, resp., are: 25°: 0.00, 1.0792, 0.0240; 35.00, 1.3773, 1.5456; 50.00, 1.2261, 2.3747; 75.70, 1.1034, 3.0783; 100.00, 1.0088, 3.8942; at 50°: 0.00, 1.5201, 0.0687; 37.70, 1.2690, 1.2150; 75.70, 1.0773, 1.7311; 100.00, 0.9890.

2.0870. Similar values at 25° for the remaining systems are: I-1,2-dichloroethane: 0.00, 1.2846, 0.7640; 25.33, 1.1495, 1.2300; 50.00, 1.0836, 1.8280; 77.10, 1.0322, 2.4670; I- PhCl : 0.00, 1.0081, 0.7124; 25.87, 1.0578, 2.3657; 50.03, 1.0439, 1.8380; 72.16, 1.0237, 2.3972; I- PhBr : 0.00, 1.4930, 1.0019; 25.14, 1.3201, 1.8435; 48.72, 1.2074, 2.1000; 74.67, 1.0916, 2.9070. *V. Ibid.* 2124-0...*a* and *d*, were detd. at 25, 35, and 40° for the system I-acetone, and at 25, 50, and 75° for the systems I-MeCOR₂ and nicotine-acetophenone (*III*). The first 2 systems have η -concn. curves similar to those mentioned in part IV except that even at the highest temp. investigated the curves are still convex towards the concn. axis. For the system I-II the curves are S-shaped, with the curvature decreasing with increasing temp. Values for mole % nicotine, *d*, and *a*, resp., at 25° are as follows: I-acetone: 0.00, 0.7818, 0.3332; 24.98, 0.8973, 0.6975; 30.05, 0.9408, 1.2797; 74.45, 0.9797, 2.2737; 100.00, 1.0088, 3.8642; I-MeCOR₂: 0.00, 0.8034, 0.4004; 24.63, 0.8922, 0.7187; 48.78, 0.8488, 1.2459; 73.74, 0.9961, 2.9002; I-II: 0.00, 1.0242, 1.0370; 24.26, 1.0223, 2.1076; 49.34, 1.0182, 2.9460; 74.31, 1.0118, 3.6271.

Arndt J. Miller

PASOVSKAYA, G.B.; UDOVENKO, V.V.

Simplified methods for the determination of alkali metals in presence
of magnesium. Trudy SAGU no.27:81-88 '51. (MLRA 9:5)
(Alkali metals) (Conductometric analysis)

PASOVSKAYA, G.B.; UDOVENKO, V.V.

Conductometric simultaneous determination of magnesium and alkali metals. Trudy Kom. anal. khim. 4:196-204 '52. (MIRA 11:6)
(Magnesium)
(Alkali metals)
(Conductometric analysis)

UDOVENKO, V. V.

Chemical Abst.
Vol. 48 No. 8
Apr. 25, 1954
Analytical Chemistry

Conductometric determination of potassium with sodium
perchlorate. V. V. Udoventko and G. B. Pasovskaya (Central
Asian State Univ., Tashkent). J. Anal. Chem. (U.S.S.R.)
7, 181-2 (1952) (Engl. translation). See C.A. 47, 1536.
H. L. H.

(3)

MF 54
11-54

UDOVENKO, V., FATULINA, L. G.

Solubility

Solubility in the system ethyl alcohol - 1,2-ethylene dichloride - water. Zhur.fiz.khim., 16, No. 6, 1952.

Monthly List of Russian Accessions, Library of Congress, November 1952. Unclassified.

UDOVENKO, V. V.

Udovenko, V. V., Airapetova, R. P., Malakhova, V. T.- "Physico-chemical analysis of the systems: phenol-monochloro-acetic acid, and phenol-trichloroacetic acid." (p. 1759)

SO: Journal of General Chemistry, (Zhurnal Obshchei Khimii), 1952, Vol. 22, No. 10

Chemical Abst.
Vol. 48 No. 9
May 10, 1954
General and Physical Chemistry

(b) Chem

Physicochemical analysis of the systems phenol-mono-chloroacetic acid and phenol-trichloroacetic acid. V. V. Udoenko, R. P. Arapetova, and V. T. Malakhova. J. Gen. Chem. (U.S.S.R.) 22, 1801-2 (1952) (Engl. translation). —See C.A. 47, 2028f. H. L. II

UDOVENKO, V.V.

3 Chemad

Chem Abv V48
1-25-54
Inorganic Chemistry

Some compounds of pyridine with cupric chloride. V. V. Ulovenko¹ and E. M. Osipova² (Mid-Asian State Univ., Tashkent). Zhur. Obshch. Khim. 27, 2935-7 (1952).-- Equimol. amts. of anhyd. CuCl₂ and C₆H₅N, in abs. alc. or dry Me₂CO form light-blue CuCl₂.2C₆H₅N (I), which slowly (in a few hrs.) changes into gray CuCl₂.C₆H₅N (II). With HCl, I forms 1.2HCl, yellow prisms, and dark brown crystals of 3CuCl₂.2C₆H₅N.2H₂O (III). This reaction can take place in moist HCl vapor, but not in dry HCl gas. II is not sol. in org. solvents; in H₂O, it is decompd. into I and CuCl₂. Heated above 200°, I loses C₆H₅N becoming anhyd. CuCl₂. III is obtained from aq. solns. contg. strictly stoichiometric amts. of CuCl₂ and C₆H₅N and an excess of HCl; without excess HCl, only II is ptd., but it redissolves in excess HCl and, on evapn., gives III, which can be recrystd. from H₂O. In alc., III decomp. to I, and in Me₂CO to 1.2HCl, insol. in C₆H₆, CCl₄, etc. III m. 194-5° (decompn.). In air, III loses H₂O and HCl, becoming 3CuCl₂.2C₆H₅N (IV), a light-green powder, which cannot be obtained directly from CuCl₂ and C₆H₅N. In H₂O, IV decomp. to I. With HCl, it forms III. It is stable in air at room temp., decomp. above 100°, and m. 272-3° (decompn.). Blue CuCl₂.5C₆H₅N was obtained with CuCl₂ and a large excess of C₆H₅N, within 10 days at a temp. not over 12-15°, faster at -10°. N. Thon

UDOVENKO, V. V.

Chemical Abst.
Vol. 48 No. 9
May 10, 1954
inorganic Chemistry

Some compounds of pyridine with cupric chloride. V. V. Udoventko and E. M. Osipova. J. Russ. Chem. U.S.S.R. 27, 2146-50 (1952) (Engl. translation).—See C.A. 48, 490b. H. L. H.

(2) ref. m

9-2 gfp

*General 71
Chemistry 72*

CA

Solubility in the system ethyl alcohol-1,2-dichloroethane-water. V. V. Udovenko and L. G. Fatkulina (Sredneasiat. Gosudarst. Univ., Tashkent). Zhur. Fiz. Khim. 26, 892-7 (1952). Solv. of H₂O in C₂HCl₂ was find. between 10° (0.0357 wt. %) and 00° (0.818%). Solv. of C₂HCl₂ in H₂O was, e.g., 0.877% at 20° and 1.380% at 72.5°. Over 50 ternary poly. were measured. E.g., EtOH 21.7%, H₂O 76.24% dissolved 1.490% C₂HCl₂ at 25°; EtOH 32.3, H₂O 47.7% dissolved 18.45% C₂HCl₂ at 34°; EtOH 31.5, C₂HCl₂ 68.5% dissolved 12.14% H₂O at 83°; and EtOH 10.4, C₂HCl₂ 89.4% dissolved 1.07% H₂O at 44°. Also d. and s. of many solns. at 30° and 40° and d. of solns. at 60° and 80° were detd. The compn. of conjugated layers is listed. Thus, at 30° the liquid H₂O 92.7, EtOH 6.3, C₂HCl₂ 1.0% was in equil. with H₂O 0.5, EtOH 1.0, C₂HCl₂ 98.5% and 51.5, 39.5, 9.0% in equil. with 3.0, 19.0, 84.0%. At 40°, 88.8, 9.0, 1.3% was in equil. with 7.0, 22.3, 70.7%. At 60° 80.8, and 30.0, 41.1, 10.0% with 7.0, 22.3, 70.7%. At 80° 80.8, 17.0, 2.2% was in equil. with 0.9, 4.2, 94.9% and 41.7, 39.1, 19.2% with 7.2, 22.0, 70.8%. At 60° 90.1, 7.0, 2.0% was in equil. with 0.6, 1.8, 97.7%. The calcd. compn. of the crit. solns. was at 30° H₂O 18.3, EtOH 34.3, C₂HCl₂ 47.4% at 40° 19.2, 32.9, 47.9%, and at 60° 19.6, 31.4, 49.0%. J. J. Bikerman.

UDOVENKO, V. V. FATKULINA, L. G.

Chemical Apparatus.

Apparatus for determination of pressure and composition of saturated vapor of layer-forming liquid systems. Zhur.fiz.khim. 26, no. 2, 1952.

9. Monthly List of Russian Accessions, Library of Congress, September 1952 Unclassified.

UDOVENKO, V. V., FAKULINA, L. G.

Phase Rule and Equilibrium.

Vapor pressure of ternary systems. Part 1. System ethyl alcohol - 1,
2-dichloroethanebenzene. Zhur. fiz. khim. 26 no. 5, 1952.

9. Monthly List of Russian Accessions, Library of Congress, November 1952 Unclassified.

1. UDOVENKO, V. Y.: FATKULINA, L. G.
2. USSR (600)
4. Systems (Chemistry)
7. Vapor pressure of ternary systems. Part 2. System ethyl alcohol-1, 2-dichloroethane water. Zhur. fiz. khim. 26 no. 10, 1952.
9. Monthly List of Russian Accessions, Library of Congress, _____ March 1953, Uncl.

UDOVENKO, V. V.

FA 242T3

USSR/Chemistry - Ternary Systems

Nov 52

"The Equilibrium of Liquid-Liquid in a Ternary System," V. V. Udoeneko and I. G. Fatkulina, Central Asian State U, Tashkent

"Zhur Fiz Khim" Vol 26, No 11, pp 1569-1572

The authors examined the eq of L. A. Rotinyan and showed that in the case of a paraboloidal curve of sepn in ternary systems, the nodes can be parallel to the side of the triangle when the liquid phases are in equil. On the basis of the above-mentioned eq, the authors concluded that the critical point of mutual solv in ternary systems can also be located at the vertex of the binodal curve of sepn.

242T3

"APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7

UDOVENKO, V.V.; PASOVSKAYA, G.B.

Conductance method for the study of the adsorption of electrolytes.
Trudy SAGU no.33:35-37 '52. (MLRA 9:5)
(Electrolytes) (Adsorption)

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7"

UDOVENKO, V.V.

USSR:

✓Reaction of lupinine and diethylaminooethanol with copper chlorides. V. V. Udoenko and L. P. Aleksandrova (Central Asia State Univ., Tashkent). Sovetsk. Sistem. Otschete Khim. 2, 1120-3(1953).—Lupinine and dry CuCl₂ in abs. EtOH form brown CuH₄NOH·CuCl₂, decomp. 125°, which slowly decomposes on exposure to air. If twice the proportion of lupinine is used, there is obtained a green alcoholate C₁₀H₁₄NO·Cu(I) (I), m. 215° (from CH₃CH₂Cl), while the evapd. filtrate yields lupinine HCl salt, m. 211.5°. I also forms on mixing abs. EtOH soln. of lupinine with CuCl₂ in contact with atm. O₂. Addn. of abs. EtOH soln. of Et₂N-CH₂CH₂OH to EtOH soln. of CuCl₂ gave green Et₂NCH₂-CH₂O·CuCl, m. 160° (from ClCH₂CH₂Cl), while evapn. of the soln. gave the amine HCl salt. Thus the original adduct is transformed into the alcoholate under action of excess amino alc., which results in loss of HCl and formation of I analog and amine HCl salt. G. M. Kholopoff

УДОВЕРЕНІСТЬ В.В.

У ССР.

✓ Separation of mixture of alkaloids anabasine and lupinine.
V. V. Uloverko, O. I. Grinova, and L. A. Vvedenskaya
(Central Asian State Univ., Tashkent). *Soviet States*
Ustiches Khim., 2, 1123-6 (1931). — The sepn. is performed
through the Cu salts (cf. following abstr.) as follows. The
mixt. of known compn. of anabasine and lupinine is treated
with a highly concd. soln. of CuCl_2 and HCl or $\text{Cu}(\text{NO}_3)_2$ and
 HNO_3 , made up on the basis of theoretical calcn. to form
either $\text{CuCl}_2 \cdot 2\text{HCl}$ or $\text{Cu}(\text{NO}_3)_2 \cdot 2\text{A} \cdot 4\text{HNO}_3$ (A = ana-
basine), with cooling. The order of addn. is immaterial.
Since the mixt. now contains a ppt. of Cu oxide, addn. acid
is added to dissolve this (avoid excess of HNO_3). Traces of
kerosine from com. alkaloids are removed at this point by
boiling. The mixt. then yields cryst. salt of anabasine,
which treated with concd. NaOH gives the pure base, which
is purified by distn. *in vacuo*. The evapd. filtrate from the
salt is treated with concd. NaOH and extd. with CH_2Cl_2 .
Evapn. of the ext. yields cryst. lupinine. The HCl salt
gives somewhat better results. Pure anabasine bp 107-10°,
 n_D^{20} 1.5120; D_4 1.0469; lupinine m. 65°. C. M. E.

UDOVENKO, V. V.

USSR.

Reaction of anabasine with salts of copper. V. V. Udrovenco and O. I. Granitova (Central Asian State University, Tashkent). *Sovetsk. Nauki Chishchel Khim.* 2, 1127-9 (1953).—Anabasine (A) forms 2 salts with CuCl₂ and HCl and with CuBr₂ and HBr, resp. With control of the reagent proportions there were obtained: yellow CuBr₂.A.2HCl, decomp. 205° (in aq. soln. this dissociates into components on basis of cond. meas.); CuCl₂.A.HCl, green, decomp. 191-6° (completely dissolved, in aq. soln.); black CuBr₂.A.2HBr, decomp. 214°; red-brown CuBr₂.A.HBr,

decomp. 157°. Similarly was obtained blue-violet Cu(NO₃)₂.A.4HNO₃, decomp. 107°, and blue CuSO₄.2A. aq. salts.

G. M. Kosolapoff

UDOVENKO, V.V.

S S R

Interaction of anabasine with mercuric and mercurous chlorides. V. V. UDOVENKO, M. A. ADYEV and T. I.

ADYEV. Doklady Akad. Nauk Uzbek. SSR 1953, No. 5, 38-41; Referat. Zhur., Khim. 1954, No. 27, 97. HgCl₂ reacted with anabasine (I) in aq. soln. to form a white amorphous compd. HgCl₂C₁₉H₁₄N₂ (II), m.p. 211° (decompn.). The reaction proceeds with appreciable evolution of heat and unless cooled a colored compd. is obtained. II was obtained by adding dropwise an aq. soln. of I to an aq. soln. HgCl₂ cooled in ice water. Upon heating to 110° II turned yellow. It is nonhygroscopic, dissolves poorly in water (slightly better in hot water), acetone, and EtOH, insol. in benzene and toluene. After boiling, aq. solns. of II have an alk. reaction. II dissolves in I, liberating metallic Hg; addn. of Me₂CO hastens this reaction. In water acidified with AcOH or strong mineral acids II dissolved forming a compd. of different compns. Specifically aq. HCl formed HgCl₂C₁₉H₁₄N₂·2HCl, acicular crystals. HgCl₂(III) reacted with I according to: HgCl₂ + C₁₉H₁₄N₂ + HgCl₂ → C₁₉H₁₄N₂. A comparative study was made of the reaction of I, pyridine (IV), and piperidine (V) with III. The reactions were carried out with an excess of base in the absence of moisture. Immediately following the mixing, after 10 min., and after 7 days, the extent of interaction of I with III was 7, 9, and 18.9%, resp. For IV and III it were 30, 75.65, and 81.27%, and for V and III it was 11 and 15.96%, resp. In the last case, no detn. was made after 7 days. Thus, in its action on dry III, I is closer to V than to IV. In the presence of water, the interaction of I or IV with III does not exceed 1%. M. H.

"APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7

U.S. VenKo, U.U.

5
6
8

Interaction of substances with membrane filters and

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7"

Reaction of anabasine with organic acids. V. V. Udrovenco and L. A. Vvedenskaya (Central Asia State Univ., Tashkent) Zhur. Obschchet Khim. 23, 1031-4 (1973). Anabasine (I) and HCO_2H yield crystalline $\text{I} \cdot \text{HCO}_2\text{H}$, very hygroscopic and sol. in org. solvents. I and AcOH gave $\text{I} \cdot \text{AcOH}$, m. 88°. PrCO_2H similarly gave $\text{I} \cdot \text{PrCO}_2\text{H}$, very hygroscopic solid. I and $\text{i-BuCO}_2\text{H}$ gave $\text{I} \cdot \text{i-BuCO}_2\text{H}$, very hygroscopic solid. With $(\text{CO}_2\text{H})_2$ there are formed: $2\text{I} \cdot \text{C}_2\text{O}_4\text{H}_2$, m. 210°, $\text{I} \cdot \text{C}_2\text{O}_4\text{H}_2$, m. 190°, and $1 \cdot 2\text{C}_2\text{O}_4\text{H}_2$, m. 81°; all are very sol. in the usual solvents. Malonic acid gave $2\text{I} \cdot \text{C}_4\text{O}_4\text{H}_4$, m. 80°. Succinic acid gave very hygroscopic $2\text{I} \cdot \text{C}_4\text{O}_4\text{H}_4$. Tartaric acid gave solid $2\text{I} \cdot \text{C}_4\text{H}_4\text{O}_4$. Citric acid gave solid $3\text{I} \cdot \text{C}_6\text{H}_8\text{O}_7$. With naphthalene-2-sulfonic acid there are formed: $\text{I} \cdot \text{C}_10\text{H}_7\text{SO}_3\text{H}$, m. 139°, and $1 \cdot 2\text{C}_10\text{H}_7\text{SO}_3\text{H}$, m. 121°.

G. M. Kosolapoff

"APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7

UDOVENKO, V.V.

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7"

"APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7

Udovenko, V V

The strength of come ~~is~~ in water and acetone

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7"

USSR

U S S R

Heats of vaporization of three-component liquid mixtures.
V. V. Udrovko and I. G. Fatkulina (Central Aerohydrodynamic Institute, Tashkent), Zhur. fiz. khim., 27, 1063-7 (1953);
cf. C.A. 47, 6748i.

The heats of vaporization (L) of ternary mixts. contg. 10-40 mole % 3OH (I) with varying proportions of benzene (II) and 1,2-dichloroethane (III) were calc'd. by means of the Clausius-Clapeyron equation from exptl. data published earlier (*loc. cit.*). Values of L so calc'd. agree within 1% or less on the av. with those calc'd. by means of the equation $L = L_{1j} \cdot x_j + L_{2j} \cdot y_j$, where L_{ij} is the heat of vaporization of the i th component and y_j is its mole fraction in the gas phase. Both sets of values are tabulated. The equation $L = (L_{1j}x_j + L_{2j}y_j + L_{3j}z_j)/(\rho_1x_j + \rho_2y_j + \rho_3z_j)$, where ρ_i , x_j , and z_j are the vapor pressure in the pure state, the activity coeff. and the mole fraction of the j th component in the liquid phase, resp. was derived. J. W. Lowenberg, Jr.

(1) 9

"APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7

LIDOVENKO, V. V.

U S S R .

Calorimetric titration of a mixture of anabasine and

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7"

LIDOVENKO, V. V.

USSR

Viscosity of some binary systems with nicotine. P. V. Lidovenko and S. V. Balak. *Trudy Sredaz. Gos. Akad.*, Ural, Khim. Nauki 5, No. 30, p. 161-163; *Referat. Zhar.*, Khim., 1954, No. 2018; cf. *C.A.*, 45, 933d, 46, 2601g. The viscosity and d. of mixts. of nicotine with CHCl_3 , CH_2Br_2 , tetrachloroethane, EtOH , iso-BuOH, AmOH , and cyclohexyl alc. were studied at 25, 50, and 75° and of nicotine and MeOH at 25 and 50°. The nature of the relation between the viscosity values and molar compn. indicates interaction between nicotine and CHCl_3 , ethyl ene bromide, tetrachloroethane, and Me, Et, and iso-Bu ales. The interaction of these substances with nicotine indicates their acidic nature in relation to nicotine as a base. The viscosity isotherm of nicotine mixt. with cyclohexyl alc. shows no reaction between the two. The acidic properties of the halogen derivs. increase with the no. of halogen atoms in the mol. and increases also upon the replacement of Cl by Br. In the case of ales., an increase in the length of the hydrocarbon chain attenuates their interaction with nicotine

M. Hosszú

UDOVENKO, V.V.

✓ 1842. Conductometric titration of certain alkaloids.

V. V. Udoenko and L. A. Vvedenskaya. *Ukr. Khim. Zh.* 1964, 20 (6), 684-689.—By conductometric titration with naphthalene-2-sulphonic acid in acetone solution, the alkaloids present in the following mixtures were separately determined: anabasine - lupinine, anabasine - nicotine, anabasine - nicotine - lupinine, also anabasine - lupinine in the presence of aphylline and aphylidine. The method depends on the effect of acetone in increasing the difference between the ionisation constants of the bases. It can be applied to the determination of alkaloids in technical products.

E. W. KIRKLAND

(2)

"APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7

Vdovchenko V.V.

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7"

"APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7"

"APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7

DOCUMENT FOR

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7"

AUTHORS: Udovenko, V.V. and Fialkov, Yu. Ya. DCI

TITLE: Viscosity of Germanium Tetrachloride - Ethers Systems. (Vyazkost' Sistem Chetyrekhkhloristy Germaniy - Efiry).

PERIODICAL: "Zhurnal Neorganicheskoy Khimii" (Journal of Inorganic Chemistry) Vol.II, No.2, pp.434-438 (U.S.S.R.).1956

ABSTRACT: A study of the complex-forming ability of germanium tetrachloride would be useful in enabling a comparison to be made of the properties of various addition products of IV group element tetrahalides; the rules governing the formation of these products could also be deduced. In connection with this studies have been made of the viscosities of binary systems of which one component was always germanium tetrachloride, the others being various organic compounds. All materials were carefully dried to avoid hydrolysis of the tetrachloride. Measurements were made of the viscosity and also the density of the following systems: germanium tetrachloride - acetoacetic ester, germanium tetrachloride - anisole at 20, 30 and 40°C; germanium tetrachloride - dioxane at 25 and 40°C, and germanium tetrachloride - diethyl ether and germanium - tetrachloride - dimethyl sulphide at 200°C. It was found that in contrast to titanium and tin tetrachlorides, germanium tetrachloride does not form addition products with the above type of compounds; this is explained as being connected with the increase in the

Card 1/2

581

Viscosity of Germanium Tetrachloride - Ethers Systems. (Cont.)
screening of the central atom in germanium tetrachloride.
There are thirty-one references, of which fifteen are Russian.
One Figure and Five Tables.
The work was carried out at Kiev Polytechnical Institute.
Received 17 September, 1956.

Card 2/2

UDOVENKO, V.V.

Viscosity of binary systems with chloral. I. V. V.
Udovenko and P. I. Kolesnikov
Reported in Russ. J. Phys. Chem. 1959, 33, 1001
of the system CCl_3CHO (molar) at 5, 10, and 20%, and
of CCl_3CHO (molar) at 5, 10, and 20% were done. The
systems show component interaction although a cryst
adduct with cyclohexanone was the only definitely soluble
product, in this case the viscosity is observed in irrational
max. at 20 mole % aldehyde. A equimolar mixt. of com-
ponents forms a 1:1 adduct, in 94% alc. (Dowesford and
Cronje, C.A. 42, 2316). The system with PhCH_2OH failed
to yield a cryst. adduct but gave a clear max. in the vis-
cosity curve at 70 mole % alc. G. M. Kolesnikov

Udovenko V. V.

Viscosity of binary systems from chloroform, II

Udovenko and R. I. Klimenko, *Bull. Acad. Sci. USSR*, 1960, No. 10, p. 2261

5306A viscosity of the system, CCl_3CHO -PhOH at 40°, 60°, and 80°, and CCl_3CHO -*m*-MeC₆H₄OH, and CCl_3CHO -*p*-MeC₆H₄OH at 50°, 59° and 75°, was dekd. There was indication that CCl_3CHO forms complexes with the 4 phenols that dissociate into components when the temp. rises. The complexes are formed through H bonding and are not measurable. The system CCl_3CHO -PhOH yields a little $\text{CCl}_3(\text{CH}_2\text{CH}_2\text{OH})_2$, m. 203°, owing to condensation under the influence of traces of HCl. Similarly a small amt. of 2,5-(*i*O₂M₂)C₆H₃(OH)₂CCl₃, m. 147°, was isolated. The system of PhOH- CCl_3CHO shows an irrational viscosity max. at 25 mole % CCl_3CHO ; this indicated a compd. $\text{CCl}_3\text{CHO}\cdot\text{PhOH}$; the system with *p*-cresol has an irrational max. that does not yield any rational component proportion; with *m*-cresol the system has an irrational max. at 25 mole % CCl_3CHO , as does that with *p*-cresol.

G. M. Kondratenko

MDOVENKO V V.

Viscosity of binary systems with chloro- III. V. V. Udo-
vits and R. A. Miller, Certified Anal. Supplied by
Ferakent and Politekhn. Inst., Moscow, 285, 1952, 245 N. 11th
St., St. Paul 2, Minn. Viscosity and density measurements
made for the following systems: $\text{CH}_3\text{COCl}-\text{CH}_3\text{COCl}$ at 25°,
60°; $\text{McCOC}-\text{CCl}_3-\text{CHO}$ at 25°, 50°, and 75°; $\text{C}_6\text{H}_5\text{COCl}-$
 $\text{C}_6\text{H}_5\text{COCl}$ at 65°, 75°, and 80°; $\text{C}_6\text{H}_5\text{COCl}-\text{CH}_3\text{COCl}$ at 25°, 50°, and 75°. The only evi-
dence of interaction between $\text{C}_6\text{H}_5\text{COCl}$ and CH_3COCl is found in the
viscosity measurements.

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7"

UDOVENKO V. V.
USSR/Physical Chemistry - Thermodynamics, Thermochemistry,
Equilibria, Physical-Chemical Analysis, Phase Transitions. B-8

Abs Jour : Referat Zhur - Khimiya, No 1, 1958, 414

Author : V.V. Udoventko, Yu.Ya. Fialkov.

Inst :

Title : Viscosity of Systems Germanium Tetrachloride - Ethers and
Esters.

Orig Pub : Zh. neorgan. khimii, 1957, 2, No 2, 434-438

Abstract : The viscosity and density of binary systems composed by germanium tetrachloride (I) with ethyl acetate (II), anisole (III), dioxane (IV), diethyl ether (V) and dimethylsulfide (VI) were measured. The systems I - II and I - III were studied at 20, 30 and 40°, the system I - IV was studied at 25 and 40°, and the systems I - V and I - VI were studied at 20°. All the operations of preparing the solutions and carrying out the measurements were done under airtight conditions. Viscosity was measured

Card 1/2

USSR/Physical Chemistry - Thermodynamics, Thermochemistry,
Equilibria, Physical-Chemical Analysis, Phase Transitions. B-8

Abs Jour : Ref Zhur - Khimiya, No 1, 1958, 414

in a closed viscosimeter for volatile liquids described earlier (Toropov A.P., Zh. prikl. khimii, 1939, 12, 1744). The viscosity isotherms of the studied systems are convex with reference to the composition axis, which indicates the absence of a chemical interaction between the components. The authors connect the absence of interaction with an increase of the screening of the central atom in I.

Card 2/2

"APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7"

446

AUTHORS:

Udovenko, V. V., and Khomenko, R. I.

TITLE:

Viscosity of Binary Systems with Chloral.4 (Vyazkost' binarnykh sistem s khloralem. IV.)

PERIODICAL:

Zhurnal Obshchey Khimii, 1957, Vol. 27, No. 1, pp. 37-40 (U.S.S.R.)

ABSTRACT:

Investigations were conducted to determine the viscosity and density of binary systems consisting of chloral with acetophenone (boiling point 80.0° at 12 mm); benzophenone (b. p. 161.0° at 11 mm); benzaldehyde (b. p. 177.2° at 731 mm) and salicylaldehyde (b. p. 193.0° at 727 mm). The viscosity/density measurements were carried out at temperatures of 25, 50 and 75° and the results obtained are given in tables (for each mixture separately). The isothermal viscosity/density curves are either slightly convex or rectilinear, depending upon the temperature. None of the systems investigated showed any definite reactions between the components which could lead to the formation of any specific chemical compound.

Four tables. There are 5 references, of which 4 are Slavic.

Card 1/2

446

Viscosity of Binary Systems with Chloral.4

ASSOCIATION: Central Asiatic State University and the Kiev Polytechnicum
(Sredneaziatskiy Gosudarstvennyy Universitet i Kiyevskiy
Politekhnicheskiy Institut).

PRESENTED BY:

SUBMITTED: January 3, 1956

AVAILABLE:

Card 2/2

UDOVENKO, J. V.

79-2-9/58

AUTHORS:

Udovenko, V. V., and Khomenko, R. I.

TITLE:

Viscosity of Binary Systems with Chloral. Part 5. (Vyazkost' binarnykh sistem s khloralem. V.)

PERIODICAL:

Zhurnal Obshchey Khimii, 1957, vol 27, No 2, pp. 322-325 (U.S.S.R.)

ABSTRACT:

In order to explain the nature of the reaction between chloral and ether, ethyl formate, chloral - ethyl acetate, chloral-ethyl ether, chloral - anisole and chloral - acetoacetic ester, chloral-ethyl acetate, chloral - 75°. Viscosity was found to be constant in the first four systems and in the fifth one it varies with time. A reaction between the components was established in all systems investigated but only the reaction of the chloral-acetoacetic ester system is clearly expressed and this is explained by the presence in the molecule of the latter of a hydroxyl group.

It was established that acetoacetic ester in the presence of pyridine reacts with chloral forming a compound which during distillation in vacuum

Card 1/2

Viscosity of Binary Systems with Chloral. Part 5 79-2-9/58
decomposes into chloral and acetoacetic ester. Optical investigation
of acetoacetic ester solutions in pyridine showed that the latter like
alcohol, reacts with the pyridine by means of the hydroxyl group. Also
the reaction between the chloral and the acetoacetic acid is due to this
hydroxyl group of enol form.

4 tables, 1 graph. There are 2 references, of which 2 are Slavic.

ASSOCIATION: The Kiev Polytechnicum

PRESENTED BY:

SUBMITTED: January 25, 1956

AVAILABLE: Library of Congress

Card 2/2

UDOVENKO, V.V.: KHOMENKO, R.I.

Viscosity of binary systems containing chloral. Part 6. Zhru. ob. khim.
27 no.3:583-585 Mr '57. (MIRA 10:6)

1. Sredneaziatskiy gosudarstvennyy universitet i Kievskiy politekhnicheskiy institut.
(Systems (Chemistry)) (Chloral)

UDOVENKO, V. V.

USSR/Inorganic Chemistry. Complex Compounds.

C

Abs Jour: Ref. Zhur-Khimya, No 1, 1958, 677.

Author : Udoventko, V.V., Fialkov, Yu. Ya.

Inst : Kiev Politech Inst.

Title : Interaction of Silicon Tetrachloride with Acetic and Benzoic Anhydrides.

Orig Pub: Zh. Obshch. Khimii, 1957, 27, No 4, 905-906.

Abstract: It was established that SiCl_4 can interact directly with anhydrides of organic acids thereby forming mixed anhydrides of organic and orthosilicic acids. This reaction proceeds under less rigid conditions than the reactions of interaction of SiCl_4 with corresponding acids. By means of the above reaction the tetraacetate and tetrabenoate of silicon were obtained in good yields.

Card : 1/1

-17-

AUTHORS:

Udovenko, V. V., Fialkov, Yu. Ya.

79-28 -3-54/61

TITLE:

The Viscosity of Binary Systems With a Substitution Reaction (Vyazkost' dvoynikh sistem s obmennym vzaimodeystviyem)

PERIODICAL:

Zhurnal Obshchey Khimii, 1958, Vo. 28, Nr 3,
pp. 814-818 (USSR)

ABSTRACT:

The viscosity diagrams of binary systems are to a great extent systematized at present and are often used for the explanation of the occurring processes. When systems with chemical reactions are considered the present classification (Ref. 1) provides diagram types for such systems in which chemical reactions take place with a decrease of the molecular number, e. g. in the system water-chloral, or for systems in which the molecular number does not change, e. g. in the system acetic anhydride-water. Such systems have been little investigated. Therefore N. A. Trifonov suggested such model systems as, among other, diethyl-water systems conducive to visualization of the type of the diagram of viscosity when only

Card 1/3

The Viscosity of Binary Systems With a Substitution Reaction

79-28 -3-54/61

one chemical compound without a decrease of the molecular number is formed. It must be noted that in the system acetic anhydride-water the number of molecules remains the same after the chemical reaction had taken place as two molecules of acetic acid are formed from the molecules of this anhydride and water. The reaction leading to the formation of a chemical compound is not the only possible^{one} for reactions where the molecular number remains unchanged. Not less frequent is the substitution reaction where the final products are two chemical compounds. Systems of this kind are of great interest for the theory of physical and chemical analysis; they have, however, not been investigated by any chemical scientist with respect to the viscosity method. Below, data on the viscosity of systems are mentioned in which one component is silicontetrachloride and the other one of the following compounds: methylal (dimethoxymethane), acetal (1,1-diethoxyethane) and acetic anhydride. According to the methods of viscosity and density the binary systems silicontetrachloride-methylal at 20 and 30°C, silicontetrachloride-acetal at 20, 30 and 40°C and silicontetrachloride-acetic acid anhydride in benzene

Card 2/3

The Viscosity of Binary Systems With a Substitution Reaction 79-23 5-54/6:

as indifferent medium at 20°C were investigated. The authors found that in the reaction of silicontetrachloride with methylal the final products are: dichlorodimethoxysilane and chlorodimethylether, and with acetal: dichloroethoxysilane and chlorodiethylether. This reaction represents a new method of the synthesis of dichlorodialkoxysilane which differs from the existing ones by its good yield and by the purity of the products. There are 1 figure, 3 tables and 5 references; 4 of which are Soviet.

ASSOCIATION: Kiyevskiy politekhnicheskiy institut (Kiyev Polytechnical Institute)

SUBMITTED: March 3, 1957.

Card 3/3

AUTHORS: Udovenko, V. V., Aleksandrova, L. P. SOV/76-32-8-26/37

TITLE: The Solubility in the System Formic Acid - 1,2-Dichloro Ethane - Water (Rastvorimost' v sisteme murav'inaya kislota - 1,2-dikhloretan - voda)

PERIODICAL: Zhurnal fizicheskoy khimii, 1958, Vol. 32, Nr 8, pp. 1889-1892
(USSR)

ABSTRACT: The mutual solubility in the above mentioned ternary system was investigated within the temperature range of from 20 to 70° according to the polythermal method by V. F. Alekseyev. The data of the mutual solubility of water and dichloro ethane were taken from other publications. In the determinations of the solubility eight polythermal lines were taken at different constant ratios between formic acid and dichloro ethane, and the results obtained were given in a table. There exists a greater laminated section in the system which decreases on an increase in temperature. For determining the intersections in the system the compositions of the layers of equilibrium were determined by a titration of the acid in both layers at 30, 45 and 60°C. The kinetic points of the mutual solubility on the curves of the separation of layers were determined according

Card 1/2

SOV/76-32-8-26/37

The Solubility in the System Formic Acid - 1,2-Dichloro Ethane - Water

to the method mentioned above. The coefficients of distribution were calculated. The difference observed between the system $H_2O-C_2H_5OH-C_2H_4Cl_2$ and that investigated was explained by the greater coefficient of distribution of formic acid as compared to that of ethanol. There are 5 tables and 3 references, 2 of which are Soviet.

ASSOCIATION: Kiyevskiy politekhnicheskiy institut (Kiyev Polytechnical Institute)

SUBMITTED: March 25, 1957

Card 2/2

5(4)

SOV/78-4-2-17/40

AUTHORS:

Udovenko, V. V., Artemenko, M. V.

TITLE:

On the Interaction of Copper Chloride With Monoethanolamine
(O vzaimodeystvii khlornoy medi s monoetanolaminom)

PERIODICAL:

Zhurnal neorganicheskoy khimii, 1959, Vol 4, Nr 2,
pp 352-355 (USSR)

ABSTRACT:

On the interaction of monoethanolamine with copper salts complex compounds of various compositions are formed. The following compounds are formed by copper chloride with ethanol amine combined with hydrochloric acid: $[\text{CH}_2\text{OHCH}_2\text{NH}_3]\text{CuCl}_3$ and $(\text{CH}_2\text{OHCH}_2\text{NH}_3)_2\cdot\text{CuCl}_4$. On the interaction of a concentrated alcoholic solution of copper chloride with monoethanol amine the compounds $\text{CH}_2\text{NH}_2\text{CH}_2\text{OCuCl}$ and $(\text{CH}_2\text{OHCH}_2\text{NH}_3)\text{CuCl}_3$ are formed. In weak solutions of copper chloride in acetone the compound $\text{Cu}(\text{CH}_2\text{OHCH}_2\text{NH}_2)_4\text{Cl}_2$ is formed. This compound crystallizes in well formed blue crystals whose melting point is 103° ; it is soluble in alcohol and water, and insoluble in acetone. On the interaction of copper chloride with $\text{Cu}(\text{CH}_2\text{OHCH}_2\text{NH}_2)_4\text{Cl}_2$

Card 1/2