

S/076/62/036/005/006/013
B101/B110

Electrochemical evolution of...

(2) Oscilloscopic measurement of the potential by an ENO-1 (ENO-1) oscilloscope, synchronously connected with a sawtooth pulse generator, showed no change of the polarization curve for n-type Si, and an increase of the potential by 0.35 v for p-type Si. (3) The oscillograms for current insertion are equal for both types at $I_0 = 10^{-4}$ a/cm². At $I_0 = 10^{-3}$ a/cm², the curve for p-type Si shows a distinct peak 2 v high. (4) The anodic charging curves for Si polarized at -0.5 v show a retardation of the potential at $I_0 \geq 5 \cdot 10^{-5}$ a/cm². This suggests the formation of a surface compound from Si and H at -0.5 v. Two processes are possible for H₂ evolution: (A) $Si + e_{val}^- + H^+ \rightarrow SiH$; $SiH + e^- + H^+ \rightarrow Si + H_2 \uparrow$. The second reaction is much retarded for p-type Si. (B) Hydrogen forms dipoles with outward-directed negative poles on the Si surface. With n-type Si, the negative charge of the surface is compensated by the positive charge of the surface barrier, and further hydrogen adsorption is restricted. With p-type Si, the positive pole of the dipole is a hole. As p-type dipoles do not reach into the body of the semiconductor the formation of

Card 2/3

Electrochemical evolution of...

S/076/62/036/005/006/013
B101/B110

further dipoles and further hydrogen adsorption is possible. There are
4 figures.

SUBMITTED: July 27, 1960

Card 3/3

X

YEFIMOV, Ye.A.; YERUSALIMCHIK, I.G.; SOKOLOVA, G.P. (Moscow)

State of the surface of anodically polarized silicon in hydro-
fluoric acid solutions. Zhur. fiz. khim. 36 no.6:1219-1221
Je'62 (MIRA 17:7)

h0044

S/076/62/036/008/005/011
B101/B144

24,7700

AUTHORS: Yefimov, Ye. A., and Yerusolimchik, I. G.

TITLE: Effect of the bichromate ion on the anodic dissolution of germanium

PERIODICAL: Zhurnal fizicheskoy khimii, v. 36, no. 8, 1962, 1791 - 1794

TEXT: Proceeding from observations made by F. Beck, H. Gerischer (Z. Electrochem., 63, 943, 1959), the behavior of p- and n-type Ge (resistivity 1.0 ohm.cm, diffusion length 0.5 mm) in 0.1 N H₂SO₄ was studied in the presence of 0.15 - 0.03 M K₂Cr₂O₇ at room temperature.

Results: (1) With p-type Ge, the potential of anodic dissolution increased in the presence of the bichromate by ~0.2 v for the whole range investigated (0 - 1.6 ma/cm²). (2) With n-type Ge, the potential of anodic dissolution increased whereas the saturation current dropped to nearly one-half. Exposure of the Ge electrode to light eliminated the bichromate effect. (3) On thin Ge electrodes with p-n junction a small bichromate effect with reverse bias and a reduction of the saturation

Card 1/2

Effect of the bichromate ion...

S/076/62/036/008/005/011
B101/B144

current with open p-n circuit were observed. Conclusion: The $\text{Cr}_2\text{O}_7^{2-}$ anion is adsorbed on the positively charged Ge surface. Since the valence electrons of the ion are drawn to the oxygen atoms, the Cr^{6+} center attracts electrons and repels holes. This inhibits the anodic dissolution and reduces the saturation current. On exposure to light, this effect is compensated by the intense generation of holes on the surface. There are 3 figures. J

SUBMITTED: December 8, 1961

Card 2/2

YERUSALIMCHIK 1.5
PHASE I BOOK EXPLOITATION

SOV/6448

Yefimov, Yevgeniy Aleksandrovich, and Josif Grigor'yevich Yerusalimehik

Elektrokhimiya germaniya i kremniya (Electrochemistry of Germanium and Silicon) Moscow, Goskhimizdat, 1963. 180 p. Errata slip inserted. 5000 copies printed.

Ed.: A. T. Kochnev; Tech. Ed.: V. V. Kogan.

PURPOSE: The book is intended for scientific workers, engineers, and technicians working in the semiconductor industry. It may also be useful to advanced students specializing both in semiconductor engineering and in electrochemistry.

COVERAGE: The book is a generalization of investigations carried out by Soviet and non-Soviet scientists in a new area of physical chemistry, the electrochemistry of semiconductors such as germanium and silicon. It offers a systematic outline of the structure of the electric double layer at the semiconductor-electrolyte interface and the kinetics of the anodic dissolution of germanium and

Card 1/8
✓

Electrochemistry of Germanium and Silicon

SOV/6448

silicon and provides data on other electrochemical reactions occurring on germanium and silicon electrodes. A special chapter has been devoted to a discussion of electrochemical operations performed in the production of semiconductor devices.. The authors thank Ye. N. Paleolog, Candidate of Chemical Sciences, for his valuable comments. Each chapter is accompanied by references.

TABLE OF CONTENTS:

Foreword	5
Ch. I. Structure of the Semiconductor-Electrolyte Interface	7
1. Surface properties of a semiconductor	7
2. Charged layers at semiconductor-electrolyte boundary	12
Bibliography	22

Card 2/7

YEFIMOV, Ye.A.; YERUSALIMCHIK, I.G. (Moscow)

Effect of electrical and structural inhomogeneities on the
process of anodic dissolution of germanium. Zhur. fiz. khim.
38 no.3:589-592 Mr '64. (MIRA 17:7)

ACCESSION NR: AP4033398

8/0016/64/038/003/0589/0592

AUTHOR: Yefimov, Ye. A. (Moscow); Yerusalimchik, I. G. (Moscow)

TITLE: Effects of electric and structural heterogeneities on the anodic dissolution of germanium.

SOURCE: Zhurnal fizicheskoy khimii, v. 38, no. 3, 1964, 589-592

TOPIC TAGS: germanium, anodic dissolution, polarization, anodic polarization, hole, electric heterogeneity, structural heterogeneity

ABSTRACT: The purpose of this investigation was to find the cause of the discrepancies between the theoretically calculated limiting current for the anodic dissolution of germanium and the much greater experimentally observed current. Since the ordinary single crystals of germanium are not strictly homogenous, specially grown crystals of n-germanium with $\rho = 3 \text{ ohm}\cdot\text{cm}$ and length of the order of 0.7 mm, containing a minimum amount of impurities and the density of dislocations of 50 disloc./cm² and also germanium with the same electric and physical parameters but having a density of dislocations $\sim 6 \cdot 10^4 \text{ disloc./cm}^2$ were used for this investigation. The anodic polarization curves were obtained by the potentio-

Card 1/4

ACCESSION NR: AP4033398

static method in 0.1 N H_2SO_4 (Fig. 1). The experimental results show that the increase of the limiting current during the anodic dissolution of germanium is associated with an additional generation of holes on some parts of the electrode surface. A higher concentration of Cu and Ni on such parts of the electrode may lead to the formation of the high resistance micro regions where the acceptor impurities compensate for the main part of donor impurities or it may lead to segregation of Cu and Ni into a separation phase, primarily at the places of disruption of the crystal lattice. In the areas of germanium on which the conductivity is close to the bulk conductivity, the limiting current due to holes is much greater than on bulk n-germanium. Upon increase of anodic polarization these zones may completely change the type of conductivity. The p-regions which occur at the n-germanium electrolyte interface will carry a large fraction of current, thus individual areas of the electrode surface dissolve more rapidly than others. A segregation of Cu and Ni in a separate phase in the germanium crystal may be produced due to break-through and local generation. This was verified by measuring the photoelectric potential as a function of the potential of germanium electrode in 0.1 N H_2SO_4 . "The authors express their gratitude to L. I. Kolesnik and Yu. A. Kontsevoy for their help and valuable suggestions during discussion of the results." Orig. art. has: 2 figures.

Card 2/4

ACCESSION NR: AP4033398

ASSOCIATION: None

SUBMITTED: 14Jan63

SUB CODE: IC

NO REF SOV: 005

ENCL: 01

OTHER: 004


Card 3/4

ACCESSION NR: AP4033398

ENCLOSURE: 01

Fig. 1. Anodic polarization curves obtained by the potentiostatic method in 0.1 N H_2SO_4 with n-germanium electrodes. Resistivity of germanium was 3 ohm·cm.

- 1) $L = 0.7$ mm, 50 disloc./cm²
- 2) $L = 0.7$ mm, $6 \cdot 10^4$ disloc./cm²
- 3) $L = 0.03$ mm, 10^7 disloc./cm²
- 4) $L = 0.03$ mm, (specimen 2 after removal of Cu & Ni impurities)



Card 4/4

ACCESSION NR: AP4033404

S/0076/64/038/003/0720/072 3

AUTHORS: Yefimov, Ye.A.; Yerusalinchik, I.G.; Gorgoraki, Ye.I.

TITLE: Reduction of persulfate ion at a germanium cathode

SOURCE: Zhurnal fizicheskoy khimii, v. 38, no. 3, 1964, 720-723

TOPIC TAGS: persulfate ion reduction, reduction, germanium cathode, n type germanium, p type germanium

ABSTRACT: Because of the contradictory data given in literature on the reduction of persulfate ion at germanium electrode, this reaction was studied by the potentiostatic polarization method and also via measurement of the photoelectric potential of the germanium electrode. This permitted determination of the magnitude of the curvature of the energy zone on the electrode surface. Electrodes from n- and p-type germanium with specific resistance of 1.5 ohm. λ_{om} and diffusion zone length of 0.7 mm were used. A series of experiments were made on a degenerated polycrystalline germanium which does not have semiconductor properties and also using electrodes with p-n transition. Polarization curves taken in 0.001 N $K_2S_2O_8$

Card 1/3

ACCOSSION NR: AP4033404

on n- and p-degenerated germanium show that under given conditions the reduction process does not depend on the type of electrode conductivity and that a limiting current of $\sim 0.35 \text{ ma/cm}^2$ is the normal specific current for persulfate ion diffusion to the electrode surface. The addition of an indifferent electrolyte to a 0.001 N $\text{K}_2\text{S}_2\text{O}_8$ solution decreases somewhat the inhibition of the electrochemical reaction. It was found that on increasing the concentration of the persulfate ion in the solution, the polarization curves for p- and n-germanium begin to differ and at $\varphi = -0.2$ to -0.1v the rate of reaction increases. With increase of the concentration of ammonium persulfate the photopotential increases and the value of the potential of flat zone is displaced toward the more positive potentials for the p- and n-type germanium electrodes. Since the polarization curves on n- and p-germanium corresponds to potentials -0.2 to -0.1v , it was concluded that in both cases the reaction is inhibited. On the basis of the lack of limiting current for the diffusion of electrons in the p-germanium it was assumed that electrons of the valence zone take part in the reduction or the rate of surface recombination at the electrolyte boundary is very great. Orig. art. has: 7 figures.

Card 2/3

ACCESSION NR: AP4033404

ASSOCIATION: None

SUBMITTED: 14Jan63

DATE ACQ: 15May64

ENCL: 00

SUB CODE: GC, GP

NR REF SOV: 003

OTHER: 002

Card

3/3

YEFIMOV, Ye.A.; YERUSALIMCHIK, I.G.; GORGORAKI, Ye.I.

Photoelectric potential of a germanium electrode as dependent
on the composition and concentration of electrolytes. Zhur.
fiz. khim. 38 no.5:1271-1273 My '64. (MIRA 18:12)

1. Submitted Jan. 14, 1963.

YEFIMOV, Ye.A.; YERUSALIMCHIK, I.G.; SOKOLOVA, G.P. (Moskva)

Electrochemical behavior of the silicon electrode in solutions
of oxidation agents. Zhur. fiz. khim. 38 no.9:2172-2175 S '64.
(MIRA 17:12)

YEFIMOV, Ye.A.; YERUSALIMCHIK, I.G. (Moscow)

Reduction of organic compounds on the platinum cathode. Zhur.
fiz. khim. 38 no.12:2868-2874 D '64.

(MIRA 18:2)

1. The first of these is the

1. The first of these is the

2. The second of these is the

3. The third of these is the

4. The fourth of these is the

5. The fifth of these is the

6. The sixth of these is the

7. The seventh of these is the

8. The eighth of these is the

9. The ninth of these is the

10. The tenth of these is the

11. The eleventh of these is the

12. The twelfth of these is the

13. The thirteenth of these is the

14. The fourteenth of these is the

15. The fifteenth of these is the

16. The sixteenth of these is the

17. The seventeenth of these is the

18. The eighteenth of these is the

19. The nineteenth of these is the

20. The twentieth of these is the

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466	467	468	469	470	471	472	473	474	475	476	477	478	479	480	481	482	483	484	485	486	487	488	489	490	491	492	493	494	495	496	497	498	499	500	501	502	503	504	505	506	507	508	509	510	511	512	513	514	515	516	517	518	519	520	521	522	523	52
--	---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	----

"APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962910018-4

SECRET

SECRET

APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962910018-4"

"APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962910018-4

APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962910018-4"

YEFIMOV, Ye.A.; YERUSALIMCHIK, I.G.

Electrochemical processes on an arsenic electrode. Elektr.
khimika 1 no.9:1133-1137 S '65. (MIRA 18:10)

YERUSALIMCHIK, Kh.I., prof., doktor med.nauk,

Involvement of the nervous system in nonpenetrating wounds of the
heart and pericardium. Voen.-med.zhur. no.9:26-29 S '59. (MIRA 13:1)

(HEART, wds. & inj.)
(NERVOUS SYSTEM, pathol.)

YERUSALIMCHIK, Kh.I. (Moskva)

Disturbance of cerebral circulation due to infarction of the
myocardium (cerebrocardiac syndrome). Kh.I.Erusalimchik.
Kaz. med. zhur. no.1:16-20 Ja-F'61 (MIRA 16:11)

*L

YERUSALIMCHIK, Kh.G., prof.

Difficulties in the diagnosis of compression of the spinal cord and its roots in tuberculous spondylitis; clinical and anatomical parallels. Sov.med. 23 no.6:55-62 Je '59. (MIRA 12:9)

1. Iz kliniki nervnykh bolezney (dir. - prof.I.N.Filimonov) II Moskovskogo meditsinskogo instituta imeni N.I.Pirogova.
(TUBERCULOSIS, SPINAL compl.)
(SPINAL CORD dis.)

YERUSALIMOV, M. Ye.

231T26

USSR/Electricity - Insulators
X Ray Tests

Oct 52

"A New Method for Prophylactic Tests of Equipment Insulators," Docent I. K. Fedchenko, Cand Tech Sci, M. Ye. Iyerusalimov

"Elektrichestvo" No 10, pp 45-49

Discusses a method for X-ray tests of mastic-filled equipment insulators which was developed in the High-Voltage Lab, Kiev Polytech Inst. Establishes the optimum conditions for penetration of the complex structure porcelain-mastic-jetmax

231T26

[The latter is a laminated plastic made from paper and synthetic bakelite resin]. Submitted 22 Nov 51.

231T26

YERUSHALIMOV, M. Ye.

YERUSHALIMOV, M. Ye.; FEDCHENKO, I. K.

Electric Lines

Controlling the condition of high voltage lead-ins, Elek. sta. 24, No. 1, 1953.

9. Monthly List of Russian Accessions, Library of Congress, May 1953. Unclassified.

YERUSALIMOV, M. Ye.

AID P - 2424

Subject : USSR/Electricity

Card 1/1 Pub. 26 - 23/33

Authors : Ierusalimov, M. E., Kand. Tech. Sci., and Fedchenko, I.K.,
Kand. Tech. Sci.

Title : On co-ordination of transmission line insulation

Periodical : Elek sta 5, 54-55, My 1955

Abstract : The authors criticize M. M. Nekrasov's article appearing in the No 12, 1953 issue of this journal on co-ordination of insulation of transmission lines. They maintain that Nekrasov's suggestions if applied would aggravate the situation. The article is accompanied by an answer from Nekrasov.

Institution: None

Submitted : No date

EXHAUSTION, M. Ya.

FEDCHENKO, I.K., doktor tekhnicheskikh nauk, professor; IYERUSALIMOV, M.Ya.,
kandidat tekhnicheskikh nauk, dotsent; ANDRIYASHEV, K.Ya., inzhener;
MARKEVICH, V.P., inzhener.

X ray examination of high-voltage insulators. Elektrichestvo no.8:
78-79 Ag '56. (MLRA 9:10)

1.Kiyevskiy ordena Lenina politekhnicheskiiy institut (for Fedchenko,
Iyerusalimov. 2.Kiyevenergo (for Andriyashov, Markevich).
(Electric insulators and insulation) (X rays--Industrial applications)

8(3)

SOV/112-59-3-4366

Translation from: Referativnyy zhurnal. Elektrotehnika, 1959, Nr 3,
pp 12-13 (USSR)

AUTHOR: Fedchenko, I. K., and Iyerusalimov, M. Ye.

TITLE: Investigation Into the Cause of Deterioration of High-Voltage Porcelain
Insulators on the Oil Circuit Breakers (issledovaniye prichin razrusheniya
farforovykh izolyatorov vysokogo napryazheniya na maslyanykh vyklyuchatelyakh)

PERIODICAL: Izv. Kiyevsk. politekhn. in-ta, 1957, Vol 22, pp 380-393

ABSTRACT: At ~~Elyavenergo~~ (Kiyev Power System) substations, cracks in bushing
porcelain insulators of VMD-35 oil circuit breakers were detected. The
cracks started at the support flange and spread upward over the porcelain. By
using the x-ray flaw-detection method, it was found that the mortar used for
fastening the porcelain penetrated inside the bushing and filled a part of the
hollow space between the "getinaks" core and the porcelain wall. The cement
belt extended over the flange by 2-12 cm; in some bushing insulators, the

Card 1/3

8(3)

SOV/112-59-3-4366

Investigation Into the Cause of Deterioration of High-Voltage Porcelain Insulators . . .

cement layer reached the level of the first insulator rib. The cement belt inside the insulator crowded out the insulating compound and caused cracking of the porcelain because the temperature coefficient of expansion for cement is considerably higher than that for porcelain. Computations have shown that with the maximum permissible temperature $+80^{\circ}\text{C}$ of the current-carrying tube and the temperature $+35^{\circ}\text{C}$ of the ambient air, the tensile stress on the internal surface of the porcelain may go as high as 234 kg/cm^2 , which considerably exceeds the permissible value. Bushing tests on other oil circuit breakers made by conventional methods (measurement of $\text{tg}\delta$, high-voltage test) did not detect the cement-belt faulty insulators. Roentgenoscopy of 32 insulators helped to detect 20 faulty insulators with cement between the porcelain bushing and the "getinaks" insulation. Insulator roentgenoscopy was made by means of a portable x-ray outfit in a housing $70 \times 30 \times 30 \text{ cm}$, weighing 40 kg. The insulator roentgenoscopy was performed with a voltage 60 kv on the x-ray tube.

Card 2/3

8(3)

SOV/112-59-3-4366

Investigation Into the Cause of Deterioration of High-Voltage Porcelain Insulators . . .

the maximum tube current 5 ma, at a focal length of 30 cm, and with an exposition of 3 minutes. The x-ray-outfit transformer was supplied from a 220 v line. The roentgenoscopy of 6 bushings of one circuit breaker took less than one hour. Bibliography: 8 items.

N.V.N.

Card 3/3

IYERUSALIMOV, M.Ye.; RYTSLIN, A.M.; MORGIN, P.I.

Radiomaterialogy of high-voltage insulation used in power systems.
Izv. KPI 22:394-396 '57. (MIRA 11:3)
(Electric insulators and insulation--Testing)
(Radiology, Industrial)

IYERUSALIMOV, M.Ye.

Shifting electric power-transmission lines of 10-110 kv.
on wooden supports to an increased voltage rating. Izv. KPI
26:441-453 '57.

(NIRA 11:6)

1. Kafedra tekhniki vysokikh napryazheniy Kiyevskogo politekhnicheskogo instituta.

(Electric lines)

FEDCHENKO, I.K., doktor tekhn. nauk prof.; IYERUSALIMOV, M.Ye., kand. tekhn. nauk dots.

Characteristics of the electric strength of large spark gaps
at constant high voltage. Izv. vys. ucheb. zav.; energ. 2 no.7:
33-40 JI '59. (MIRA 13:1)

1.Kiyevskiy ordena Lenina politekhnicheskoy institut.
(Electric spark)

IYERUSALIMOV, M.Ye., kand. tekhn. nauk

Improvement of the checking operation of line suspension
insulators before their installation. Energ. i elektrotekh.
prom. no.2:48-50 Ap-Je '63. (MIRA 16:7)

1. Kiyevskiy politekhnicheskii institut.
(Electric lines—Overhead)

LITVINOV, L.N., kand.tekhn.nauk; SOLOV'YEV, A.I., inzh.; ^vIERUSALIMOV, Ye.P.,
inzh.

Driving piles without a pile driver using the UR-1250 diesel
hammer. Transp. stroi. 13 no.2:17-18 F '63. (MIRA 16:3)
(Piling (Civil engineering))

PONOMARENKO, A.V., ispolnyayushchiy obyazannosti dotsenta; VINOGRADOV, P.V.;
starshiy nauchnyy sotrudnik; MIKHAYLOV, K.G., agronom-entomolog;
IYERUSALIMSKAYA, K.P., studentka

Controlling soil pests in checkrowed corn fields. Zashch. rast.
ot vred. i bol. 5 no.4:24-27 Ap '60. (MIRA 13:9)

1. Rostovskiy universitet (for Ponomarenko, Iyerusalimskaya).
2. Zernogradskaya selektsionnaya stantsiya (for Vinogradov).
3. Sal'skiy nablyudatel'nyy punkt (for Mikhaylov).
(Corn (Maize))---Diseases and pests)

BEZRUK, V.M.; KOTYLEV, Yu.L.; GROT, A.I.; ZNAMENSKIY, A.I.; YERUSALIMSKAYA,
M.F.; GERBURT-GEYBOVICH, A.V., redaktor; KOVALIKHINA, N.F., tekhnicheskii redaktor

[Building roads on saline soils and shifting sands] Stroitel'stvo
dorog na zasolennykh gruntakh i podvizhnykh peskakh. Moskva,
Avtotransizdat, 1953. 202 p. (MLRA 7:8)

1. Moscow. Dorozhnyy nauchno-issledovatel'skiy institut.
(Road construction)

YERUSALIMSKAYA, M. F.

USSR/Agriculture - Soil testing instruments

Card 1/1 Pub. 123 - 8/17

Authors : Yerusalsinskaya, M. F.

Title : The simplest salt measuring device

Periodical : Vest. AN Kaz. SSR, 11, 61-66, Nov 1954

Abstract : A simple device for measuring the solinity of soil quickly (5-6 minutes) is described. The device operates on the electrolysis principle. Three USSR references (1932-1953). Graphs; table; illustration.

Institution :

Presented by : Active Member of the Acad. of Sc. of the Kaz. SSR K. I. Goryaev

YERUSALIMSKAYA, M.F., inzhener

Quality of soil packing in earth embankments. Avt.dor.18 no.5:
11-12 S'55. (MLRA 9:1)

(Embankments)

YERUSALIMSKAYA, M.F.

IYERUSALIMSKAYA, M.F.

Effect of the degree of salinity on the hydrating capacity of soils.
Izv. AN Kazakh. SSSR. Ser. gor. dela, met., stroi. i stroimat. no. 10:81-
90 '56. (MIRA 10:1)
(Alkali lands) (Hydration) (Kazakhstan--Road construction)

1748
KHARCHENKO, V.A.; IYERUSALIMSKAYA, M.P.

Northern Kazakhstan soils as a road building material. Inv. AN
Kazakh. SSR. Ser. gor. dela, met., stroi. i stroimat. no.3:114-
136 '57. (MIRA 10:11)
(Kazakhstan--Solonets soils) (Road construction)

IYERUSALIMSKAYA, M.F.

Soil-cement road beds to be constructed in northern Kazakhstan
regions. Trudy Inst. stroi. i stroimat. AN Kazakh. SSR 2:72-77
'59. (MIRA 12:10)

(Kazakhstan--Roads, Soil-cement)

IYERUSALIMSKAYA, M.F.

Road construction in Kazakhstan. Avt. dor. 24 no.7:7-8 J1 '61.
(MIRA 14:7)

(Kazakhstan--Road construction)

YERUSALIMSKAYA, S.M.

Method of counting thrombocytes. Vrach. delo no. 9:132-133
s:63. (MIRA 16:10)

I. Gorodskaya klinicheskaya bol'nitsa Sovetskogo rayona g.
Kiyeva.

(BLOOD PLATELETS)

"APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962910018-4

APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962910018-4"

YERUSALIMSKIY, A.L.

Etiology and pathogenesis of malignant tumors (hypothesis of
"local crisis of reproduction"). Vrach. delo no.2:101-104 F '61.
(MIRA 14:3)

(CANCER)

YERUSALIMSKIY, A.L. [deceased]; KAZ'MIN, S.D.

Effect of some antitumoral preparations on the kinetics of
chain reactions. Ukr.khim.zhur. 31 no.5:521-524 '65.
(MIRA 18:12)

1. Submitted October 22, 1964.

YERUSALIMSKIY, A. M.

Motorcycles. Leningrad, Gos. nauchno-tekhn. izd-vo mashinostroit. lit-ry, 1945.
295 p: (55-16281)

TL440.I3

●YERUSALIMSKIY, A.M.; IVANOV, A.A., inzhener, otvetstvennyy redaktor;
POLYAKOV, V.S., kandidat tekhnicheskikh nauk, glavnyy redaktor;
TISHKOVA, M.V., tekhnicheskiy redaktor

[Theory, construction and calculations for motorcycles] Teoriia,
konstruktsiia i raschet mototsikla. Izd. 2-e, ispr. 1 dop. Moskva,
Gos. nauchno-tekhn. izd-vo mashinostroit. lit-ry, 1947. 414 p.
(Motorcycle industry) (MLRA 8:2)

YERUSALIMSKIY, A. M.

Graficheskaya gramota rabochego-izobretatelya. Leningrad, Mashgiz, 1950.
63 p. diags.

Graphic record of a worker and inventor.

DLC: T339.I3

SO: Manufacturing and Mechanical Engineering in the Soviet Union, Library
of Congress, 1953.

ARANOVSKIY, M.G.; ORLENKO, N.I.; SHUKIN, L.S.; IYERUSALIMSKIY, A.M., dotsent,
redaktor.

[Drafting in machine construction] Cherteshnoe khoziaistvo v mashinostroenii.
Leningrad, Nauchno-tekhn. izd-vo mashinostroit. lit-ry [Leningradskoe otd-nie]
1953. 103 p. (MIRA 6:10)

(Machinery--Drawing) (Drawing-room practice)

Y
~~SYRUSALIMSKIY, A.M.~~; TSVETKOV, A.T., redaktor; DAVRILOV, S.S., tekhnicheskii redaktor

[Descriptive geometry] Nachertatel'naya geometriia. Moskva, Gos.
izd-vo tekhniko-teoret. lit-ry, 1954. 304 p. (MLRA 8:4)
(Geometry, Descriptive)

YERUSALIMSKIY, A.M.

FEDORENKO, Viktor Alekseyevich; SHOSHIN, Aleksandr Ivanovich; IYERUSALIMSKIY,
-A.M., professor, redaktor; GOFMAN, Ye.E., redaktor isdatel'stva;
SOKOLOVA, L.V., tekhnicheskiiy redaktor

[Machinery drawing handbook] Spravochnik po mashinostroitel'nomu
chercheniyu. Pod red. A.M.Ierusalimskogo. Izd. 4-oe, ispr. i dop.
Moskva, Gos. nauchno-tekhn. izd-vo mashinostroit. lit-ry, 1956.
208 p. (MLRA 10:4)

(Machinery--Drawing)

YERUSALIMSKIY, A.M.

VYSOTSKAYA, N.H.; YERUSALIMSKIY, A.M.; HEVEL'SON, R.A.; FEDORENKO, V.A.;
GOFMAN, Ye.K., redaktor; FUGACHEV, A.A., inzhener, retsenzent;
POL'SKAYA, R.G., tekhnicheskiiy redaktor

[Technical projections for articles made of sheet metal] Tekhnicheskies razvertki izdelii iz listovogo materiala. Pod obshchei red. A.M.Ierusalinskogo. Moskva, Gos.nauchno-tekhn.izd-vo mashinostroitel'noi lit-ry, 1955. 230 p. (MLRA 9:1)
(Sheet-metal work)

IYERUSALIMSKIY, A.P.

Information on a conference in Novosibirsk on the control of
diseases with a natural focus. Zhur. mikrobiol. epid. i immun.
32 no.5:157-158 My '61. (MIRA 14:6)
(COMMUNICABLE DISEASES CONGRESSES)

YERUSALIMSKIY, A.S., doktor ist. nauk, otv. red.; AYZIN, B.A.,
kand. ist. nauk, red.; GALKIN, I.S., doktor ist. nauk, red.;
GOROSHKOVA, G.N., kand. ist. nauk, red.; SMIRIN, M.M., doktor
ist. nauk, red.; TARTAKOVSKIY, B.G., red. izd-va; KOVICHKOVA,
N.D., tekhn. red.

[German labor movement in the modern period] Germanskoe rabo-
chee dvizhenie v novoe vremia; sbornik statei i materialov.
Moskva, Izd-vo Akad. nauk SSSR, 1962. 405 p. (MIRA 15:10)

1. Akademiya nauk SSSR. Institut istorii.
(Germany--Labor and laboring classes)

YERUSALIMSKIY, B.

Machine for over-all mechanization of housing construction.
NT0 2 no.9:28-29 S '60. (MIRA 13:9)

1. Zamestitel' nachal'nika Spetsial'nogo konstruktorsko-tekhnicheskogo byuro; chlen soveta nauchno-tekhnicheskogo obshchestva zavoda No.5 Glavtonnel'metrostroya Ministerstva transportnogo stroitel'stva.

(Building machinery)

YERUSALIMSKIY, B. G., SPIVAK, P. Yo.

"Measuring the Neutron Multiplication Factor for Thermal Fission of Uranium and Plutonium," a paper presented at the Atoms for Peace Conference, Geneva, Switzerland, 1955

124 of substitution company in requirements specified
for the oil industry. H. G. Krommeltail and D. E. Hen-
richs. *Chem. Abstr.* 1985, 115, 181-93 (Pub.
Energy, Science Div. Tech. Rep. 1985, 181-93 (Pub.
1985) (Engl. translation).--See C.A. 90, 654f. H. M. R.


"APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962910018-4

APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962910018-4"

"APPROVED FOR RELEASE: 03/15/2001 CIA-RDP86-00513R001962910018-4



APPROVED FOR RELEASE: 03/15/2001 CIA-RDP86-00513R001962910018-4"

YERUSALIMSKIY, B. L.

USSR/Chemistry - Reaction processes

Card 1/1 Pub. 151 - 14/37

Authors : Dolgoplosk, B. A.; Yerusalimskiy, B. L.; Krol', V. A.; and Romanov, L. M.

Title : Reaction of free radical in solutions. Part 2.- Relative activity of free radicals during reaction with isopropyl benzene, cyclohexene and polymers

Periodical : Zhur. ob. khim. 24/10, 1775-1782, Oct. 1954

Abstract : Data regarding the relative activity of numerous free radicals, which are distinguished by their entirely different reactivity characteristics, were obtained by studying the products of thermal decomposition of diazamine compounds in solution. A series of free radical activities was established in the reaction of separation of the H-atom from various compounds. The reason why allyl, crotyl, benzyl and tertiary-butyl free radicals do not react with isopropyl benzene, is discussed. The effect of low-active radicals (allyl, benzyl and anil) on the structure-formation of natural rubber, is explained. Twelve references: 4-USA; 5-German; 2-USSR and 1-English (1895-1953). Tables.

Institution : Acad. of Sc. USSR, All-Union Scientific Research Institute of Synthetic Rubber and Institute of High Molecular Compounds

Submitted : March 2, 1954

YERUSALIMSKIY, B. W.

The inhibition mechanism of radical chain reaction
in the presence of a radical scavenger is discussed.
The results of the experiments are presented.
The mechanism of the inhibition of the radical chain reaction
by a radical scavenger is discussed.
The results of the experiments are presented.

1/1

YER. SABINOVIT BL.

✓ Reactivity of free radicals and the role of the polymer filler.
B. A. Belogorsh, B. I. Gerasimov, V. A. Kaul, and L. I. Romanov. *Popravly Rubr. Kautsch. Latexes i Reaktivnaya Spetsializatsiya. Akad. Nauk S.S.S.R. Otdel. Khim. Nauk* 1955, 810-20. The yields of RH and RL in radical reactions of RN:NNHPh in *iso*-PrPh or cyclohexene indicate the order of descending radical reactivity at Me, Et, C₆H₅, Et, Pr, Bu, Me₂CCN, *iso*-Pr, *neo*-Bu, allyl, MeCH₂-CHCH₃, PhCH₂, and Me₂C. A similar series is obtained in the yield of final products in reactions of these radicals with benzene solns. of rubber, or in thermal treatment of rubber in vulcanization. The following yields of triidim-soln vulcanizate were obtained with: diisopentylazene 70-90%; diazaminobenzene 60-85%; methylphenyltriazene 13-44%; isopropylbenzene hydroperoxide 18-23%; Et₂O₂ 11-60%. Benzylphenyltriazene, allylphenyltriazene, and butylphenyltriazene gave 0%. G. M. Loskutov

YERUSALINSKIY, B.I., DOLGOPLASK, B. A., MELOVSKAYA, E. B., and KOVURENKO, A. P.

"Free radicals and unsaturated compounds in polymerization," a paper
presented at the 9th Congress on the Chemistry and Physics of High Polymers,
28 Jan-2 Feb 57, Moscow, Polymer Research Inst.

B-3,084,395

"APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962910018-4

APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962910018-4"

"APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962910018-4

APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962910018-4"

"APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962910018-4

APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962910018-4"

491

AUTHORS: Yerusalimskiy, B. L.; Dolgoplosk, B. A.; and Kavunenko, A. P.

TITLE: Reactions of Free Radicals in Solutions. Part 9. Dimethyldiphenyltetrazene and tetramethyltetrazene as Sources of Free Radicals with Reaction Center on a Nitrogen Atom (Reaktsii svobodnykh radikalov v rastvorakh. IX. Dimetildifeniltetrazen i tetrametiltetrazen kak istochniki svobodnykh radikalov s reaktsionnym tsentrom na atome azota)

PERIODICAL: Zhurnal Obshchey Khimii, 1957, Vol. 27, No. 1, pp. 267-270 (U.S.S.R.)

ABSTRACT: This report presents for the first time data obtained during the study of free radicals with the reaction center on a nitrogen atom. It was established that RNR type radicals are highly active in reactions leading to the separation of the H-atom from the carbon molecule, additional reactions to the vinyl double bond and are capable of exciting the polymerization process. Using dimethyldiphenyltetrazene as an example, the authors established the monomolecular nature of tetrazene decomposition during heating in a solution. The activation energy of this reaction was calculated from kinetic data by the Arrhenius equation and fixed at 33 kcal/mol. The ability of free radicals with a reaction center

Card 1/3

491

Reactions of Free Radicals in Solutions. Part 9

on a nitrogen atom to excite the polymerization process was verified on a styrene-dimethyldiphenyltetrazene system (See Fig.2). The ability of alkylnitrous radicals to react with saturated and unsaturated hydrocarbons was determined by studying the products of thermal decomposition of tetramethyltetrazene in isopropylbenzene and in alpha-methylstyrene. Tetramethyltetrazene appears to be of high thermal stability; at 130° it decomposes in the solution at a slow rate. The constant of the decomposition rate at 145° in isopropylbenzene at a 0.7 mol% concentration is $0.56 \cdot 10^{-4} \text{ sec}^{-1}$. The complete absence of dimethylamine in the products of tetramethyltetrazene decomposition in alpha-methylstyrene indicates that the free dimethylnitrous radicals do not become disproportionated.

One table, 2 graphs. There are seven references, of which 1 is Slavic.

Card 2/3

Reactions of Free Radicals in Solutions. Part 9

492

ASSOCIATION: Academy of Sciences USSR, Institute of High Molecular Compounds
(Institut Vysokomolekulyarnykh Soyedineniy Akademii Nauk SSSR)

PRESENTED BY:

SUBMITTED: February 17, 1956

AVAILABLE:

Card 3/3

"APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962910018-4

APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962910018-4"

"APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962910018-4

APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962910018-4"

YERUSALIMSKIY, B.L.

AUTHORS	Dolgoplosk, B.A. Romanov L.M., Yerusalimskiy, B.L. and Bogomol'nyy, V.Ya.	20-4-27/60
TITLE	The Use of Catalysts based on Magnesium Halogen Alkyls and Titanium Tetrachloride in the Production of Amorphous and Crystalline Polymers from α -Olefins and Dienes (Katalizatory na osnove magniygalogenalkilov i chetyrekh-khloristogo titana dlya sinteza amorfnykh i kristalliches-kikh polimerov iz α -olefinov i dienov.)	
PERIODICAL	Doklady Akademii NaukSSSR, 1957, Vol. 115, Nr 4, pp. 731-733 (USSR)	
ABSTRACT	The production of polymers from aliphatic compounds of the ethylene series on the basis of polymerization by free radicals is only possible for the first representative of this series, namely ethylene. The interaction of the free radicals with the ethylene homologues leads to the formation of low-molecular products due to the rupture reactions in the first stages of the process. Ziegler and his collaborators used the reaction between organo-aluminum compounds and titanium tetrachloride for initiating the ethylene polymerization. Further investigations in this field furnished the possibilities of initiating the olefin polymerization under formation	
CARD 1/ 4		

20-4-27/60

The Use of Catalysts based on Magnesium Halogen Alkyls and Titanium Tetrachloride in the Production of Amorphous and Crystalline Polymers from α -Olefins and Dienes

seide. The thus obtained polyethylene possesses $\eta = 2,55$ melting point $130 - 138^{\circ}\text{C}$, ultimate strength 335 Kg/cm^2 and relative extension 730% . These indices are analogous to those of polyethylene which is obtained by means of the ordinary Ziegler catalyst. Of great interest are data which were obtained by the authors from the polymerization of propylene. It was proved by the authors that on this occasion in the presence of magnesium chloroethyl and tetrachlorotitanium about the same quantities of an amorphous and a crystalline (isotactic) polymer-form develop. At the same time a fraction was isolated which is insoluble in boiling hexane and whose crystallizability was spectroscopically and X-ray structurally proved. In the case of the polymerization of isoprene various polymer forms were also isolated. The simultaneous production of cis-1,4-polyisoprene and trans-1,4-polyisoprene or of the amorphous and the isotactic polyisoprene directly prove that in the system coexist various catalytic

CARD 3/4

YERUSALIMSKIY, B. L.

AUTHORS: Dolgoplosk, B. A., Yerusalinskiy, B. L., 62-53-4-13/32
Tinyakova, Ye. I.

TITLE: Generation of Free Radicals in Solutions and Their
Reactions in Model Systems (Generirovaniye svobodnykh
radikalov v rastvorakh i ikh reaktsii v model'nykh
sistemakh). Report of the Conference on Chemical
Sciences of the AS USSR on October 30, 1957 (Doklad
na sessii otdeleniya khimicheskikh nauk Akademii nauk
SSSR, 30 oktyabrya 1957)

PERIODICAL: Izvestiya Akademii Nauk SSSR, Otdeleniye Khimicheskikh
Nauk, 1958, Nr 4, pp. 469-481 (USSR)

ABSTRACT: The present paper gives the final results of the work
of the authors with - in the field of oxidation - and
reduction initiation of radical processes and the in-
vestigation of a number of reactions of free radicals.
Corresponding to their action the oxidation and reduc-
tion systems are given in 3 groups (Ref. 1): To the
first kind belong systems in which the reaction takes
place between the reducing agent and the oxidizing agent

Card 1/3

02-504-17/52

Generation of Free Radicals in Solutions and Their Reactions in
Model Systems. **Report** of the Conference on Chemical Sciences
of the AS USSR on October 30, 1957

by forming a radical (see formulae 1,2,3). The detailed description of the first type (system with peroxides) follows. Also systems in which also metal salts with varying valence take part (as oxidizing agents) also belong here. The systems of the second kind are of interest in theoretical and practical respects (second type). Among them is also a system which acts with hydroquinone taking part. This system was utilized industrially (initiation of polymerizations in emulsions). There is still a number of other systems in which the reactions take part between the oxidizing agent and the reducing agent by formation of 2 radicals. Those systems belong to the third kind which have a participation of the diazoamino compounds. 2. Systems with participation of ethylenediamine and polyethylenepolyamine. 3. Systems with participation of sulfur and oxygen (as oxidizing agents). After classification of the systems according to their mechanisms the report deals with the different reactions of alkyl- and heteroradicals with various mo=

Card 2/3

62-58-4-13/32

Generation of Free Radicals in Solutions and Their Reactions in Model Systems. **Report** of the Conference on Chemical Sciences of the AS USSR on October 30, 1957

monomers and polymers on which occasion a break of the bonds C-H, C=C, C - C and S - S is formed. Finally the part played by the cell in the heat stability of polymers was investigated. There are 2 tables and 44 references, 40 of which are Soviet.

ASSOCIATION: Institut vysokomolekulyarnykh soyedineniy Akademii nauk SSSR (Institute for High-Molecular Compounds, AS USSR)

SUBMITTED: December 23, 1957

AVAILABLE: Library of Congress

1. Chemical conference--Report 2. Free radicals--Solutions--Reactions 3. Free radicals--Solutions--Generation

Card 3/3

AUTHORS: Milovskaya, Ye. B., Yerusalimskiy, B.L., SOV/20-120-2-31/63
Dolgoplosk, B. A.

TITLE: The Reactions of Free Radicals in Solutions (Reaktsii
svobodnykh radikalov v rastvorakh) The Interaction of Free
Radicals With Internal and External Double Bonds (Vzaimodeyst-
viye svobodnykh radikalov s vnutrennimi i vneshnimi dvoynymi
svyazyami)

PERIODICAL: Doklady Akademii nauk SSSR, 1958, Vol. 120, Nr 2, pp. 336-338
(USSR)

ABSTRACT: This paper discusses data which characterize the relative ac-
tivity of the internal and external double bonds in the re-
action of interaction with free radicals. The first part of
this paper discusses the interaction of the free methyl radical
with hydrocarbons and polymers. The intensity of this inter-
action was estimated indirectly by comparing the decrease of
methane (produced according to the reaction $R + LH \rightarrow RH + L$),
with the yield of CH_4 when the process takes place in a sat-
urated hydrocarbon. The internal double bond is by far less
capable for the addition of free radicals than a double bond
of the vinyl type. The investigation of the interaction of free
methyl radicals with polymers led to the same conclusion. In
swollen polystyrene containing 10% isopropylbenzene the methane

Card 1/3

The Reactions of Free Radicals in Solutions.
The Interaction of Free Radicals With Internal and External Double Bonds

SOV/20-120-2-31/63

yield amounts to only 9% of the theoretical value, whereas the yield in pure isopropylbenzene amounts to 49%. Transition to very viscous media leads to a decrease of the rôle of effective radical reactions. The second part of this paper deals with the interaction of the free dimethylcyanomethyl radical with 2-butene. The addition of this radical to an internal double bond is realized only to a very insignificant extent. Under the conditions of the experiments discussed in this paper these radicals have a tendency to recombine. Finally the experimental part is discussed. The authors describe the decomposition of methyl-phenyltriazene in several media and also the products of the decomposition of azoisobutyric acid in 2-butene. There are 2 tables and 4 references, 2 of which are Soviet.

ASSOCIATION: Institut vysokomolekulyarnykh soedineniy Akademii nauk SSSR
(Institute of High-Molecular Compounds, AS USSR)

PRESENTED: December 24, 1957, by B. A. Kazanskiy, Member, Academy of
Sciences, USSR

Card 2/3

The Reactions of Free Radicals in Solution.
The Interaction of Free Radicals With Internal and External Double Bonds

SOV/20-120-2-31/63

SUBMITTED: June 20, 1957

1. Free radicals--Chemical reactions 2. Methyl radicals--Chemical
reactions 3. Polymers--Chemical reactions 4. Hydrocarbons
--Chemical reactions

Card 3/3

SOV/20-120-4-26/67

AUTHORS: Dolgoplosk, B. A., Yernsalimskiy, B. L., Milovskaya, Ye. B.,
Belonovskaya, G. P.

TITLE: The Cell Effect and the Thermal Stability of Polymers
(Effekt kletki i termostabil'nost' polimerov)

PERIODICAL: Doklady Akademii nauk SSSR, 1958, Vol. 120, Nr 4, pp. 783-785
(USSR)

ABSTRACT: A great number of organic substances are known whose thermal stability is much weaker in the solution or melt than in the solid state. The most typical examples are compounds with unstable bindings (peroxides, azo- and diazo-compounds) which begin to decompose only at their melting temperature. At the same time they decompose much quicker in solutions and at a much lower temperature (Table 1). According to the authors' opinion the following experimental results render it possible to relate the mentioned phenomenon to a rapid increase of the cell effect (= reaction of the primary recombination of the free radicals) in viscous and solid media. As was proved already earlier the thermal decomposition of methyl-phenyl triazene in a medium of hydrocarbons leads

Card 1/4

SOV/20-120-4-26/67

The Cell Effect and the Thermal Stability of Polymers

to the formation of methane and methyl-aniline (Ref 5). It is most probable that the latter forms as a result of recombination of the radicals which are released at the moment of decomposition in the "cell". The authors proved that in the case of decomposition of methyl-phenyl triazene in systems of hydrocarbon polymers the methane yield decreases with increasing viscosity of the medium. At the same time it was proved that the yield of the product of primary recombination, namely of the methylaniline increases (Table 2). The above mentioned data give evidence as to a considerable influence of the viscosity of the medium on the efficiency of interaction in the cell. The results obtained render possible the discussion of a possible influence of the state of aggregation on the thermal stability of those substances that contain unstable bindings (Table 1) as well as of the polymers that have a high fusing temperature. The difference in behavior of such compounds in solid state and in solution (or melt) may be explained by means of the particularly important part played by the cell effect in solid state. Polyparaxylylene decomposes only after having been melted (at 425°). In the solution this is the case already at 302°. These polymers

Card 2/4

NOV/20-120-4-26/61

The Cell Effect and the Thermal Stability of Polymers

are apparently "overheated"; only after surpassing the temperature of vitrification they undergo a destructive decomposition when the viscosity of the system decreases considerably. Hence we may conclude that the thermal stability of polymers with a high melting temperature displays abrupt jumps in connection with the transition from solid state into an elastic one and from the elastic state into the solution. From the above mentioned it may be concluded that the task of increasing the thermal stability of carbon atom chain polymers in vitrified state consists above all in increasing their melting temperature. A high thermal stability of rubber-like polymers can apparently only be reached by the stability of the skeleton bindings of the main chain. There are 3 tables and 8 references, 2 of which are Soviet.

ASSOCIATION: Institut vysokomolekulyarnykh soyedineniy Akademii nauk SSSR
(Institute of High-Molecular Compounds AS USSR)

PRESENTED: January 6, 1958, by V. A. Kargin, Member, Academy of Sciences,
Card 3/4 USSR

The Cell Effect and the Thermal Stability of Polymers

SOV/20-120-4-26/67

SUBMITTED: January 4, 1958

1. Polymers--Thermodynamic properties
2. Polymers--Decomposition
3. Polymers--Molecular structure
4. Free radical--Chemical effects

Card 4/4

YERUSALIMSKIY, B. L.

820111
S/062/60/000/02/08/012
B003/B066

5.3200

AUTHORS:

Dolgoplosk, B. A., Yerusalimskiy, B. L., Kuren'gina, T. N.,
Tinyakova, Ye. I.

TITLE:

Reactions of Free Radicals in Solutions. 15th Report.
Destruction Mechanism of Polymers by Free Radicals

PERIODICAL: Izvestiya Akademii nauk SSSR. Otdeleniye khimicheskikh nauk,
1960, No. 2, pp. 311 - 316

TEXT: The authors investigated the destruction of polyisobutylene dis-
solved in ethyl benzene under the action of disulfides, benzoyl peroxide,
isopropyl benzene-hydroperoxide, triazenes, dimethyl-diphenyl-tetrazene,
iron- and cobalt naphthenate. The destructive effect of the individual
agents may be seen from the diagrams in Figs. 1, 2, and 3. The following
conclusions may be drawn from the investigations and pertinent papers by
other authors: The destructive effect is most intense in such free
radicals as are especially active in the reaction of H-separation. The
destruction takes place in such a manner that first a H-atom is separated
from the polymer chain and, secondly, the C-C bonds of the polymer radical

Card 1/2

Reactions of Free Radicals in Solutions.

15th Report. Destruction Mechanism of Polymers
by Free Radicals

82944
S/062/60/000/02/08/012
B003/B066

thus formed are spontaneously freed. The authors mention a paper by
L. M. Romanov (Ref. 2). There are 3 figures and 14 references: 8 Soviet,
4 American and British, and 2 German.

ASSOCIATION: Institut vysokomolekulyarnykh soyedineniy Akademii nauk
SSSR (Institute of High-molecular Compounds of the Academy
of Sciences USSR)

SUBMITTED: June 19, 1958

Card 2/2

VAN FO SUN [Wang-Fo-sung]; DOLGOPIOSK, B.A.; YERUSALIMSKIY, B.L.

Polymerization of isoprene induced by organomagnesium compounds.
Vysokom. soed. 2 no.4:541-545 Ap '60. (MIRA 13:11)

1. Institut vysokomolekulyarnykh soyedineniy AN SSSR.
(Isoprene) (Magnesium organic compounds)
(Polymerization)

S/062/60/000/03/04/007
B008/B006

AUTHORS: Van Fo-sun, Dolgoplosk, B. A., Yarusalinskiy, B. L.
TITLE: Reactions of Organo-metallic Compounds With Salts of
Heavy Metals. 1. Interaction of Ethyl Magnesium Bromide
With Halides of Titanium and Cobalt
PERIODICAL: Izvestiya Akademii nauk SSSR. Otdeleniye khimicheskikh
nauk, 1960, No. 3, pp. 469-473

TEXT: An investigation of the interaction of ethyl magnesium bromide with titanium trichloride and titanium tetrachloride in the presence of acceptors for free radicals was carried out. As such, nitrogen oxide, styrene, and α -methyl styrene were used. The experiments were carried out in the apparatus schematically shown in Fig. 1. Data on the interaction of magnesium ethyl bromide and titanium tetrachloride at 20°C in the presence of the above acceptors are given in Table 1. Data on the interaction of ethyl magnesium bromide and titanium trichloride at 100°C in the presence of nitrogen oxide and α -methyl styrene are listed in Table 2. Also, the interaction of ethyl magnesium bromide and cobalt

Card 1/2

Reactions of Organo-metallic Compounds With
Salts of Heavy Metals. 1. Interaction of
Ethyl Magnesium Bromide With Halides of
Titanium and Cobalt

S/062/60/000/03/04/007
B008/B006

chloride in the presence of the three above-mentioned acceptors in the temperature range -20°C - 40°C was investigated (Table 3). The data given in Tables 1 to 3 allow the assumption that the reaction of ethyl magnesium bromide with titanium- and cobalt salts - at least at the temperatures applied in the experiments - does not proceed via radical stages. It was seen in the experiments that the total yields of ethane and ethylene, and their proportion are not affected by the presence of the above acceptors. The authors mention a paper by N. V. Kondyrev and D. A. Fomina (Ref. 1), and one by Yu. V. Koryakin (Ref. 11). There are 1 figure, 3 tables, and 12 references, 4 of which are Soviet.

ASSOCIATION: Institut vysokomolekulyarnykh soyedineniy Akademii nauk
SSSR (Institute of High-molecular Compounds of the
Academy of Sciences, USSR)

SUBMITTED: July 14, 1958

Card 2/2

15.9201 2109,2209,1372

84511

S/190/60/002/004/012/020
B004/B056

AUTHORS: Van Fo-sun, Dolgoplosk, B. A., Yerusalimskiy, B. L.
TITLE: Polymerization of Isoprene Under the Influence of
Organomagnesium Compounds
PERIODICAL: Vysokomolekulyarnyye soyedineniya, 1960, Vol. 2, No. 4,
pp. 541-545

TEXT: The authors report on the polymerization of isoprene in cumene carried out by means of ether-free butylmagnesium chloride, bromide, iodide, dibutylmagnesium, and mixtures of butylmagnesium iodide and dibutylmagnesium, as well as of phenylmagnesium chloride and diphenylmagnesium at 90°C and a concentration of the monomeric isoprene of 60 mole%. Results are given in Table 1: yield, 4 - 50%; duration of the reaction, 20 - 40 hours; ratio between monomer and organomagnesium compound, 1 : 0.007 - 1 : 0.044; vitrification temperature, -3 to -10°C; intrinsic viscosity, 0.75 - 0.95; degree of unsaturation, 77 - 83%; content of 3,4-bonds, 93 - 98% (determined by means of infrared

Card 1/3

84511

S/190/60/002/004/012/020

B004/B056

Polymerization of Isoprene Under the
Influence of Organomagnesium Compounds

spectroscopy by Ye. I. Pokrovskiy at the authors' institute). The polyisoprene obtained in this way was perfectly soluble in benzene, probably because of intramolecular cyclization of the polymer chain. It mainly contained 3,4-bonds, so that the vitrification temperature is considerably increased in comparison to 1,4-polyisoprene. The kind of the halogen and radical of the organomagnesium compound did not affect the structure of the polymer. Fig. 1 shows the infrared spectrum of a polyisoprene sample. By adding complexing reagents, such as diethyl ether or triethylamine, the polymerization is retarded with an increase in the concentration of the reagents (Fig. 2, Table 2). Besides, a decrease in 3,4-bonds to 83 - 88% occurs (Table 3). In the experimental part, the authors describe the synthesis of ether-free organomagnesium compounds in cumene at 130 - 140°C, as well as in paraffin hydrocarbons in sealed ampoules at 135°C. There are 2 figures, 3 tables, and 5 references: 1 Soviet, 1 US, 1 British, and 2 German.

ASSOCIATION: Institut vysokomolekulyarnykh soyedineniy AN SSSR
(Institute of High-molecular Compounds AS USSR)

Card 2/3

84511

Polymerization of Isoprene Under the
Influence of Organomagnesium Compounds

S/190/60/002/004/012/020
B004/B056

SUBMITTED: December 18, 1959

Card 3/3

S/062/60/000/009/014/021
B023/B064

AUTHORS: Yerusalimskiy, B. L., Kavunenko, A. P., and Dolgoplosk, B.A.
TITLE: Reactions of the Free Radicals in Solutions. Communication
17. Effect of the Viscosity of the Medium on the Primary
Recombination of Free Radicals
PERIODICAL: Izvestiya Akademii nauk SSSR. Otdeleniye khimicheskikh
nauk, 1960, No. 9, pp. 1672-1674

TEXT: The authors investigated in how far the methane- and methyl aniline yields depend on the molecular weight and concentration of the polymer in the case of thermal splitting of methyl-phenyl triazene in the cumene - polystyrene system. As is shown by a previous paper of the authors (Ref.2), in solutions with 60% polystyrene (molecular weight 5000 to 200,000), the reaction leads to a reduction of the methane yield as compared to the data obtained from the use of a pure solvent. The methyl aniline yield remains, however, the same as that obtained in the absence of the polymer. Only in the solution of polystyrene with a molecular weight of 600,000, and a polymer concentration of 60%, the methyl aniline yield increases, while the

Card 1/3

Reactions of the Free Radicals in Solutions.
Communication 17. Effect of the Viscosity of
the Medium on the Primary Recombination of
Free Radicals

S/062/60/000/009/014/021
B023/B064

methane yield decreases considerably (Table 1). Consequently, the change of yields in methane solutions, containing polystyrene with a molecular weight of up to 200,000, cannot be considered as a result of the increase in viscosity of the medium. This would have certainly led to a higher yield of the product of methyl aniline primary recombination. The reduction of the yield is more likely to be due to the difference between the relative activity of polystyrene and that of cumene than to hydrogen donors. This is in agreement with published data, according to which the H atoms in polystyrene are less mobile than in cumene (Ref. 3). The authors proved that also in systems containing considerably lower polystyrene concentrations, the methane yield is reduced. The amount of the yield depends, as is shown in Table 2, on the concentration only. The molecular weight of the polymer has no effect upon the amount of the yield. In systems with a high viscosity, the importance of the primary recombination of free radicals increases. This becomes obvious by the fact that the methyl aniline yield increases, while the methane yield decreases at the same time. There are 2 tables and 5 references:

Card 2/3