

ASSOCIATION: none

Card 1/2

L 53738-65

ACCESSION NR: AP5015488

SUBMITTED: 17 Jul 63

ENCL: 00

NO REF SOY: 000

OTHER: 001

Card 2/2

YULIN, M.K.; VOL'NITSKIY, A.B.

Refining liquid alkyl phenols from the production of *p*-tert-butyl phenol. Neftepar. i neftekhim. no.1:31-33 1965.

1. Institut gorylovikh iskopayemykh, Moscow.

YULIN, S. I.

133-10-23/26

AUTHOR: Yulin, S. I., and Stetsenko, N. A. Engineers.

TITLE: The Influence of the Reduction Rate on Destruction of Carbide Lattice in the X12 and X12Φ Steels.
(Vliyaniye Stepeni Ukovki Na Razrusheniye Karbidnoy Setki v Stalyakh X12 i X12Φ.)

PERIODICAL: Stal', 1957, No.10, pp. 948-950 (USSR).

ABSTRACT: During forging of ingots of X12 and X12Φ1 steels of 300 and 500 kg in weight into rounds 40-180 mm and strip 60 x 300 x 180 mm, a network of eutectic carbides appeared. Causes of this phenomenon were investigated. It was found that the distribution of carbides in ingots of the above two steels along the radius of a cross-section of an ingot is independent from the weight of the ingot. In cast ingot metal the network of eutectic carbides increases from the surface towards the centre. Cast structure of X12 steel has a coarser eutectic carbide network than that of X12Φ1 steel (for ingots of the same weight). Hot mechanical working (forging) destroys the brittle network of eutectic carbides, but does not make the distribution of carbides in a cross-section more uniform. In the centre of forgings coarser grains with coarsely precipitated carbides on their boundaries are formed. Therefore, remains of Card 1/2the broken carbide network can be preserved there. An

133-10-23/26

The Influence of the Reduction Rate on Destruction of Carbide
(Cont.)

increase in the ingot diameter (and thus of weight), permits increasing the degree of reduction on obtaining forgings of the same diameter and thus to attain a higher degree of destruction of the network. There are 5 figures.

ASSOCIATION: Dneprospetsstal' Works. (Zavod Dneprospetsstal').

AVAILABLE: Library of Congress

Card 2/2

OKHOTIN, A.; YULIN, V.

It is time to solve these problems. Vnesh. torg. 43 no.12:19-22 '63.
(MIRA 17:2)

8(0)

SOV/112-59-4-7546

Translation from: Referativnyy zhurnal. Elektrotehnika, 1959, Nr 4, p 156 (USSR)

AUTHOR: Yulin, V. I.

TITLE: Automating the Measurement of the Amount of Dust Passing a Pneumatic Pipe

PERIODICAL: Byul. tsvetn. metallurgii, 1957, Nr 23, pp 21-22

ABSTRACT: A relay is described that operates on the accumulation of a certain amount of dust in a bunker. The relay has a cone-shaped tin hood fastened to a stem that carries a spring. As the amount of dust in the bunker increases, the hood and stem are moved in their guides and close contacts; the contacts, through an intermediate relay, turn on the bunker shutter drive. The shutter shaft is connected with a meter calibrated in units of weight. The relay size is: 150 x 150 x 180 mm. The service life under heavy operating conditions is 6 months.

M.A.U.

Card 1/1

PARAFONONOV, L.S.; SERIKOV, A.G.; YULINA, A.V.; RODIONOVA, N.V.,
telegrafistka, udarnik kommunisticheskogo truda;
RASKATAYEVA, M.F.; BPLYGIN, I.V.

We are discussing the project of the program of the CPSU.
Vest. svyazi 21 no.9:7-9 S '61. (MIRA 14:9)

1. Nachal'nik Nauchno-issledovatel'skogo instituta telefonnoy
svyazi Ministerstva svyazi SSSR (for Parafonov). 2. Glavnyy
inzhener Moskovskoy gorodskoy telefonnoy seti (for Serikov).
3. Rukovoditel' brigady kommuni ticheskogo truda TSentral'nogo
telegrafa SSSR (for Yulina). 4. TSentral'nyy telegraf SSSR
(for Rodionova). 5. Rukovoditel' brigady kommunisticheskogo
truda TSentral'nogo telegrafa SSSR (for Raskatayeva). 6.
Glavnyy inzhener Kiyevskogo oblastnogo upravleniya svyazi
(for Bulygin).

(Telecommunication)

GILEV, S.S.; SKVORTSOVA, Z.A.; SLEPOVA, V.A.; YULINA, L.N.

Photoelectric wedge densitometer. Usp.nauch.fot.no.4:82-87 '55.
(Densitometers) (MIRA 9:4)

YULINA, H. S.

Dissertation defended for the Candidate of Philosophical Sciences
at the Institute of Philosophy

"Criticism of the Philosophy of George Santayana."

Vestnik Akad. Nauk, No. 4, 1963, pp 119-145

YULISH, G.A.

Asthenoamnesic syndrome following general hypothermia
(freezing). Voen.-med. zhur. no. 1:74-75 Ja '66
(MIRA 19:2)

YULISH, G.A., podpolkovnik med. sluzhby

Syringomyelia reported from a garrison hospital. Voen.-med. zhur.
no. 11:75 N '57. (MIRA 11:4)

(SYRINGOMYELIA)

YULISH, K. (Praga)

Effect of minor changes in the shape of a blade on the frequency
of its bending vibrations. Izv. AN SSSR. Otd. tekhn. nauk. Mekh. 1
mashinostr. no. 4:104-108 J1-Ag '61. (MIRA 14:8)
(Blades--Vibration)

MONTITSKIY, R., starshiy nauchnyy sotrudnik; GORYACHEVA, M., mladshit
nauchnyy sotrudnik; YULIUS, A., mladshiy nauchnyy sotrudnik

Packing materials out of polymers. Sov.torg. 33 no.9:
48-50 S '59, (MIRA 12:12)

1. Nauchno-issledovatel'skiy institut trgovli i obshchest-
vennogo pitaniya.
(Synthetic products) (Packaging)

ROSKIN, G.I.; YULIUS, A.A.

Colloidolabile state of ribonucleic acid during the development,
intensive functioning, and malignant degeneration of cells.
Ark. anat. gist. i embr. 32 no.4:19-22 O-D '55. (MLRA 9:5)

1. Biologo-pochvennyy fakul'tet Moskovskogo universiteta imeni.
Lomonosova.

- (BREAST, neoplasms,
exper., ribonucleic acid in mouse mammary tumor cells,
comparison with embryonic & normal tissues)
- (BREAST, metabolism,
ribonucleic acid in embryonic, normal & malignant mouse
mammary tissues)
- (NUCLEIC ACIDS, metabolism,
mammary tissues, comparison in embryonic, normal &
malignant mouse mammary tissues)
- (NEOPLASMS, experimental,
mammary cancer in mouse, ribonucleic acid in tumor
tissue, comparison with embryonic & normal tissues)

YULIUS, A.A.; BAZHINOV, A.G.

Sterilization of solid media for fungus culture by β -propiolactone.
Mikrobiologiya 32 no.1:143-147 '63 (MIRA 17:3)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut fermentnoy i spirtovoy promyshlennosti.

LAPAN, A.P.; LARINA, V.A.; PISANOVA, L.I.; FURMAN, S.; YUL'KEVICH, L.P.

Phenols from waste waters of semicoking and other. Izv. Fiz.-
khim. nauch.-issl. inst. Izv. un. 4 no.2:233-254 '59.

(MIRA 16:8)

(Industrial wastes—Analysis) (Phenols)

YULKOVA, A. A.

"The Effect of Thermal Cycling on Dimensional and Structural Stability of Various Metals and Alloys", by A. A. Bochvar, G. J. Sergayev, A. A. Yulkova, L. I. Kolobneva, G. I. Tomson.

Report presented at 2nd UN Atoms-for-Peace Conference, Geneva, 9-13 Sept 1958

YUL'MAMEDOV, G.; NEVRAYEV, V. Yu. (Eng.)

"Results of Investigation of Work of Asynchronous Electric Drive with Power Supply from Source,"

paper read at the Session of the Acad. Sci. USSR, on Scientific Problems of Automatic Production, 15-20 October 1956.

Avtomatika i telemekhanika, No. 2, p. 182-192, 1957.

9015229

YUL'METOV, Sh.F.

*Find of Orbiculoidea in the Vereya beds of Tatarstan.
Paleont. zhur. no.4:93-94 '65. (MIRA 19:1)

1. Tatarskiy neftyanoy nauchno-issledovatel'skiy institut.
Submitted May 2, 1964.

YUL'MET'YEV, M.G.

Reliable way for the increase of the butterfat percentage of cows. Agrobiologiya no.6:879-893 N-D '65.

(MIRA 18:12)

1. Direktor uchebno-opytного khozyaystva Kazanskogo sel'skokhozyaystvennogo instituta imeni M.Gor'kogo.

L 6745-65 IMA.R./MWT(1)/ENC.200.20011-2/ENC.200.20011-2
Po-4 IJP(c)/USD/AS(mp)-2, 19D/ENC.200.20011-2/ENC.200.20011-2

ACCESSION NR: AP4043870

8/01/89/1

AUTHORS: Yul'met'yev, R. M.; Valiyev, R. A.

TITLE: Spin relaxation in Brownian rotational complexes in a liquid

SOURCE: IVUZ. Fizika, no. 4, 1989, 88-91

TOPIC TAGS: Brownian rotational spin lattice magnetic relaxation, electron resonance, relaxation

ABSTRACT: The Kubo and Tomita method (Kubo, 1958, 1954) is used to calculate the spin-lattice relaxation line $\omega_0/2$ and the spin-lattice relaxation of spins whose Stark interaction is described by an arbitrary rank. By way of a particular example, an interaction of fourth rank (with an electron magnetic moment). The calculations are made for $\omega_0/2$ and $\omega_0/2$.

L 6745-65

ACCESSION NR: AP4043870

liquid, having the form of spherical, symmetrical
drops. It is shown that the rate of magnetization
fusion of Brownian rotation of the paramagnetic
is a complicated process which takes place under
relaxations due to the anisotropic parts of the
spin energy, and also the interaction between the
thermal vibrations of the paramagnetic complex.
exact formula is derived for the rate of Stark
allowance for the shape of the paramagnetic complex
establish, by comparison with experimental data.
predominates under certain conditions. (Sov. Phys. Dokl.)

ASSOCIATION: Kazanski gospedinstitut (Kazanskii
Institute)

SUBMITTED: 11Feb63

SUB CODE: SA. NP

NR REF NO: 1

000 2/2

YUL'MET'YEV, R.M.

Nuclear magnetic relaxation in a liquid resulting from intermolecular interaction of nuclear spins. Zhur. strukt. khim. 6 no. 4:637-639 JI-Ag '65 (MIRA 19:1)

1. Kazanskiy pedagogicheskiy institut. Submitted November 12, 1964.

YUL'MET'YEV, R.M.

Quasi-classical approximation in the theory of slow neutron
scattering. Ukr. fiz. zhur. 10 no.11:1168-1175 N '65.
(MIRA 18:12)

1. Kazanskiy pedagogicheskiy institut. Submitted February 20,
1965.

S/056/63/044/002/022/055
B102/B186

AUTHORS: Valiyev, K. A., Timerov, R. Kh., Yul'met'yev, R. M.
TITLE: The influence of the molecular shape on the magnetic relaxation rate in liquids. II
PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, v. 44, no. 2, 1963, 522-529

TEXT: The authors continue previous investigations (ZhETF, 42, 503, 1962; Optika i spektroskopiya, 13, 505, 1962) on the Brownian rotation of molecules in a fluid. The probabilities of relaxative transitions between magnetic sublevels of nonspherical fluid particles have been calculated. These results are now used to determine the magnetic-resonance line widths and longitudinal relaxation times for such particles. The Kubo-Tomita method (J. Phys. Soc. 9, 888, 1954) is applied to obtain a relation between the relaxation times $T_{1,2}$ and the main values $D_{1,2,3}$ of the tensor D_{ii} or rotational diffusion that characterizes the Brownian rotation of the molecules. The calculations are made for quadrupole and dipole spin-
Card 1/4

The influence of the ...

S/056/63/044/002/022/065
B102/B186

spin interactions, and also for anisotropic g-factors, hyperfine and Stark interaction constants. The magnetic-resonance line half-width

$$\Delta\omega_{1/2} = 1/T_2 = \sum_{\beta} \sigma_{1\beta}^2 \tau_{1\beta} \quad \text{and} \quad 1/T = \sum_{\beta} \sigma_{0\beta}^2 \tau_{0\beta}. \quad \text{With}$$

$$f_{\alpha\beta}(\tau) = \langle \varphi_{-\beta}(\tau) \varphi_{\beta}(0) \rangle / \langle |\varphi_{\beta}|^2 \rangle = f(\tau) =$$

$$= \frac{1}{8\pi^3} \int \varphi_{\beta}(\alpha^0, \beta^0, \gamma^0) \varphi_{-\beta}(\alpha, \beta, \gamma) G(\alpha, \beta, \gamma; \tau | \alpha^0, \beta^0, \gamma^0; 0) \times$$

(15)

$$\times \sin \alpha^0 d\alpha^0 d\beta^0 d\gamma^0 \sin \alpha d\alpha d\beta d\gamma / \frac{2}{15} g_{\beta}^2.$$

$$\tau_{\alpha\beta} = \tau_{\beta} = \sum_i \Omega_i \rho(D_{ki}, \rho) = \sum_i \Omega_i D_{ki} (D_{ki}^2 + \beta^2 \omega_i^2)^{-1}. \quad (17),$$

$$\frac{1}{T_2} = \frac{1}{25} \left(\frac{eQg_{\beta}}{\hbar} \right)^2 \frac{I(I+1) - 3/4}{I^2(2I-1)^2} \sum_i \Omega_i [\rho(D_{ki}, 0) + \frac{5}{3} \rho(D_{ki}, 1) + \frac{2}{3} \rho(D_{ki}, 2)] \quad (21)$$

$$\frac{1}{T_1} = \frac{1}{25} \left(\frac{eQg_{\beta}}{\hbar} \right)^2 \frac{I(I+1) - 3/4}{I^2(2I-1)^2} \sum_i \Omega_i \left[\frac{2}{3} \rho(D_{ki}, 1) + \frac{8}{3} \rho(D_{ki}, 2) \right]. \quad (22)$$

Card 2/4

The influence of the ...

S/056/63/044/002/022/065
B102/B186

is obtained. For magnetic relaxation caused by innermolecular spin-spin interaction

$$\frac{1}{T_2(I)} = \sum_{l=-2}^2 \Omega_l(I) \sum_{p=-2}^2 \sigma_{lp}^2(I) D_{ll}(D_{ll}^2 + \beta^2 \omega_l^2)^{-1}, \quad (36)$$

$$\begin{aligned} \sigma_{11}^2 &= \sigma_{1-1}^2 = \frac{2}{3} \sigma_{11}^2 = \frac{2}{3} \sigma_{10}^2 = \frac{2}{3} \sigma_{01}^2 = 2\sigma_{0\pm 1}^2 = \frac{2}{3} \sigma^2, \\ \sigma_{1-2}^2 &= \sigma_{00}^2 = 0; \quad \sigma^2 = \frac{2}{3} I(I+1) \gamma^4 \hbar^2 r_{ij}^{-6}. \end{aligned} \quad (37)$$

is obtained in the case of equivalent nuclei. In the case of electron resonance in liquids, line width and relaxation time are given by

$$\begin{aligned} \Delta\omega_{l,m} &= \frac{2}{15} \hbar^{-1} \sum_{l=-2}^2 \left\{ \left[\frac{2}{3} g_p^2 + \frac{1}{2} g_d^2 f_m \right] \Omega_l^{(n,m)} \rho(D_{ll}, \omega_{l,m}) + \right. \\ &+ \left. \left[\frac{1}{2} g_p^2 + \frac{7}{12} g_d^2 f_m \right] \Omega_l^{(-1,m)} \rho(D_{ll}, \omega_{-1,m}) + \frac{2}{3} \left(S(S+1) - \frac{3}{4} \right) g_d^2 \Omega_l^2 \times \right. \end{aligned} \quad (44)$$

Card 3/4

S/056/63/044/002/022/065
B102/B186.

The influence of the ...

$$\begin{aligned} & \times \{ \rho(D_{kl}, \omega_{0,0}) + \frac{4}{3} \rho(D_{kl}, \omega_{10}) + \frac{2}{3} \rho(D_{kl}, \omega_{2,0}) \} \} \\ T_{1m}^{-1} = & \frac{4}{15} \hbar^{-2} \sum_{l=-2}^2 \left\{ \left[\frac{1}{2} g_p^2 + \frac{7}{15} g_d^2 f_m \right] \Omega_l^{-1, m} \rho(D_{kl}, \omega_{-1, m}) + \right. \\ & \left. + \frac{4}{5} (S(S+1) - \frac{3}{4}) \Omega_l^d g_d^2 \{ \rho(D_{kl}, \omega_{1,0}) + 4 \rho(D_{kl}, \omega_{2,0}) \} \right\}. \end{aligned} \quad (45).$$

It may be seen that for $S > 1/2$ the main contribution to the line width is due to Stark interaction of the particle spins. Apart from the broadening caused by the Brownian rotation, there is also a broadening due to the interaction between spin and inner oscillations of the molecule. The latter is equal for all hyperfine components, as is the case for the Stark broadening.

ASSOCIATION: Kazanskiy pedagogicheskiy institut (Kazan' Pedagogical Institute), Fiziko-tekhnicheskiy institut Kazanskogo filiala Akademii nauk SSSR (Physicotechnical Institute of the Kazan' Branch of the Academy of Sciences USSR)
June 11, 1962

SUBMITTED:
Card 4/4

YUL'NET'YEV, R.M.: VAI'YEV, A.A.

Spin relaxation in NMR of paramagnetic complexes
in fluids. Izv. vys. ucheb. zav.; 112, no. 1:85-93, 1964
(MDR 17:8)

1. Kiyevskiy gosudarstvennyy universitet.

VALIYEV, K.A.; TIMEROV, R.Kh.; YUL'MET'YEV, R.M.

Effect of the shape of molecules on the rate of magnetic
relaxation in liquids. Part 2. Zhur. eksp. i teor. fiz.
44 no.2:522-529 F '63. (MIRA 16:7)

1. Kazanskiy pedagogicheskiy institut i Fiziko-tekhnicheskiy
institut Kazanskogo filiala AN SSSR.

YUL'MET'YEV, R.M.

Effect of the collective motion of particles on translation
diffusion in liquids. Zhur.strukt.khim. 6 no.5:676-683 S-0
'65. (MIRA 18:12)

1. Kazanskiy pedagogicheskiy institut. Submitted May 13, 1964.

I 10262-66 ENT(1) LTP(c) BW/ERS/CC/RE

1. The first part of the document is a list of names.

2. The second part of the document is a list of names.

3. The third part of the document is a list of names.

4. The fourth part of the document is a list of names.

5. The fifth part of the document is a list of names.

6. The sixth part of the document is a list of names.

7. The seventh part of the document is a list of names.

8. The eighth part of the document is a list of names.

9. The ninth part of the document is a list of names.

10. The tenth part of the document is a list of names.

11. The eleventh part of the document is a list of names.

12. The twelfth part of the document is a list of names.

ACC NR:

AP5001651

CHG 11/1

ATTN: STERIL

11/16/83

Card 2/2

FISHER, I.Z.; YUL'FETEV, R.H.

Use of an electronic computer in studying thermal motion
in a fluid. Usp. fiz. nauk 87 no.2:374-378 0 '65.

(MIRA 18:11)

VAYNSHTEYN, E.Ye.; PAVLENKO, A.S.; TURANSKAYA, N.V.; YULOVA, T.G.

Effect of the distribution of rare earth elements in rocks on petrochemical factors and its significance for the solution of petrogenetic problems. Geokhimiia no.12:1077-1086 '61.

(MIRA 15:3)

1. Vernadskiy Institute of Geochemistry and Analytical Chemistry, Academy of Sciences, U.S.S.R., Moscow.
(Rare earth metals) (Petrology)

YULOVSKI, A. A.

Calculation of artillery projectiles
Leningrad, 1933, 67 p. (54-45765)

UF750.18

YULOVSKIY, N. V.

PA 19T79

USSR/Vacuum Tubes, Transmitting
Filaments

Jul 1946

"Alternating Current Filament Supply for Transmitter Tubes," N. V. Yulovskiy, 3 pp

"Vestnik Svyazi - Elektro Svyaz'" No 7 (76)

Well illustrated article which shows the difference between feeding transmitter tube filaments by an alternating or direct current. The former is said to be more advantageous. This article makes reference to an article published in "Vestnik Svyaz" No 3, 1945 which gave conditions under which an alternating current had to be used as well as the type of equipment.

19T79

YULOVSKIY, N.V.
Category : USSR/Radiophysics - Generation and conversion of radio-frequency oscillations

I-4

Abs Jour : Ref Zhur - Fizika, No 1, 1957, No 1842

Author : Zaryanov, N.V., Yulovskiy, N.V.

Title : Use of Beam Power Tetrodes in Short-Wave Transmitters

Orig Pub : Vestn, svyazi, 1956, No 7, 3-5

Abstract : No abstract

Card : 1/1

YULOVSKIY, N.V., inzhener.

Obtaining and maintaining high quality indices in radiobroadcasting
transmitters. Vest. svyazi 17 no.4:5-7 Ap '57. (MLRA 10:5)
(Radio--Transmitters and transmission)

PHASE I BOOK EXPLOITATION 920

U.S.S.R. Ministerstvo svyazi. Tekhnicheskoye upravleniye

Korotkovolnovyye radioperedayushchiye ustroystva; informatsionnyy sbornik. (Shortwave Radio Transmitting Apparatus; Collection of Information) Moscow, Svyaz'izdat, 1958. 150 p. (Series: Tekhnika svyazi) 14,500 copies printed.

Resp. Ed.: Rozentsveyg, I.Ye.; Ed.: Novikova, Ye.S.; Tech. Ed.: Shefer, G.I.

PURPOSE: The book is intended for engineering and technical personnel working in radio communication.

COVERAGE: The monograph contains two articles describing two different broadcast transmitters of 120 kw and 50 kw for operation in the short-wave band. The book provides general information and discusses the electrical and acoustic indices, as well as elementary circuits, of the high-frequency and low-frequency stages. It also presents elementary power supply circuits, control circuits and

Card 1/4

Shortwave Radio Transmitting Apparatus (Cont.) 920

blocking and signaling circuits of the transmitters and describes their operation. The articles also describe improvements of the equipment for the KVM Transmitter. No personalities are mentioned. There are no references.

TABLE OF CONTENTS:

Foreword

2

N.V.Yulovskiy. A High-power (120-kw) Short-wave Broadcast Transmitter (Type KVM)

3

The article contains a block diagram of the transmitter and describes the operation of various high-frequency stages. The article also discusses the master oscillator section, which includes such components as a crystal oscillator, high-frequency amplifier, frequency calibrator, selenium rectifier, bias rectifier, voltage stabilizer, voltage rectifier, and control block. The author describes the low-frequency stage which includes a modulator, submodulator, low-frequency amplifier, and

Card 2/4

Shortwave Radio Transmitting Apparatus (Cont.) 920

linear amplifier. He also describes the power supply circuits for the tube filaments, plates and grids and discusses the circuit of a thyatron rectifier. Systems for transmitter power distribution, control, blocking, and signaling are described and a discussion of the water-cooling of tubes and coils is presented. Methods of switching-on tube filament circuits and low- and high-power rectifiers are discussed. The article also describes safety circuits and the arrangement of transmitter equipment in various cabinets. A description of changes and improvements in the initial design of the transmitter is given.

N.N.II'ina. A 50-kw Short-wave Broadcast Transmitter

105

The transmitter described in this article represents a modern set-up used for high-quality broadcasting. It has certain advantages over the 120-kw transmitter discussed in the preceding article. A block diagram of the transmitter is given and a description of the operation of various high-frequency stages is presented. The article describes a VChD-100 type master oscillator circuit and the low-frequency stage, which includes

Card 3/4

Shortwave Radio Transmitting Apparatus (Cont.) 920

the circuits of the modulator, linear amplifier, low-frequency filter, level indicator, and signaling key. It also presents the d-c and a-c power supply circuits and discusses high- and low-power thyatron rectifiers. It describes the control, blocking, and signaling systems of the transmitter as well as the air- and water-cooling systems. A discussion of the arrangement of transmitter equipment in various cabinets is given and the operation of the entire set-up is described.

AVAILABLE: Library of Congress (TK6553.R9)

JP/whl
12-8-58

Card 4/4

YULOVSKIY, N.Y., ingh.

Use of anode self-modulation in shortwave radio transmitters. Vest.
svyazi 20 no.10:9-12 0 '60. (MIRA 13:11)

(Modulation (Electronics))

(Radio, Shortwave--Transmitters and transmission)

S/109/62/007/004/000/012
D230/D302

9.4230

AUTHORS: Gaponov, A.V., and Yulpatov, V.K.

TITLE: Interaction of closed electron beams with an electromagnetic field in hollow cavities

PERIODICAL: Radiotekhnika i elektronika, v. 7, no. 4, 1962,
631 - 643

TEXT: Expressions are obtained for the fields excited in a cavity by a thin closed electron beam and for the case of an arbitrarily-distributed electron stream in a cavity volume, the condition being that in an unexcited state all electrons move in closed trajectories. In deriving the equation for the frequencies of normal cavity oscillations with a closed electron beam, the examination is limited to the case when the unexcited electron ring is stationary and does not change with time; it is assumed that the current in the unexcited beam is constant. The system is in equilibrium when the field in the resonator is alternating and the current in the beam is constant. The investigation of the equilibrium of the system reduces to studying small electron oscillations close to the unexcited state.
Card 1/3

Interaction of closed electron beams ... S/109/62/007/004/006/018
D230/D302

ted trajectories under the action of a matched alternating e.m. field. The resonator losses can be calculated from the characteristic equation. The possibility of self-excitation of the microwave oscillations in the cavity with a ring electron beam is examined; in this case the resonant interaction between the electron stream and the e.m. field in the cavity is studied when one of the natural beam frequencies is close to the partial frequency of the cavity. A third-order equation having complex roots is deduced showing that self-oscillation can take place in a microwave with an electron beam. Certain cases of interaction between the ring electron beam and the e.m. cavity fields are considered. In a quadrupole electric cavity field with the beam lying in the plane $z = 0$ the requisite spatial grouping of the electrons is caused by the z -component of the field falling into the retarding field always in the same phase. When the ring electron beam is in a constant cavity field it is shown that self-excitation depends on a relativistic effect. All examples of self-excited oscillations are discussed in terms of the electron gyro-frequency. Self-excitation can also take place on harmonics of the gyro-magnetic frequency; for this purpose the corresponding spatial field harmonics in the region occupied by the beam

Card 2/3

Interaction of closed electron beams ... S/109/62/007/004/005/012
D230/D302

should be different from zero; generally, this is the case in any constant field. There are 5 figures and 20 references: 18 Soviet-bloc and 2 non-Soviet-bloc.

SUBMITTED: August 28, 1961

Card 3/3

8/24/77
F14-07177

AUTHOR: Yulpatov, V.K.

TITLE: On the dispersion equation for a
curvilinear electron beam

PERIODICAL: Izvestiya vysshego uchebnogo zav.
v.o, no.1, 1963, 95-103

TEXT: Previous investigations of interaction of curvilinear
electron beams and electromagnetic fields in waveguides in the

small-signal approximation have not taken into account the
influence of space charge. This problem is solved in the
present article for waveguides with arbitrary cross-sections.
The dispersion equation is derived in the form of a determinant
of variations, permitting the influence of the space charge
of the beam to be taken into account. The case of a beam
and waveguide are infinitely long and the fields and particle
trajectories are parallel to the axis of the waveguide.
The trajectories of the particles are shifted in space. The expression for
the dispersion equation is obtained. A qualitative idea of the dispersion
characteristics is given.

On the dispersion equations for ...

simplest cases it is extremely difficult to find the coefficients. One case where these coefficients are found is when the interaction occurs at not too high a frequency. It is found that in interaction of the synchronous wave with the asynchronous wave, the space charge instability.

ASSOCIATION: Nauchno-issledovatel'skiy institut pri Gor'kovskom universitete
(Scientific Research Institute of Gor'kiy University)

SUBMITTED: July 9, 1962

Card 2/2

1.11.10-1
150-11P/01

SECRET (S--A--T)

ACCESSION NO. 150-11P/01

AUTHOR. Yelgatox, V. K.

TITLE. [illegible]
[illegible]

SECRET (S--A--T)
410-413

was considered. It was pointed out that the possibility of [illegible]

ACCESSION NR: AP 30001AR

ACCESSION NR: 30001AR
ACCESSION NR: 30001AR

ACCESSION NR: 30001AR

YULUSHEV, I.G.

Effect of mixed organic-mineral fertilizers on buckwheat. *Agrobiologiya*
no.4:609-611 J1-Ag '64. (MIRA 17:12)

1. Kirovskiy sel'skokhozyaystvennyy institut.

S/065/61/000/008/004/009
E030/E335

AUTHORS: Silich, M.I., Sidorov, I.P., Martynova, L.L.,
Bukarov, A.R., Yulusov, A.A. and Kisil', I.M.

TITLE: Improved Process for Obtaining Alcohols by the
Oxo-synthesis Method With Suspended Catalyst

PERIODICAL: Khimiya i tekhnologiya topliv i masel, 1961,
No. 8, pp. 19.- 24

TEXT: The authors mention briefly the drawbacks of the
existing technological schemes for obtaining alcohols by
oxo-synthesis. The main drawbacks of the scheme with
suspended catalyst are the erosion of the throttle elements,
the need for paste pumps for transporting the catalyst (which
is in suspension in the liquid) and the existence of a
filtering section which work intermittently. Periodic switching
between gas and liquid streams, a complicated automatic
control and the decomposition of the cobalt carbonyl (decobal-
tisation) are the chief drawbacks of the other two schemes.
The present paper deals with improving the scheme with sus-
pended catalyst. The tests were carried out on a model and
in a pilot plant. In the present process the synthesis occurs
Card 1/4

Improved Process

S/065/61/000/008/004/009
E030/E335

in the liquid phase and therefore a solvent is used which is isobutyl alcohol at the start of the reaction, changing to the final product as the reaction proceeds. In the laboratory tests a propane-propylene feedstock with 74 to 85% propylene was used, the ratio of raw material to solvent being nearly 1:2 and that of CO to hydrogen 1:1.2. In the pilot plant, synthesis gas was used as feed, with the ratio of hydrogen to carbon monoxide varying between 0.5:1 to 7.5:1, the other parameters being nearly the same as those in the laboratory tests. In order to eliminate the deficiency in the filter system, a re-cycle system using a centrifugal separator was introduced. This system (developed in conjunction with NIIKhIMMASH under the direction of Senior Engineer G.K.Ivanova) enables the filters to work for long periods without cleansing and, by returning the catalyst-rich fraction to the reactor, diminishes the quantity of product going for decobaltisation, filtering, hydrogenization and rectification. Thus, the process of obtaining butyl alcohols is carried out in three stages: 1) production of cobalt carbonyls and hydroformylation of propylene; 2) decomposition of cobalt carbonyls

Card 2/4

Improved Process

S/065/61/000/008/004/009
E030/E335

(decobaltization) and 3) hydrogenization of aldehydes and alcohols. In the previous two-stage process only alcohols were obtained as the final product; in the present three-stage one aldehydes also are obtained. It has been shown that by hydroformylation at 300 atm. and 125 °C the content of n-aldehydes in the final product increases. It has also been found that at temperatures of 110 to 140 °C and pressures of 25 to 100 atm the catalyst decomposes completely. At 135 °C and 300 atm. propylene converts to n-aldehydes (63%), iso-aldehydes (21%), high aldehydes (11.4%) and by-products (46%), the ratio of n- to iso-aldehydes being 3:1. With decreasing pressures this ratio decreases, being 2.2:1 at 250 atm. and 1.6:1 at 200 atm. During the oxo-reaction carried out in the pilot plant at temperatures between 135 and 160 °C, a pressure of synthesis gas of 180-200 atm., content of catalyst of 1-2% and contact time 45 min., a product with a ratio of n- to iso-aldehydes of approximately 2:1 was obtained. This product hydrogenated in a mixture of butyl alcohols in the same ratio. G.N. Klinova, A.D. Yerofeyeva, N.M. Malygina, A.I. Khokhlov, A.I. Zaytseva, T.V. Yelisova and A.I. Busygina
Card 3/4

Improved Process

S/065/61/000/008/004/009
E030/E335

participated in the tests. There are 3 figures, 2 tables and 11 references: 4 Soviet and 7 non-Soviet. The four latest English-language references quoted are: Ref. 3 - H. Keulemans - U.S. Patent No. 2587858, 1952; Ref. 4 - I. Mertzweeler, W.M. Smith, U.S. Patent No. 2725401, 1955; Ref. 6 - Petroleum 16, No. 10, 291, 1953; Ref. 7 - I. Kirshenbaum, K.L. Hughes - Petr. Refin., 37, No. 6, 209, 1958.

ASSOCIATION: GIAP, LKhK and OKBA

Card 4/4

SILICH, M.I.; SIDOROV, I.P.; MARTYNOVA, L.L.; BUKAROV, A.R.;
YULISOV, A.A.; KISIL', I.M.; Prinimali uchastiye: KLINOVA, G.N.;
YEROFEYEVA, A.D.; MALYGINA, N.M.; KHOKHLOV, A.I.; ZAYTSEVA, A.I.;
YELISOVA, T.V.; BUSYGINA, A.I.

Improved technological system with a suspended catalyst
for the production of alcohol by oxo synthesis method. Khim.i
tekh.topl.i masel 6 no.8:19-24 Ag '61. (MIRA 14:8)

1. Gosudarstvennyy institut azotnoy promyshlennosti; LKhK;
Opytno-konstruktorskoye byuro po avtomatike.
(Alcohols) (Oxo process)

30221

S/087/61/000/019/066/085
B117/B110

1/1.0150

AUTHORS: Klimenok, B. V., Yul'yakshina, K. G.

TITLE: The problem of the influence of organic sulfur compounds on the rate of deparaffination of the diesel fraction with carbamide

PERIODICAL: Referativnyy zhurnal. Khimiya, no. 19, 1961, 421 - 422, abstract 19M158 (Sb. "Khimiya seraorgan. soyedineniy, soderzhashchikhsya v neftyakh i nefteproduktakh". M., AN SSSR, 1959, 150 - 153)

TEXT: Two individual substances whose boiling points lie within the boiling range of the diesel fraction (230 - 350°C) were used to study the influence of organic sulfur compounds on the rate of complex formation during the reaction of the diesel fraction with aqueous carbamide solution: 2,8-dimethyl-5-thianonane (I) and dibenzo thiophene (II). The latter were introduced into the diesel fraction which had been carefully purified with aluminosilicate. The maximum concentration of each organic sulfur compound was 4% by weight. It was found that I and II, unlike tars, have

Card 1/2

The problem of the influence...

30221
S/081/61/000/019/066/085
B117/B110

virtually no effect on the rate of complex formation. Furthermore, it was shown that I and II do not display any surface-active properties. In view of the behavior of the two above-mentioned compounds, the authors voiced the assumption that organic sulfur compounds do not affect the process of complex formation in deparaffination with aqueous carbamide solution. [Abstracter's note: Complete translation.] X

Card 2/2

1. YUL'YANISHCHEV, M. I.

2. USSR (600)

4. Apricot

7. Non-related hybridization in apricots. *Agrobiologia* No. 5 1952

9. Monthly List of Russian Accessions, Library of Congress, January 1953. Unclassified.

YUL'YANKIN, I., RA'IKO, T.

Milking

Milking cows by machine in the pasture. Kolkh. proiz. 12 no. 6, 1952.

Monthly List of Russian Accessions, Library of Congress, October, 1952, UNCLASSIFIED

YUL'YANKIN, I. Eng.

Hydraulic Rama

Hydraulic ram. Kolkh. proizv. 12 no. 9, 1952.

9. Monthly List of Russian Accessions, Library of Congress, December 1958. Unclassified.

1. YUL'YANOV, M. I.
2. SSSR (600)
4. Neuroses
7. Certain hematological shifts in neurosis; first communication.
Klin. med. 30 No. 10, 1952
9. Monthly Lists of Russian Accessions, Library of Congress, March 1953, Unclassified.

YUL'YEV, F.

Mysterious neighbor. Znan.sila.36 no.3:6-7 Mr '61.
(Venus(Planet))

(MIRA 14:3)

TIMOFEYEV, M., yurist; KRIVTSOV, G., yurist; YUL'YEV, I. (g.Saratov)

Our consultations. Sov. profsoiuzy 18 no.8:46 '62. (MIRA 15:4)
(Employees, Dismissal of) (Overtime) (Exigent domain)

SOKOLOV, L.H., kandidat tekhnicheskikh nauk; ZAISSKIY, V.I., professor;
YULYUTIN, V.P., professor, doktor.

Resistance to deformation of industrial titanium. Sbor.Inst.stali
no.33:142-153 '55. (MLRA 9:6)

1.Kafedra kovki i shtampovki i Kafedra metallurgii redkikh metallov.
(Titanium--Testing)

AUTHOR: Ponomosov, A. V.; Yum', A. A.; Murashova, I. I.

TITLE: Electrolytic production of low dispersion nickel

CITED SOURCE: Tr. 7 Vses. nauchno-tekhn. konf. po metallurgii, Yerevan, 1984, 78-82

TOPIC TAGS: powder metallurgy, powder metal process, electrolyte, electrolysis, dispersion characteristics

TRANSLATION: A new sulfate-chloride electrolyte is used for production of nickel. The composition of the electrolyte is: 90-100 grams/liter NiSO_4 , 200 grams/liter NH_4Cl , 200 grams/liter NaCl . The presence of NH_4Cl maintains a constant pH over the whole time of the process.

Card 1/2

L 57543-65

ACCESSION NR: AR5015153

electrical conductivity of the electrolyte. The greater part to the current was attained at a current density of 100 A/cm². The specific consumption of electric power in the laboratory was 2200 kilowatt hours/T. The particle size of the powder is sufficiently wide limits by varying the concentration of the electrolyte, the temperature of the electrolyte, and the current density. In the process conditions, there are required a constant current and the removal of a determined amount of electrolyte from the electrolyte. 7. Kvin

SUB CODE: MM

ENCL: 00

SUB CODE: MM

ENCL: 00

Cord 2/2

YUMADILOV, A.Yu.

Completion of wells by the sand jet method in fields of the Oil
Field Administration of the Chekmagush Petroleum Trust. Nefteprom.
delo no.7:13-14 '65. (MLRA 18:8)

1. Neftepromyslovoye upravleniye "Chekmagushneft".

YUMAGULOV, B. L.

Cand Agr Sci - (diss) "Basic problems of agrotechniques of corn
in the north of the Tselinnyy Kray." Alma-Ata, 1961. 100 p.
(Ministry of Agriculture Kazakh SSR, Kazakhistan: Scientific
search Inst of Farming imeni V. R. Viliyeva. 1961.
not given; (KL, 7-61 sup, 253)

YUMAGULOV, G.L.

Farming in virgin areas without clean fallow. Zemledelie 24
no.2:10-17 F '62. (MIRA 15:3)

1. Severo-Kazakhstanskaya oblastnaya gosudarstvennaya sel'sko-
khozyaystvennaya opytnaya stantsiya.
(Following)

KOZHEVNIKOV, A.R., prof.; POPOVA, G.I., dots.; VOROZHTSOV, I.P.,
kand. tekhn. nauk, dots.; GERASENKOV, B.I., kand. sel'-
khoz. nauk; YUMAGULOV, G.L., kand. sel'khoz. nauk;
MAR'YASOV, V.G., assistant; VINOGRADOVA, N.I., kand. sel'-
khoz. nauk; ROKTANEN, L.P., dots., kand. biol. nauk;
KOKHOMSKIY, F.M., Geroy Sotsialisticheskogo Truda, zasl.
zootekhnika RSFSR; MAKHNOVSKIY, M.K., dots., kand. ekon.
nauk; ARTAMONOV, F.D., assistant; MAKAROVA, I.V., red.

[Corn in the Virgin Territory and Western Siberia] Kukuruza
v tselinnom krae i Zapadnoi Sibiri. Moskva, Kolos, 1965.
229 p. (MIRA 18:9)

1. Omskiy sel'skokhozyaystvennyy institut im. S.M.Kirova
(for Kozhevnikov, Popova, Mar'yasov, Vinogradova, Kokhomskiy,
Makhnovskiy, Artamonov).
2. Zamestitel' direktora po nauchnoy
rabote Severo-Kazakhstanskoy opytnoy stantsii (for Yumagulov).
3. Zaveduyushchiy laboratoriyey kukuruzy Sibirskogo nauchno-
issledovatel'skogo instituta sel'skogo khozyaystva (for
Gerasenkov).
4. Tselinogradskiy sel'skokhozyaystvennyy institut
(for Roktanen).

AKHMETOV, K.T.; KORNEYEV, V.F.; POPOV, N.A.; YUMAKAYEV, Sh.I.

Accelerating processes of leaching zinc calcines and an increase in
labor productivity. Trudy Alt. GIMII AN Kazakh. SSR 14:178-190 '63.
(MIRA 16:9)

(Zinc—Heat treatment) (Leaching)

S/661/61/000/006/036/031
D202/D302

AUTHORS: Alashkevich, M. L., Leznov, N. S., Yumakova, A. Ye. and
Andrianov, K. A.

TITLE: Physico-mechanical properties of linear polydiethylsi-
loxanes

SOURCE: Khimiya i prakticheskoye primeneniye kremneorganichskikh
soyedineniy; trudy konferentsii. no. 6: Doklady, diskus-
sii, resheniye. II Vses. konfer. po khimii i prakt. prim.
kremneorg. soyed., Len., 1958. Leningrad, Izd-vo AN SSSR,
1961, 171-172

TEXT: A supplement to a previous report in no. 2, p. 20, of this
publication. The authors compare the properties of polymethyl-phe-
nyl-siloxanes with those of polydiethyl-siloxanes used as high-va-
cuum pump fluids. No experimental details are given. It was found
that the first compounds have marked advantages over the second,
although cyclic polymers, formed during their synthesis, unfavorab-
ly affect the thermal stability of both. ✓

Card 1/1

CA

YUMANOVA, L.V.

The theory of hydrolysis of chlorides. The minimum hydrolysis of magnesium chloride in the drying of its hydrates. I. G. Sherbekov and L. V. Yumayeva. *Khimiya* (1954, No. 4, 37-41). The hydrolysis decreases (1) with a decrease of the proportion of the water that is to be removed, (2) with a decrease of temp. to which the hydrate is heated and (3) with a decrease of the original water content. The hydrolysis depends on the velocity of the heating. The hydrolysis is min. when the temp. is increased at the rate of 1° per min. Twelve references. A. Benoit

***** METALIPYCNAL LITERATURE CLASSIFICATION *****

YUMANOVA, L. V.

000

1. SHCHERBAKOV, I. G., YUMANOVA, L. V.

2. USSR (600)

Ural Scientific-Research Chemical Institute. "The Cathode Process in the Formation of a Diaphragm Coating" Tsvet. Met. 14, No. 6, 1939.

9. ~~Report~~ Report U-1506, 4 Oct. 1951.

YUMANOVA, L. V.

YUMANOVA, L. V., MIKULINSKIY, A. S., I GEL'D, P. V.

36181 O temperature shikhty i davlenii gazov v karbidnoy pechi. V sb: Teoriya i praktika rudnoy elektrotermii. Sverdlovsk-Moskva, 1948, S. 91-94.

SO: Letopis' Zhurnal'nykh Statey, No. 49, 1949

YUMANOVA, L. V. and MIKULINSKIY, A. S.

Yumanova, L. V. and Mikulinskiy, A. S. "The method and results of investigation of the space under the electrodes of a carbide furnace", (Report), Scobschch. o nauch. rabotakh chlenov Vsesoyuz. khim. o-va im. Mendeleyeva, 1949, Issue 1, p. 23-25.

SO: U-4630, 16 Sept. 53, (Letopis 'Zhurnal 'nykh Statey, No. 23, 1949).

USSR/Chemical Technology - Chemical Products and Their Applications, Mineral Salts. Oxides. Acids. Bases. I-6

Abs Jour : Ref Zhur - Khimiya, No 3, 1957, 8790

Author : Yumanova, L.V. and Mikulinskiy, A.S.

Inst : Uralsk Science Research Chemical Institute.

Title : Phase Changes and Chemical Changes in the Charge During the Reduction of Oxides of Calcium and Silicon by Carbon.

Orig Pub : Tr. Ural'skogo n-i. khim. in-ta, 1954, No 2, 166-176

Abstract : A laboratory method has been developed for the investigation of the reduction of the oxides taking into account the filtering effect of the charge in industrial furnaces using electrodes inserted in the charge. Some information has been obtained on the reduction of CaO and the mechanism of the reduction process. Most of the CaC_2 is formed in the lower high-temperature zone of the furnace where the liquid phase collects. An increase in temperature from 1800 to 1900° increases the conversion to carbide

Card 1/2

USSR/Chemical Technology - Chemical Products and Their
Application, Mineral Salts. Oxides. Acids. Bases.

I-6

Abs Jour : Ref Zhur - Khimiya, No 3, 1957, 8790

above 10-30%. The excess C in the top carbonaceous layer of the charge is the product of the dissociation of CaC_2 . The C and elementary Ca diffuse to the upper layers of the charge and cover the surface of the CaO and C chunks and diffuse into the interior of the latter. SiO_2 and its dissociation products are reduced in the vapor state. When sufficient iron filings are added to the charge to form 45% FeSi, metallic beads are formed, the carborization of the Fe being accompanied by its enrichment in Si in the upper layers of the charge at 1100-1200°.

Card 2/2

SOV/137-58-11-22255

Translation from: Referativnyy zhurnal. Metallurgiya, 1958, Nr 11, p 59 (USSR)

AUTHORS: Yumanova, L. V., Germaidze, M. S.

TITLE: Distillation of Potassium From a Potassium-lead Alloy (O distillyatsii kaliya iz splava svinets-kaliy)

PERIODICAL: Tr. Ural'skogo n.-i. khim. in-ta, 1957 (1958), Nr 5, pp 66-80

ABSTRACT: The flowsheet for producing K through the medium of a K-Pb alloy includes electrolytic production of the alloy and distillation of the K therefrom. A study is made of the distillation of K from alloys containing ~ 6-13% K. Conditions of distillation permitting reduction in K content to 0.1-0.4% are recommended, as follows: Temperatures 630-680°C, residual pressure 0.04-1.5 mm Hg. It is noted that condensation of K on a water-cooled condenser occurs in the 200-300° interval. It is observed that the process rate is determined by the size of the surface of evaporation; within certain limits, the thickness of the alloy layer is not of decisive significance owing to the natural turbulence within the alloy which is caused by the reduction in specific gravity as the concentration of alloy is changed. Flowsheets and descriptions are provided of

Card 1/2

SOV/137-58-11-22255

Distillation of Potassium From a Potassium-lead Alloy

semicontinuous laboratory distillation equipment now in use and of pilot-plant equipment being planned. Both permit the distillation operations of transferring the alloy, and the K, to proceed under air-tight conditions. In large-scale laboratory experiments using semicontinuous equipment a K yield of 84% with a loss of 0.2% thereof is attained. The Pb content of the condensate is 0.36%. It is found that contamination of the K by lead diminishes with an increase in the height of the distillation chamber.

L. P.

Card 2/2

YUMARKOV, M.; KVIHT, M.

The project has been crowned with success. Izobr.1 rats. no.2:
31-32 F '60. (MIRA 13:8)
(Aluminum hydroxide)

YUMASHEV, A., mayor.

Executing flight maneuvers with a jet plane. Kryl.rod. 8 no.1:11-13
Ja '57.

(MLBA 10:5)

(Jet planes--Piloting)

TOKMALAYEV, S.P., dotsent [deceased]; KUZNELEV, N.S., dotsent; OSTROVI-
TIANOV, K.V., akademik; ALEKSEYEV, A.M., dotsent; KUDROV, V.M.;
LEONT'YEV, L.A. Prinsipali uchastiye: BELYATEVA, Z.N., kand.ekon.
nauk; MRACHKOVSKAYA, I.M., kand.ekonom.nauk; RYNDINA, M.N.,
kand.ekonom.nauk; SHIRINSKIY, I.D., kand.ekonom.nauk; red.;
YUMASHEV, A.I., kand.ekonom.nauk; PROKOP'YEV, S.P., red.; NAUMOV,
K.M., tekhn.red.

[Capitalist production method] Kapitalisticheskiy sposob pro-
izvodstva. Moskva. Pt.2. 1960. 357 p. (MIRA 13:10)

1. Kommunisticheskaya partiya Sovetskogo Soyuz. Vysshaya
partiynaya shkola. 2. Chlen-korrespondent Akademii nauk SSSR (for
Leont'yev).

(Economics)

YUMASHEV, Aleksandr Ivanovich; KOROTKEVICH, G.Ya., red.; NAUMOV,
K.M., tekhn. red.

[Socialist reorganization of agriculture in the U.S.S.R.
and the people's democracies] Sotsialisticheskoe preo-
brazovanie sel'skogo khoziaistva v SSSR i stranakh narod-
noi demokratii; uchebnyi material. Moskva, Izd-vo VPSn i
AON pri TsK KPSS, 1962. 106 p. (MIRA 16:4)
(Agriculture)

YUMASHEV, A. P.

AID P - 3676

Subject : USSR/Aeronautics
Card 1/1 Pub. 135 - 3/22
Author : Yumashev, A. P., Maj.
Title : The attack of a fighter on a bomber
Periodical : Vest. vozd. flota, 1, 12-14, Ja 1956
Abstract : The author discusses in general terms several methods of attack of fighters on a modern bomber. Some figures are given.
Institution : None
Submitted : No date

YUMASHEV, A. [P]

Subject : USSR/Aeronautics - Flying training (jets) AID P - 5549
Card 1/1 Pub. 58 - 8/20
Author : Yumashev, A., Major
Title : Special features of aerial acrobatics executed in jets
Periodical : Kryl. rod., 1, 11-13, Ja 1957
Abstract : Professional advices to pilots of jet planes as to the
carrying out of some elementary aerial acrobatics (wing
over, loop, half loops, etc.) 5 designs.
Institution : None
Submitted : No date

YUMASHEV, G.S., kand.med.nauk; GOLUBEVA, I.V., kand.med.nauk

Case of spontaneous osteolysis of the scapula. Ortop. travz.
protez, 24 no. 7:51-52 J1 '63. (MIRA 17:2)

1. Iz 2-go khirurgicheskogo otdeleniya (zav. - prof. Ya. G. Dubrov) Moskovskogo oblastnogo klinicheskogo instituta imeni M.P.Vladimirovskogo (fir. - P.M.Leonenko). Adres avtorov: Moskva I-11, ul. Shchepkina, d. 61/2, Moskovskiy oblastnoy nauchno-issledovatel'skiy klinicheskiy institut, 20-y korpus.

YUMASHEV, G. S.

"Effect of Certain Infections and Intoxications on Carcinoma in Rabbits."
Cand Med Sci, Ryazan State Medical Inst, Moscow, 1953. (RZhBiol, No 5 Nov 54)

Survey of Scientific and Technical Dissertations Defended at USSR Higher
Educational Institutions (11)

SO: Sum. No. 521, 2 Jun 55

YUMASHEV, G.S., kandidat meditsinskikh nauk; YEGOROV, G.V.

Analysis of agricultural accidents in Ostashevo District of Moscow Province during 8 years. Khirurgia no.8:55-57 Ag '54. (MLRA 7:11)

1. Iz 2-go khirurgicheskogo otdeleniya (zav. chlen-korrespondent Akademii meditsinskikh nauk SSSR prof. N.N.Blokhin) Moskovskogo oblastnogo nauchno-issledovatel'skogo klinicheskogo instituta imeni M.F.Vladimirovskogo. 2. Glavnyy khirurg Ostashevskogo rayona Moskovskoy oblasti (for Yumashev)

(AGRICULTURE,

accid. in Russia, statist.)

(ACCIDENTS,

agriculture, statist. in Russia)

YUMASHEV, G.S., kandidat meditsinskikh nauk

Distribution of the calcium isotope in tubular bones in fractures with subsequent intramedullary metal nailing; experimental investigations. Ortop.travm. i protez. no.3:3-6 My-Je '55 (MLRA 8:10)

1. Iz 2-go khirurgicheskogo otdeleniya (zav.chlen-korrespondent AMN SSSR prof. E.N.Blokhin) Moskovskogo oblastnogo nauchno-issledovatel'skogo klinicheskogo instituta im. Vladimirskogo

(FRACTURES, experimental,

intramedullary nailing, eff. on radiocalcium metab.)

(CALCIUM, radioactive,

in bones after intramedullary nailing of exper.fract.)

(BONES, metabolism,

radiocalcium, eff. of intramedullary nailing of exper.fract.)

YUMASHEV, G.S., kandidat meditsinskikh nauk

Experimental transplantation of homotransplants prepared by quick refrigeration and drying. Ortop.travm. i protez. 17 no.6:134 H-D '56.
(MLRA 10:2)

1. Iz 2-y khirurgicheskoy kliniki (zaveduyushchiy - professor Ya.G. Dubrov) Moskovskogo oblastnogo nauchno-issledovatel'skogo klinicheskogo instituta im. M.F.Vladimirovskogo (direktor - kandidat meditsinskikh nauk P.M.Leonenko)
(SKIN GRAFTING)

YUMASHEV, G. S.

YUMASHEV, G.S., kandidat meditsinskikh nauk

Some complications connected with endoprotheses. Ortop.travm. i
protez. 18 no.3:60-61 My-Je '57. (MIRA 10:9)

1. Iz 2-go khirurgicheskogo otdeleniya (zav. - prof. Ya.G.Dubov)
Moskovskogo oblastnogo nauchno-issledovatel'skogo klinicheskogo
instituta imeni M.F.Vladimirovskogo (dir. - kandidat meditsinskikh nauk
P.M.Ionenko)

(HIP, surg.
arthroplasty, compl. in use of internal prothesis)

and L. preservation

YUMASHEV, G.S.

EXCERPTA MEDICA Sec 9 Vol. 12/6 Surgery June 58

3145. (786) CLINICAL USE OF LYOPHILIZED OSSEOUS HOMOLOGOUS GRAFTS
(Russian text) - Yumashev G. S. - VESTN KHIR. 1957, 78/4 (65-68)
Tables 1 illus. 3

Bone grafts taken from cadaver and worked up by freeze drying (minus 78-196 C.) were clinically used and assessed. Twenty men and 4 women were operated upon for pseudarthroses and non-union of upper and lower extremity fractures. Lyophilized homologous grafts yielded the best results in non-union of thigh and leg long bones (a good take in 10 out of 12 cases). Pseudarthroses showed poorer results, especially when the graft failed to provide an ideal fit with the contiguous surface of the recipient bone bed.

YUMASHEV, Georgiy Stepanovich; ARONOV, Issak Grigor'yevich

[Organization of aid for the injured in villages; a manual for
physicians and feldshers] Organizatsiya travmatologicheskoi
pomoshchi na selye; kratkoe posobie dlia vrachei i fel'dsherov.
[Vladimir] Vladimirskoe knizhnoe izd-vo, 1957. 83 p. (MIRA 11:4)
(WOUNDS--TREATMENT)